# On the pre- $\lambda$ -ring structure on the Grothendieck ring of stacks and the power structures over it

## S. M. Gusein-Zade, I. Luengo and A. Melle-Hernández

To the memory of Prof. Torsten Ekedahl

# Abstract

We describe a pre- $\lambda$ -structure on the Grothendieck ring of stacks (originally studied by Torsten Ekedahl) and the corresponding power structures over it, discuss some of their properties and give some explicit formulae for the Kapranov zeta-function for some stacks. In particular, we show that the *n*th symmetric power of the class of the classifying stack BGL(1) of the group GL(1) coincides, up to a power of the class  $\mathbb{L}$  of the affine line, with the class of the classifying stack BGL(n).

# 1. Introduction

Let  $K_0(\mathcal{V}_{\mathbf{k}})$  be the Grothendieck ring of quasi-projective varieties over an algebraically closed field  $\mathbf{k}$ . This is the Abelian group generated by the classes [X] of all quasi-projective  $\mathbf{k}$ -varieties X modulo the relations:

- (1) if varieties X and Y are isomorphic, then [X] = [Y];
- (2) if Y is a Zariski closed subvariety of X, then  $[X] = [Y] + [X \setminus Y]$ .

The multiplication in  $K_0(\mathcal{V}_{\mathbf{k}})$  is defined by the Cartesian product of varieties:  $[X_1] \cdot [X_2] = [X_1 \times X_2]$ . The class  $[\mathbb{A}^1_{\mathbf{k}}] \in K_0(\mathcal{V}_{\mathbf{k}})$  of the affine line is denoted by  $\mathbb{L}$ .

In [8], there was defined a notion of a power structure over a ring and there was described a natural power structure over the Grothendieck ring  $K_0(\mathcal{V}_{\mathbb{C}})$  of complex quasi-projective varieties. This means that, for a series  $A(T) = 1 + a_1T + a_2T^2 + \ldots \in 1 + T \cdot K_0(\mathcal{V}_{\mathbb{C}})[T]$  and for an element  $m \in K_0(\mathcal{V}_{\mathbb{C}})$ , one defines a series  $(A(T))^m \in 1 + T \cdot K_0(\mathcal{V}_{\mathbb{C}})[T]$  so that all the usual properties of the exponential function hold. Here, we assume the base field **k** to be an arbitrary algebraically closed field and therefore we indicate changes required for that.

A special property of this power structure which is important for applications is its effectiveness. This means that if all the coefficients  $a_i$  of the series A(T) and the exponent m are represented by classes of quasi-projective varieties (that is,  $a_i = [A_i]$  and m = [M]), then all the coefficients of the series  $A(T)^m$  are represented by classes of quasi-projective varieties too, not by classes of virtual varieties. (This is connected with the fact that the power structure can be defined over the Grothendieck semiring of quasi-projective varieties.) The geometric description given in (1.2) has been used, in particular, in [1, 2, 9, 10]. This property looks somewhat mysterious. A power structure over a ring R can be also defined by a  $pre-\lambda$ -structure

Received 2 August 2012; published online 11 January 2013.

<sup>2010</sup> Mathematics Subject Classification 14C05 (primary), 14G10 (secondary).

The first author was partially supported by the Russian government grant 11.G34.31.0005, RFBR-10-01-00678, NSh-8462.2010.1, by Simons-IUM fellowship and by the research programme of the Universidad Complutense - Grupo Santander. The last two authors were partially supported by the grants MTM2010-21740-C02-01 and Grupo Singular CCG07-UCM/ESP-2695-921020.

on R. The pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_{\mathbb{C}})$  which induces the power structure described in [8] is defined by the Kapranov zeta-function introduced in [13]:

$$\zeta_X(T) := 1 + [S^1 X] \cdot T + [S^2 X] \cdot T^2 + [S^3 X] \cdot T^3 + \dots,$$
(1.1)

where  $S^k X := X^k / S_k$  is the *k*th symmetric power of the variety *X*. Indeed the use of another (natural) pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_{\mathbb{C}})$  (opposite to the one defined by (1.1)) induces a power structure over it which is not effective (see Statement 1 in Section 3).

For a field **k** of positive characteristic, the Kapranov zeta-function (1.1) also defines a pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_{\mathbf{k}})$  (see [6, Proposition 1.1(i)]) and therefore a power structure over it. However, in [6] it is explained that (over a field of positive characteristic) one should consider the following, a little bit different, pre- $\lambda$ -structure on  $K_0(\mathcal{V}_{\mathbf{k}})$ .

Let  $X^k = \prod_{\{k_i\}} X^k_{\{k_i\}}$  be the natural representation of the Cartesian power  $X^k$  of X as the union of the strata corresponding to all the partitions of k. The stratum corresponding to the partition  $\{k_i\}: \sum ik_i = k$  consists of the k-tuples  $(x_1, \ldots, x_k)$  with (exactly)  $k_i$  groups with (exactly) i equal elements. For the partition  $\{k_i\}$ , let  $\prod_i X^{k_i} \setminus \Delta$  be the complement of the large diagonal which consists of  $\sum_i k_i$ -tuples of points of X with at least two coinciding ones. The group  $\prod_i S_{k_i}$  acts freely on  $\prod_i X^{k_i} \setminus \Delta$ . One has  $X^k_{\{k_i\}}/S_k \cong (\prod_i X^{k_i} \setminus \Delta)/\prod_i S_{k_i}$ . One should use  $[S^k X]^* = \sum_{\{k_i\}} [(\prod_i X^{k_i}) \setminus \Delta)/\prod_i S_{k_i}]$  instead of  $[S^k X]$  in definition (1.1). For the base field  $\mathbf{k}$  of characteristic zero, the classes  $[S^k X]$  and  $[S^k X]^*$  coincide. Moreover, one can see that the geometric definition of the power structure in  $[\mathbf{8}]$ , being literally extended to a field of an arbitrary characteristic, corresponds, in fact, to this modification of the Kapranov zeta-function. Indeed in  $[\mathbf{8}]$ , the series  $(A(T))^{[X]}$  for  $A(T) = 1 + \sum_{i=1}^{\infty} [A_i] T^i$  where  $[A_i]$  are quasi-projective varieties and for a quasi-projective variety X is defined as

$$(A(T))^{[X]} := 1 + \sum_{k=1}^{\infty} \left\{ \sum_{\{k_i\}:\sum i k_i = k} \left[ \left( \left(\prod_i X^{k_i}\right) \setminus \Delta \right) \times \prod_i A_i^{k_i} \middle/ \prod_i S_{k_i} \right] \right\} \cdot T^k, \quad (1.2)$$

where the group  $S_{k_i}$  acts by permuting corresponding  $k_i$  factors in  $\prod_i X^{k_i} \supset (\prod_i X^{k_i}) \setminus \Delta$  and the spaces  $A_i$  simultaneously. If all  $A_i$  are points (that is,  $[A_i] = 1$ ), the coefficient at  $T^k$  in  $(1 + T + T^2 + \ldots)^{[X]} = (1 - T)^{-[X]}$  is just  $\sum_{\{k_i\}} [(\prod_i X^{k_i} \setminus \Delta) / \prod_i S_{k_i}] = [S^k X]^*$ . There are natural generalizations of the pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_k)$  to

There are natural generalizations of the pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_k)$  to other situations, say to the Grothendieck ring  $K_0(\operatorname{Stck}_{\mathbb{C}})$  of algebraic stacks of finite type over  $\mathbb{C}$  all of whose automorphism group schemes are affine. This pre- $\lambda$ -structure was defined in [6]. The proof there was implicit. Moreover, it contained a certain inaccuracy (the equation for the symmetric power of a product in Corollary 4.4 used in the construction) which had to be corrected. It was corrected in a new extended version of [6]. (It was sent by Prof. T. Ekedahl to the authors in a personal communication.) It is a matter of regret that this version did not appear in the arXiv and was not published (at least up to now). The pre- $\lambda$ -structure described in [6] and, in particular, Corollary 4.4. are used, for instance, in [4].

In this paper, we discuss the pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\operatorname{Stck}_k)$  of stacks and the corresponding power structure over it, describe some of their properties and give some explicit formulae for the Kapranov zeta-function for some stacks. In particular, we show that the *k*th symmetric power of the class of the classifying stack BGL(1) of the group  $\operatorname{GL}(1)$ coincides, up to a power of the class  $\mathbb{L}$  of the affine line, with the class of the classifying stack BGL(k). This leads to the statement that in a natural sense the *k*th symmetric power of the infinite-dimensional projective space  $\mathbb{CP}^{\infty}$  is piecewise isomorphic to the infinite-dimensional Grassmannian  $\operatorname{Gr}(k, \infty)$ . Though the corresponding pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\operatorname{Stck}_k)$  is effective (that is, all the coefficients of the series  $\zeta_X(T)$  are represented by effective stacks), the corresponding power structure fails to be effective (see Statement 2 in Section 3).

#### S. M. GUSEIN-ZADE ET AL.

### 2. Kapranov zeta-function for some stacks

A pre- $\lambda$ -structure on a ring R is given by a series  $\lambda_a(T) \in 1 + T \cdot R[T]$  defined for each  $a \in R$  so that

(1)  $\lambda_a(T) = 1 + aT \mod T^2;$ (2)  $\lambda_{a+b}(T) = \lambda_a(T)\lambda_b(T)$  for  $a, b \in R.$ 

defined by the Kapranov zeta-function:

A natural pre- $\lambda$ -structure on the Grothendieck ring  $K_0(\mathcal{V}_{\mathbf{k}})$  of quasi-projective varieties is

$$\zeta_X(T) := 1 + [S^1 X] \cdot T + [S^2 X] \cdot T^2 + [S^3 X] \cdot T^3 + \dots$$

(where  $S^k X := X^k/S_k$  is the *k*th symmetric power of the variety *X*) modified, for fields of positive characteristic, in the above-described way. The Kapranov zeta-function possesses the property  $\zeta_{\mathbb{L}^s X}(T) = \zeta_X(\mathbb{L}^s T)$ . In [7, Lemma 4.4, 8, Statement 3], this property was proved for  $\mathbf{k} = \mathbb{C}$ . The proof is based on Statement 2 therein: for any integer  $s \ge 0$ ,  $(A(\mathbb{L}^s T))^{[X]} = (A(T)^{[X]})|_{t \mapsto \mathbb{L}^s t}$ . One considers the corresponding parts

$$V = \left( \left( \prod_{i} X^{k_i} \right) \setminus \Delta \right) \times \prod_{i} A_i^{k_i} / \prod_{i} S_{k_i}$$

and

$$\tilde{V} = \left( \left( \prod_{i} X^{k_i} \right) \setminus \Delta \right) \times \prod_{i} (\mathbb{L}^{si} \times A_i)^{k_i} / \prod_{i} S_{k_i}$$

of the coefficients of the series  $(A(T))^{[X]}$  and  $(A(\mathbb{L}^s T))^{[X]}$ . The natural map  $\tilde{V} \to V$  is a Zariski locally trivial vector bundle of rank sk. In [8], this is deduced from [15]. For the base field **k** of positive characteristic, one can use [14, Section 7, Proposition 7].

The property  $\zeta_{\mathbb{L}^s X}(T) = \zeta_X(\mathbb{L}^s T)$  permits to 'extend' the pre- $\lambda$ -structure to the localization  $K_0(\mathcal{V}_{\mathbf{k}})[\mathbb{L}^{-1}]$  of the Grothendieck ring  $K_0(\mathcal{V}_{\mathbf{k}})$  in the multiplicative set  $\{\mathbb{L}^n\}$ . Thus, for an element  $M \in K_0(\mathcal{V}_{\mathbf{k}})[\mathbb{L}^{-1}]$  its Kapranov zeta-function is defined as  $\zeta_M(T) := \zeta_{\mathbb{L}^s M}(\mathbb{L}^{-s}T)$  for s large enough so that  $\mathbb{L}^s M \in K_0(\mathcal{V}_{\mathbf{k}})$ . In particular,

$$\zeta_{\mathbb{L}^{-s}}(T) = 1 + \mathbb{L}^{-s} \cdot T + \mathbb{L}^{-2s} \cdot T^2 + \mathbb{L}^{-3s} \cdot T^3 + \ldots = \frac{1}{1 - \mathbb{L}^{-s}T}.$$

In the Grothendieck ring  $K_0(\mathcal{V}_{\mathbf{k}})$ , one has  $[\operatorname{GL}(n)] = (\mathbb{L}^n - 1)(\mathbb{L}^n - \mathbb{L}) \dots (\mathbb{L}^n - \mathbb{L}^{n-1}).$ 

Let  $K_0(\text{Stck}_k)$  be the Grothendieck group of algebraic stacks of finite type over k all of whose automorphism group schemes are affine with relations:

- (1) [X] depends only on the isomorphism class of X;
- (2) [X] = [Y] + [U] if Y is a closed substack of X and U is its complement;
- (3) if  $E \to X$  is a vector bundle of constant rank n, then  $[E] = \mathbb{L}^n[X]$ , where  $\mathbb{L}$  also denotes the class  $[\mathbb{A}^1_{\mathbf{k}}] \in K_0(\operatorname{Stck}_{\mathbf{k}})$  of the affine line.

The multiplication in  $K_0(\text{Stck}_{\mathbf{k}})$  is defined by the fibred product of stacks:  $[X_1] \cdot [X_2] = [X_1 \times X_2]$ . The following property holds in  $K_0(\text{Stck}_{\mathbf{k}})$ ; see [5]. Let  $\mathcal{Z}_{\mathbf{k}}$  be the class of connected group schemes for which torsors (principal *G*-bundles in the étale topology) are trivial in the Zariski topology (for example, GL(n) and SL(n) are in  $\mathcal{Z}_{\mathbf{k}}$ ). For a group  $G \in \mathcal{Z}_{\mathbf{k}}$ , *G* is affine and connected. Then, for any group  $G \in \mathcal{Z}_{\mathbf{k}}$ , [G][BG] = 1 where *BG* is the classifying stack of the group *G* (for example,  $[BGL(1)] = 1/(\mathbb{L} - 1)$ ). Thus, in particular,  $\mathbb{L}$  and  $\mathbb{L}^n - 1$  for any  $n \ge 1$  are invertible in  $K_0(\text{Stck}_{\mathbf{k}})$ .

There is a natural ring homomorphism from  $K_0(\mathcal{V}_k)$  to  $K_0(\text{Stck}_k)$  by considering an algebraic variety as an algebraic stack. In [5, Theorem 1.2] (see also [3, 16]), it was shown that the natural ring homomorphism

$$K_0(\mathcal{V}_{\mathbf{k}})[\mathbb{L}^{-1}][(\mathbb{L}^n - 1)^{-1}, \forall n \ge 1] \longrightarrow K_0(\operatorname{Stck}_{\mathbf{k}})$$
(2.1)

is an isomorphism between the Grothendieck ring  $K_0(\text{Stck}_k)$  of algebraic stacks with affine stabilizers and the Grothendieck ring  $K_0(\mathcal{V}_k)$  of varieties localized by the elements  $\mathbb{L} := [\mathbb{A}^1_k]$ and  $\mathbb{L}^n - 1$  for all  $n \ge 1$ . (The fact that these rings tensored by the field  $\mathbb{Q}$  of rational numbers are isomorphic was proved in [12].)

Here, we shall give somewhat more explicit formulae for the pre- $\lambda$ -structure defined in [6] (that is, for the corresponding generalization of the Kapranov zeta-function) and compute it for some examples.

Let  $R_k(q_1,\ldots,q_k)$  be the rational function in  $q_1,\ldots,q_k$  defined by its Taylor expansion

$$R_k(q_1, \dots, q_k) := \sum_{\underline{i} = (i_1, \dots, i_k) \in \mathbb{Z}_{\geq 0}^k: \ i_s \neq i_r \ \text{for } s \neq r} q_1^{i_1} q_2^{i_2} \dots q_k^{i_k}.$$
 (2.2)

FACT 1. One can see that

$$R_k(q_1, \dots, q_k) = \sum_{\sigma \in S_k} \frac{q_{\sigma_1}^{k-1} q_{\sigma_2}^{k-2} \dots q_{\sigma_{k-1}}}{(1 - q_{\sigma_1})(1 - q_{\sigma_1} q_{\sigma_2}) \dots (1 - q_{\sigma_1} \dots q_{\sigma_k})}$$

Here  $q_1^{k-1}q_2^{k-2} \dots q_{k-1}/(1-q_1)(1-q_1q_2) \dots (1-q_1 \dots q_k)$  is the part of the sum in (2.2) over  $\underline{i} = (i_1, \dots, i_k)$  with  $i_1 > i_2 > \dots > i_k \ge 0$ .

FACT 2. For  $\sum_{j=1}^{s} n_j = k$ , define

$$R_{n_1,n_2,\ldots,n_s}(q_1,\ldots,q_s) := \sum_{\{i_r^{(j)}\}} q_1^{i_1^{(1)}} \ldots q_1^{i_{n_1}^{(1)}} q_2^{i_1^{(2)}} \ldots q_2^{i_{n_2}^{(2)}} \ldots q_s^{i_1^{(s)}} \ldots q_s^{i_{n_s}^{(s)}},$$

where the sum is over all collections of  $\{i_r^{(j)}\}$  with  $1 \le j \le s, 1 \le r \le n_j, i_1^{(j)} < i_2^{(j)} < \ldots < i_{n_j}^{(j)}, i_{r_1}^{(j_1)} \ne i_{r_2}^{(j_2)}$  for  $(j_1, r_1) \ne (j_2, r_2)$ . One can see that

$$R_{n_1,n_2,...,n_s}(q_1,...,q_s) = R_k(\underbrace{q_1,...,q_1}_{n_1},\underbrace{q_2,...,q_2}_{n_2},...,\underbrace{q_s,...,q_s}_{n_s}) / \prod_{j=1}^s n_j!.$$

Because of the isomorphism (2.1), an element of the Grothendieck ring  $K_0(\text{Stck}_{\mathbf{k}})$  of stacks is of the form  $a = M\mathbb{L}^{-m}/\prod_{i=1}^{\ell}(1-\mathbb{L}^{-n_i})$ , where  $M \in K_0(\mathcal{V}_{\mathbf{k}})$ ,  $m \in \mathbb{Z}$ ,  $n_i \in \mathbb{Z} \setminus \{0\}$ . (Without loss of generality one may assume that  $m \in \mathbb{Z}_{\geq 0}$ ,  $n_i \in \mathbb{Z}_{>0}$ ; however, for the formulae below this is not essential.)

In [6, Lemma 2.2], the corresponding series  $\zeta_a(T)$ , or rather  $(\zeta_a(-T))^{-1} (= \zeta_{-a}(-T))$ , was described by a functional equation. A closed formula for  $\zeta_a(T)$  is somewhat involved. Therefore, we shall give a formula for the Kapranov zeta-function of an element a of the form  $b \mathbb{L}^{-m}/(1 - \mathbb{L}^{-n})$ , where  $b \in K_0(\text{Stck}_k)$  (in terms of the series  $\zeta_b(T)$ ). Let us denote the coefficients of the series  $\zeta_b(T)$  by  $\sigma^k b \in K_0(\text{Stck}_k)$ :

$$\zeta_b(T) = 1 + \sum_{k=1}^{\infty} \sigma^k b \cdot T^k,$$

 $\sigma^1 b = b$ . For b = [X], where X is a quasi-projective variety,  $\sigma^k b$  is the class  $[S^k X]$  of its symmetric power  $S^k X$  (modified, for a field **k** of positive characteristic, in the described way).

REMARK 1. Note that, for  $\sum_{j=1}^{s} n_j = k$  and for arbitrary  $\ell_1, \ldots, \ell_s$ , the corresponding element  $R_{n_1, n_2, \ldots, n_s}((\mathbb{L}^{-1})^{\ell_1}, \ldots, (\mathbb{L}^{-1})^{\ell_s}) \in K_0(\operatorname{Stck}_{\mathbf{k}})$  is effective, in the sense that it is equal to the class of a quasi-projective variety divided by  $\mathbb{L}^r \prod_i [\operatorname{GL}(r_i)]$ . This follows directly from the formula for  $R_k(q_1, \ldots, q_k)$ .

PROPOSITION 1. For an element  $a = bq^m/(1-q^n)$ , where  $b \in K_0(\text{Stck}_k)$  (and  $q = \mathbb{L}^{-1}$ ), one has

$$\zeta_a(T) = 1 + \sum_{k=1}^{\infty} \left( \sum_{\underline{k}:\sum_{j=1}^s jk_j = k} R_{k_1, 2k_2, \dots, sk_s}(q^n, q^{2n}, \dots, q^{sn}) \prod_j (\sigma^j b)^{k_j} \right) q^{km} T^k,$$

where the second sum runs over all partitions  $\underline{k} = (k_1, \ldots, k_s) : \sum_{j=1}^s jk_j = k$  of the integer k.

COROLLARY 1. For an element  $a = 1/(1 - q^n) \in K_0(\operatorname{Stck}_{\mathbf{k}})$  (and  $q = \mathbb{L}^{-1}$ ), one has

$$\zeta_{1/(1-q^n)}(T) = 1 + \sum_{k=1}^{\infty} \left( \sum_{\underline{k}:\sum_{j=1}^s jk_j = k} R_{k_1, 2k_2, \dots, sk_s}(q^n, q^{2n}, \dots, q^{sn}) \right) T^k,$$

where the second sum runs over all partitions  $\underline{k} = (k_1, \ldots, k_s) : \sum_{j=1}^s jk_j = k$  of k.

Proof. According to [6], the series  $\zeta_a(T)$  is defined by the following functional equation  $\zeta_a(T)/\zeta_a(q^n T) = \zeta_b(q^m T)$  and the coefficients of this series are polynomials in  $\sigma^i b$ ,  $q^{\pm 1}$  and  $(1-q^j)^{-1}$  for all j > 0. Polynomials in  $q^{\pm 1}$  and  $(1-q^j)^{-1}$  can be defined by their Laurent expansion at q = 0. Consider the series in T, whose coefficients are elements of  $\mathbb{Z}[q^{\pm 1}, \sigma^i b][\![q]\!]$ , of the form

$$\psi(T) := \prod_{i=0}^{\infty} \zeta_b(q^{m+in}T)$$

One can see that

$$\psi(T)/\psi(q^n T) = \prod_{i=0}^{\infty} \zeta_b(q^{m+in} T) \left(\prod_{i=0}^{\infty} \zeta_b(q^{m+(i+1)n} T)\right)^{-1} = \zeta_b(q^m T).$$

Also one can show (see below) that the coefficients of the series  $\psi(T)$  are Laurent expansions in q of polynomials in  $\sigma^k b$ ,  $q^{\pm 1}$  and  $(1 - q^j)^{-1}$ . Therefore,  $\psi(T) = \zeta_a(T)$ .

Thus, one has

$$\zeta_a(T) = \prod_{i=1}^{\infty} (1 + q^{m+in} \sigma^1 b \ T + q^{2(m+in)} \sigma^2 b \ T^2 + \ldots).$$

The coefficient of  $T^k$  in this series is the sum over all partitions  $\underline{k} = (k_1, \ldots, k_s)$  of k with  $\sum_{j=1}^s jk_j = k$  of the series

$$\prod_{j} (\sigma^{j} b)^{k_{j}} \sum_{\substack{\{i_{r}^{(j)}\}\\ \{i_{r}^{(j)}\}}} q^{m+i_{1}^{(1)}n} \dots q^{m+i_{k_{1}}^{(1)}n} q^{m+2i_{1}^{(2)}n} \dots q^{m+2i_{k_{2}}^{(2)}n} \dots q^{m+si_{1}^{(2)}n} \dots q^{m+si_{k_{s}}^{(s)}n},$$

where the sum is over all collections of  $\{i_r^{(j)}\}$  with  $1 \leq j \leq s, 1 \leq r \leq n_j, i_1^{(j)} < i_2^{(j)} < \ldots < i_{n_j}^{(j)}, i_{r_1}^{(j_1)} \neq i_{r_2}^{(j_2)}$  for  $(j_1, r_1) \neq (j_2, r_2)$ . This implies the statement.

COROLLARY 2. The Kapranov zeta-function for stacks is effective in the sense that, for a stack X, the coefficients of  $\zeta_{[X]}(T)$  are represented by classes of stacks in  $K_0(\text{Stck}_k)$ .

This was shown by Ekedahl [6] who identified the coefficients of  $\zeta_{[X]}(T)$  with the classes of the stack symmetric products of X.

EXAMPLE 1. The Kapranov zeta-function of the element  $a = q^m/(1-q^n) \in K_0(\text{Stck}_k)$  is equal to

$$\zeta_{q^m/(1-q^n)}(T) = 1 + \sum_{k=1}^{\infty} \frac{q^{mk}}{(1-q^n)(1-q^{2n})\dots(1-q^{kn})} T^k.$$
(2.3)

In particular,

$$\zeta_{[BGL(1)]}(T) = \zeta_{1/(\mathbb{L}-1)}(T) = 1 + \sum_{k=1}^{\infty} \frac{\mathbb{L}^{k^2 - k}}{(\mathbb{L}^k - \mathbb{L}^{k-1}) \dots (\mathbb{L}^k - 1)} T^k.$$
(2.4)

Thus,

$$\sigma^{k}[BGL(1)] = \mathbb{L}^{k^{2}-k}[BGL(k)].$$
(2.5)

As in the proof of the proposition with b = 1 consider

$$\psi(T) = 1 + \sum_{k=1}^{\infty} \frac{q^{mk}}{(1-q^n)(1-q^{2n})\dots(1-q^{kn})} T^k,$$

and we have to show that the following functional equation  $\psi(T)/\psi(q^n T) = \zeta_1(q^m T)$  holds. Thus, equation (2.3) follows from the equalities

$$\left( 1 + \sum_{k=1}^{\infty} \frac{q^{(m+n)k}}{(1-q^n)(1-q^{2n})\dots(1-q^{kn})} T^k \right) \left( \sum_{k=0}^{\infty} q^{mk} T^k \right)$$

$$= 1 + \sum_{k=1}^{\infty} \sum_{j=0}^k \frac{q^{(m+n)j+m(k-j)}}{(1-q^n)(1-q^{2n})\dots(1-q^{jn})} T^k$$

$$= 1 + \sum_{k=1}^{\infty} \sum_{j=0}^k \frac{q^{nj}}{(1-q^n)(1-q^{2n})\dots(1-q^{jn})} q^{mk} T^k$$

$$= 1 + \sum_{k=1}^{\infty} \frac{q^{mk}}{(1-q^n)(1-q^{2n})\dots(1-q^{kn})} T^k.$$

REMARK 2. Note that the classifying space for the group  $\operatorname{GL}(1) = \mathbb{C}^*$  in the topological sense is  $\mathbb{CP}^{\infty}$ , whence the topological classifying space for the group  $\operatorname{GL}(k)$  is the (infinite-dimensional) Grassmannian  $\operatorname{Gr}(k, \infty)$ . If one defines  $\zeta_{\mathbb{CP}^{\infty}}(T)$  as

$$\zeta_{\mathbb{CP}^{\infty}}(T) := \lim_{N \to \infty} \zeta_{\mathbb{CP}^{N}}(T) = \prod_{j=0}^{\infty} \zeta_{\mathbb{L}^{j}}(T)$$

(whatever this means) and  $[\mathbf{Gr}(k,\infty)]$  as a series in  $\mathbb{L}$  equal to  $[\mathbf{Gr}(k,\infty)] = \lim_{N\to\infty} [\mathbf{Gr}(k,N)]$ , then one gets

$$\zeta_{\mathbb{CP}^{\infty}}(T) = 1 + \sum_{k=1}^{\infty} [\mathbf{Gr}(k, \infty)] T^k, \qquad (2.6)$$

which is similar to (2.4) up to dimensional factors  $\mathbb{L}$  in some powers in the coefficients. Equation (2.6) means that  $[S^k \mathbb{CP}^\infty] = [\mathbf{Gr}(k, \infty)]$ . Moreover,  $S^k \mathbb{CP}^\infty$  and  $\mathbf{Gr}(k, \infty)$  have decompositions into quasi-projective varieties (compatible with the inclusions  $S^k \mathbb{CP}^N \subset S^k \mathbb{CP}^\infty$  and  $\mathbf{Gr}(k, N) \subset \mathbf{Gr}(k, \infty)$ ) such that components of  $S^k \mathbb{CP}^\infty$  and of  $\mathbf{Gr}(k, \infty)$  are pairwise isomorphic. This uses the fact that  $S^k \mathbb{C}^n$  and  $\mathbb{C}^{kn}$  have decompositions into quasi-projective varieties such that their components are pairwise isomorphic. See a proof of these statements in [11].

#### 3. A power structure over $K_0(\text{Stck}_k)$

DEFINITION 1. A power structure over a (semi)ring R with a unit is a map  $(1 + T \cdot R[T]) \times R \to 1 + T \cdot R[T]$ :  $(A(T), m) \mapsto (A(T))^m$ , which possesses the following properties:

- (1)  $(A(T))^0 = 1;$ (2)  $(A(T))^1 = A(T);$ (3)  $(A(T) \cdot B(T))^m = (A(T))^m \cdot (B(T))^m;$ (4)  $(A(T))^{m+n} = (A(T))^m \cdot (A(T))^n;$ (5)  $(A(T))^{mn} = ((A(T))^n)^m;$ (6)  $(1+T)^m = 1 + mT$  + terms of higher degree;
- (7)  $(A(T^k))^m = (A(T))^m|_{T \mapsto T^k}.$

REMARK 3. Note that one does not require, in general, that  $(A(-T))^m = (A(T))^m|_{T \mapsto -T}$ : see the remark after Statement 1 below.

As it was explained in [9], a pre- $\lambda$ -structure  $\lambda_a(T) \in 1 + T \cdot R[T]$  on a ring R defines a power structure over it. To define the series  $(A(T))^m$  for  $A(T) = 1 + a_1T + a_2T^2 + \ldots, a_i \in R, m \in R$ , one has to represent (in a unique way) A(T) as a product of the form  $A(T) = \prod_{k=1}^{\infty} \lambda_{b_k}(T^k)$ , with  $b_i \in R$ , and then

$$(A(T))^m := \prod_{k=1}^{\infty} \lambda_{mb_k}(T^k).$$
(3.1)

To each pre- $\lambda$ -structure  $\lambda_a(T)$  on a ring R, there corresponds the so-called *opposite* pre- $\lambda$ -structure  $\lambda_a^{\text{op}}(T) := (\lambda_a(-T))^{-1}$ . Note that the power structures defined by these two pre- $\lambda$ -structures are in general different. One can easily see this fact for the standard pre- $\lambda$ -structure on the polynomial ring  $\mathbb{Z}[u_1, \ldots, u_k]$  described, for example, in [9].

One can say that there are at least four 'natural' pre- $\lambda$ -structures over the Grothendieck ring  $K_0(\mathcal{V}_k)$  of quasi-projective varieties. They are those defined by the Kapranov zeta-function

$$\zeta_M(T) := 1 + [S^1 M] \cdot T + [S^2 X] \cdot T^2 + [S^3 M] \cdot T^3 + \dots,$$

and by the generating series of the configuration space  $M_k = (M^k \setminus \Delta)/S_k$  of unordered k-tuples of different points of a quasi-projective variety M,

$$\varphi_M(T) := 1 + [M] \cdot T + [M_2] \cdot T^2 + [M_3] \cdot T^3 + \dots,$$

and their corresponding opposites.

One can see that the power structures corresponding to the first two pre- $\lambda$ -structures coincide (this is a consequence of the equation  $\varphi_M(T) = (1+T)^M$ ; see example in [8], where in this equality the power structure is defined by  $\zeta_M(T)$ ), whence those corresponding to their opposites are different from this one (and coincide with each other). The advantage of the power structure defined by the Kapranov zeta-function (or by the series  $\varphi_M(T)$ ) is the fact that it is defined over the Grothendieck semiring of quasi-projective varieties (whose elements are represented by 'genuine' varieties and not by virtual ones). We shall say that this power structure is effective. This follows from the above-mentioned geometric description of the coefficient of the series  $(A(T))^M$  for  $A(T) = 1 + [A_1] \cdot T + [A_2] \cdot T^2 + [A_3] \cdot T^3 + \ldots$ , with  $A_k$  and M being quasi-projective varieties. We do not know how one can prove the effectiveness of this power structure from the definition via  $\varphi_M(T)$ . With their opposite power structure, one has the following statement.

STATEMENT 1. The power structure over the Grothendieck ring  $K_0(\mathcal{V}_{\mathbf{k}})$  defined by the pre- $\lambda$ -structure  $\lambda_X(T) := (\zeta_X(-T))^{-1}$  is not effective.

For a class [M] of a quasi-projective variety M,  $\lambda_{[M]}(T)$  (equal to  $(\lambda_1(T))^{[M]} = (1+T)^{[M]}$ in the power structure defined by the pre- $\lambda$ -ring structure  $\lambda_X(T)$ ) is the inverse of the series  $(1 - [M]T + [S^2M]T^2 - [S^3M]T + ...)$ , that is,  $\lambda_{[M]}(T) = 1 + [M]T + ([M]^2 - [S^2M])$  $T^2 + ...$  If M is a non-singular elliptic curve C, then the Hodge–Deligne polynomial  $e([C]^2 - [S^2C])$  is equal to  $-u^2v - uv^2 + u^2 + v^2 + 2uv - u - v$ , since e([C]) = 1 - u - v + uvand  $e([S^2C])$  is the coefficient at  $T^2$  in the series (1/(1-T))(1-uT)(1-vT)(1/(1-uvT)). Such polynomial cannot be the Hodge–Deligne polynomial of a quasi-projective variety since its homogeneous part of highest degree is not of the form  $\ell(uv)^n$  with a non-negative integer  $\ell$ .

**REMARK 4.** The computations above show that

$$(1+T)^{[M]} = 1 + [M] \cdot T + [M_2] \cdot T^2 + \dots$$

is not equal, in general, to

$$(1-T)^{[M]}|_{T\mapsto -T} = 1 + [M]T + ([M]^2 - [S^2M])T^2 + \dots$$

STATEMENT 2. The power structure over the Grothendieck ring  $K_0(\text{Stck}_k)$  defined by the Kapranov zeta-function is not effective.

We shall show that the series  $(1+T)^{[BGL(1)]} = (1+T)^{1/(\mathbb{L}-1)}$  is not effective, in the sense that it contains some coefficients which are not represented by varieties. Namely, we will show that the second coefficient of such a series is not effective. One has

$$(1+T)^{1/(\mathbb{L}-1)} = \frac{(1-T)^{-1/(\mathbb{L}-1)}}{(1-T^2)^{-1/(\mathbb{L}-1)}} = \frac{\zeta_{1/(\mathbb{L}-1)}(T)}{\zeta_{1/(\mathbb{L}-1)}(T^2)}.$$

According to (2.4)

$$(1+T)^{1/(\mathbb{L}-1)} \bmod T^3 = \left(1 + \frac{1}{\mathbb{L}-1}T + \frac{\mathbb{L}^2}{(\mathbb{L}^2 - 1)(\mathbb{L}^2 - \mathbb{L})}T^2\right) \left(1 - \frac{1}{\mathbb{L}-1}T^2\right) \bmod T^3.$$

The coefficient at  $T^2$  is equal to

$$\frac{\mathbb{L}^2}{(\mathbb{L}^2 - 1)(\mathbb{L}^2 - \mathbb{L})} - \frac{1}{\mathbb{L} - 1} = \frac{-\mathbb{L}^3 + \mathbb{L}^2 + \mathbb{L}}{[\operatorname{GL}(2)]}$$

It cannot be represented by a variety since the term of highest degree in the Hodge–Deligne polynomial of the numerator has a negative coefficient.

For a quasi-projective variety X, the coefficient at  $T^n$  in the series  $(1+T)^{[X]}$  is represented by the configuration space  $(X^n \setminus \Delta)/S^n$  of n different points in X. Statement 2 gives a hint for the conjecture that the notion of the configuration spaces of different points cannot be defined for stacks (at least in a form close to that for varieties).

#### References

 K. BEHREND, J. BRYAN and B. SZENDRÖI, 'Motivic degree zero Donaldson-Thomas invariants', Invent. Math. (2012) doi:10.1007/s00222-012-0408-1.

- D. BOURQUI, 'Produit eulérien motivique et courbes rationnelles sur les variétés toriques', Compos. Math. 145 (2009) 1360–1400.
- 3. T. BRIDGELANDL, 'An introduction to motivic Hall algebras', Adv. Math. 229 (2012) 102-138.
- 4. B. DAVISON and S. MEINHARDT, 'Motivic DT-invariants for the one loop quiver with potential', Preprint, 2011, arXiv:1108.5956.
- 5. T. EKEDAHL, 'The Grothendieck group of algebraic stacks', Preprint, 2009, arXiv:0903.3143.
- 6. T. EKEDAHL, 'A geometric invariant of a finite group', Preprint, 2009, arXiv:0903.3148v1.
- 7. L. GÖTTSCHE, 'On the motive of the Hilbert scheme of points on a surface', Math. Res. Lett. 8 (2001) 613–627.
- S. M. GUSEIN-ZADE, I. LUENGO and A. MELLE-HERNÁNDEZ, 'A power structure over the Grothendieck ring of varieties', Math. Res. Lett. 11 (2004) 49–57.
- S. M. GUSEIN-ZADE, I. LUENGO and A. MELLE-HERNÁNDEZ, 'Power structure over the Grothendieck ring of varieties and generating series of Hilbert schemes of points', Michigan Math. J. 54 (2006) 353–359.
- S. M. GUSEIN-ZADE, I. LUENGO and A. MELLE-HERNÁNDEZ, 'On the power structure over the Grothendieck ring of varieties and its applications', Proc. Steklov Inst. Math. 258 (2007) 53–64.
- 11. S. M. GUSEIN-ZADE, I. LUENGO and A. MELLE-HERNÁNDEZ, 'On piecewise isomorphism of some varieties', Azerb. J. Math. 1 (2011) 91–96.
- 12. D. JOYCE, 'Motivic invariants of Artin stacks and "stack functions", Q. J. Math. 58 (2007) 345-392.
- M. KAPRANOV, 'The elliptic curve in the S-duality theory and Eisenstein series for Kac-Moody groups'. Preprint, 2000, arXiv math.AG/0001005.
- D. MUMFORD, Abelian varieties, Tata Institute of Fundamental Research Studies in Mathematics 5, Bombay (Oxford University Press, London, 1970) viii+242 pp.
- 15. J. P. SERRE 'Les espaces fibrés algébriques', Anneaux de Chow et applications, Séminaire Claude Chevalley, 2e année, tome 3 (1958) Exposé 1, 1–37.
- 16. B. TÖEN, 'Anneaux de Grothendieck des n-champs d'Artin', Preprint, 2005, arXiv:math/0509098.

S. M. Gusein-Zade Faculty of Mathematics and Mechanics Moscow State University Moscow GSP-1, 119991 Russia

sabir@mccme.ru

A. Melle-Hernández Faculty of Mathematics University Complutense of Madrid Madrid 28040 Spain

Current address: ICMAT (CSIC-UAM-UC3M-UCM) C/ Nicolás Cabrera, 13-15 Campus de Cantoblanco, UAM Madrid 28049 Spain

amelle@mat.ucm.es

I. Luengo Faculty of Mathematics University Complutense of Madrid Madrid 28040 Spain

iluengo@mat.ucm.es