EQUIVARIANT VERSIONS OF HIGHER ORDER ORBIFOLD EULER CHARACTERISTICS

S.M. GUSEIN-ZADE, I. LUENGO, AND A. MELLE-HERNÁNDEZ

ABSTRACT. There are (at least) two different approaches to define an equivariant analogue of the Euler characteristic for a space with a finite group action. The first one defines it as an element of the Burnside ring of the group. The second approach emerged from physics and includes the orbifold Euler characteristic and its higher order versions. Here we give a way to merge the two approaches together defining (in a certain setting) higher order Euler characteristics with values in the Burnside ring of a group. We give Macdonald type equations for these invariants. We also offer generalized ("motivic") versions of these invariants and formulate Macdonald type equations for them as well.

2010 Math. Subj. Class. 55M35, 32Q55, 19A22.

KEY WORDS AND PHRASES. Finite group actions, orbifold Euler characteristic, Burnside ring, complex quasi-projective varieties, wreath products, generating series.

1. INTRODUCTION

Let X be a topological space (good enough, say, a real subanalytic variety) with an action of a finite group G. There are (at least) two different approaches to define an equivariant analogue of the Euler characteristic for the pair (X, G). The first one [29] defines the equivariant Euler characteristic $\chi^G(X)$ as an element of the Burnside ring A(G) of the group G. The second approach emerged from physics (the string theory of orbifolds: [10], [11]). The orbifold Euler characteristic $\chi^{orb}(X, G)$ is defined through the fixed point sets of some subgroups of G and is an integer. Higher order (orbifold) Euler characteristics were introduced in [1] and [8] (also as integers). They can be defined through the fixed point sets of collections of commuting elements in G. Here we give a way to merge the two approaches together.

Through this paper we consider the Euler characteristic $\chi(\cdot)$ defined as the alternating sum of the dimensions of the cohomology groups with compact support.

 $\odot 2016$ Independent University of Moscow

Received June 5, 2016; in revised form June 31, 2016.

The work of the first named author (Sections 1, 2 and 4) was supported by the grant 16-11-10018 of the Russian Science Foundation. The second and third mentioned authors were supprted in part by the grant MTM2013-45710-C02-02-P.

This Euler characteristic is not a homotopy invariant in the usual sense, but an invariant of the homotopy type defined in terms of proper maps. It is an additive function on the algebra of ("good") spaces (say, on the algebra of (real) constructible sets.). For complex quasi-projective varieties this Euler characteristic coincides with the "usual" one, i.e., with the alternating sum of the dimensions of the usual cohomology groups.

There is the universal additive invariant on the algebra of complex constructible sets. It takes values in the Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$ of complex quasi-projective varieties and can be regarded as a generalized ("motivic") Euler characteristic $\chi_g(X)$. There were defined generalized ("motivic") analogues of the orbifold Euler characteristic and of its higher order generalizations. First it was essentially made in [2] (for the Hodge–Deligne polynomial) and then formulated precisely in [17] and [19]. These higher order generalized Euler characteristics take values in a certain modification of the Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$ of complex quasi-projective varieties.

The Euler characteristic satisfies the Macdonald type equation

$$1 + \sum_{k=1}^{\infty} \chi(S^k X) \cdot t^k = (1-t)^{-\chi(X)}, \tag{1}$$

where $S^k X = X^k/S_k$ is the *k*th symmetric power of *X* (see, e.g., [24]). A Macdonald type equation for a given invariant expresses the generating series of the values of the invariant for the symmetric powers of a space (or for their analogues) as a series not depending on the space (in this case $(1 - t)^{-1}$) with the exponent equal to the value of the invariant for the space itself. If the invariant takes values in a ring *R* different from a number ring (i.e., from \mathbb{Z} , \mathbb{Q} , \mathbb{R} or \mathbb{C}), Macdonald type equations can be formulated in terms of a so-called power structure over the ring. There are Macdonald type equations for the orbifold Euler characteristic and for its higher order analogues ([30], [28]) and also for the equivariant Euler characteristic with values in the Burnside ring A(G) of *G* (see Lemma 1 below). An analogue of these equations for the generalized ("motivic") higher order Euler characteristics was obtained in [19]. It was formulated in terms of the (natural) power structure over the Grothendieck ring $K_0(\text{Var}_{\mathbb{C}})$ of complex quasi-projective varieties: [15]. (Its version for the orbifold Hodge–Deligne polynomial was proved in [31].)

Here we define higher order Euler characteristics with values in the Burnside ring of a group. This is made in the setting when there are two commuting finite group actions. Namely, let a topological space X be endowed with commuting actions of two finite groups G_O and G_B (or equivalently with an action of the direct product $G_O \times G_B$). Then the action of G_O can be treated in the way similar to "the orbifold approach" and the action of G_B in the way which leads to invariants (say, equivariant Euler characteristic) with values in the Burnside ring of the group. This gives a possibility to define higher order Euler characteristics of the triple $(X; G_O, G_B)$ with values in the Burnside ring of the group G_B .

This situation can be met, for example, in the following considerations. Assume that (f, G) is a pair consisting of a quasi-homogeneous polynomial f and an abelian

group G of its (diagonal) symmetries. Such pairs are subjects of analysis, in particular, in the Berglund-Hübsch-Henningson mirror symmetry [4], [3]. The classical monodromy transformation of f is a map of a finite order from the Milnor fibre of f into itself. It commutes with the G-action. Thus one has the action of two groups: the group G and the (cyclic) group generated by the monodromy transformation. A study of the orbifold monodromy zeta function of f leads to the situation when one should apply "the orbifold approach" to the first action whence the other one should be treated in another way, see, e.g., [12].

We give Macdonald type equations for the constructed invariants (Theorem 1). We also offer generalized ("motivic") versions of these invariants and formulate Macdonald type equations for them as well (Theorem 2).

The authors are very grateful to E. A. Gorsky for carefully reading the paper and making a number of useful suggestions.

2. Power Structures over Rings

As it was indicated above, a Macdonald type equation for an invariant can be formulated in terms of a power structure over the ring of values of the invariant. Let R be a commutative associative ring with unity. A power structure over the ring R gives sense to expressions of the form $(A(t))^m$, where $A(t) = 1 + a_1t + a_2t^2 + \cdots$ is a power series with the coefficients a_i from R and m is an element of R.

Definition [15]. A power structure over the ring R is a map

 $(1 + tR[[t]]) \times R \to 1 + tR[[t]] \qquad ((A(t), m) \mapsto (A(t))^m)$

possessing the following properties:

- (1) $(1 + a_1t + \cdots)^m = 1 + ma_1t + \cdots;$ (2) $(A(t)B(t))^m = (A(t))^m (B(t))^m;$ (3) $(A(t))^{m+n} = (A(t))^m (A(t))^n;$ (4) $(A(t))^{mn} = (A(t))^m (A(t))^n;$
- (4) $(A(t))^{mn} = ((A(t))^m)^n$.

Let \mathfrak{m} be the ideal tR[[t]] in the ring R[[t]].

Definition. A power structure over the ring R is *finitely determined* if, for any $k \ge 1$, the fact that $A(t) \in 1 + \mathfrak{m}^k$ implies that $(A(t))^m \in 1 + \mathfrak{m}^k$.

The natural power structure over the ring \mathbb{Z} of integers is defined by the standard formula for a power of a series (see, e.g., [27]):

$$(1 + a_1 t + a_2 t^2 + \dots)^m =$$

= $1 + \sum_{k=1}^{\infty} \left(\sum_{\{k_i\}:\sum i k_i = k} \frac{m(m-1)\cdots(m-\sum_i k_i + 1)\cdot\prod_i a_i^{k_i}}{\prod_i k_i!} \right) \cdot t^k.$

An important example of a power structure over the Grothendieck ring of quasiprojective varieties was introduced in [15].

A quasi-projective variety is the complement of a projective variety in a projective one. The Grothendieck ring $K_0(\text{Var}_{\mathbb{C}})$ of complex quasi-projective varieties is the abelian group generated by the classes of complex quasi-projective varieties modulo the relations:

S.M. GUSEIN-ZADE, I. LUENGO, AND A. MELLE-HERNÁNDEZ

- (1) if X and X' are isomorphic, then [X] = [X'];
- (2) if $Y \subset X$ is a Zariski closed subset, then $[X] = [Y] + [X \setminus Y]$.

The multiplication in $K_0(\text{Var}_{\mathbb{C}})$ is defined by the cartesian product. The class $\mathbb{L} = [\mathbb{A}^1_{\mathbb{C}}]$ of the complex affine line plays a special role in a number of constructions connected with the Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$.

A power structure over the ring $K_0(\text{Var}_{\mathbb{C}})$ is defined in [15] by the formula

$$(1 + [A_1]t + [A_2]t^2 + \cdots)^{[M]} =$$

$$= 1 + \sum_{k=1}^{\infty} \left(\sum_{\{k_i\}:\sum ik_i = k} \left[\left(\left(M^{\sum_i k_i} \setminus \Delta \right) \times \prod_i A_i^{k_i} \right) / \prod_i S_{k_i} \right] \right) \cdot t^k, \quad (2)$$

where A_i , i = 1, 2, ..., and M are quasi-projective varieties ($[A_i]$ and [M] are their classes in the ring $K_0(\text{Var}_{\mathbb{C}}))$, Δ is the large diagonal in $M^{\sum_i k_i}$, i.e., the set of (ordered) $(\sum_i k_i)$ -tuples of points of M with at least two coinciding ones, the group S_{k_i} of permutations on k_i elements acts by the simultaneous permutations on the components of the corresponding factor M^{k_i} in $M^{\sum_i k_i} = \prod_i M^{k_i}$ and on the components of $A_i^{k_i}$.

One can see that the coefficient at t^k in the right hand side of (2) has the following interpretation. Let $I: \prod_{i=1}^{\infty} A_i \to \mathbb{Z}$ be the "tautological function" on the disjoint union $\prod_{i=1}^{\infty} A_i$ sending A_i to i. The coefficient at t^k in the right hand side of (2) is represented by the configuration space of pairs (K, ψ) , where K is a check the reference finite subset of M and ψ is a map $K \to \prod_{i=1}^{\infty} A_i$ such that $\sum_{x \in K} I(\psi(x)) = k$. This interpretation makes it much easier to prove that the equation (2) really defines check the reference a power structure (i.e., to verify the properties 2-4-from the definition) and also permits to prove some formulae for generating series of classes of Hilbert schemes of "fat points" (zero-dimensional subschemes) on quasi-projective manifolds (see, e.g., [16]). Less formally, see [14], [9], one can say that on the variety M there live particles equipped with some natural numbers (multiplicities, masses, charges, ...). A particle of multiplicity n has a complicated space of internal states which is parametrized by points of a quasi-projective variety A_n and the coefficient at t^k is the configuration space of tuples of particles with the total multiplicity k.

The notion of a power structure over a ring is closely related with the notion of a λ -ring structure. A λ -ring structure (or a pre- λ -ring structure in a certain terminology, see, e.g., [23]) is an additive-to-multiplicative homomorphism $R \rightarrow$ $1 + tR[[t]], a \mapsto \lambda_a(t) \ (\lambda_{a+b}(t) = \lambda_a(t) \cdot \lambda_b(t))$ such that $\lambda_a(t) = 1 + at + \cdots$. A $\lambda_{a+b}(t) = \lambda_a(t) \cdot \lambda_b(t)$ ring structure over a ring defines a finitely determined power structure over it in the following way. Any series $A(t) \in 1 + tR[[t]]$ can be in a unique way represented as $\prod_{i=1}^{\infty} \lambda_{b_i}(t^i)$, for some $b_i \in R$. Then one defines $(A(t))^m := \prod_{i=1}^{\infty} \lambda_{mb_i}(t^i)$. On the other hand, in general, there are many λ -ring structures corresponding to one power structure over a ring. One can show that the power structure (2) is defined by the λ -ring structure on the Grothendieck ring $K_0(\text{Var}_{\mathbb{C}})$ given by the Kapranov

4

check the reference

zeta function

$$\zeta_{[M]}(t) := 1 + \sum_{k=1}^{\infty} [S^k M] \cdot t^k,$$

where $S^k M$ is the kth symmetric power of the variety M. This follows from the following equation

$$\hat{S}_{[M]} = (1-t)^{-[M]} = (1+t+t^2+\cdots)^{[M]}.$$

Burnside ring A(G) of a finite group G is the Grothendieck ring of finite G-sets, i.e., finite sets with left actions of the group G: see, e.g., [7]. As an abelian group the Burnside ring A(G) is freely generated by the classes [G/H] of the quotients G/H for representatives H of the conjugacy classes \mathfrak{h} of subgroups of G. The multiplication is defined by the cartesian product with the diagonal action of G. There is a natural power structure over the Burnside ring A(G): see, e.g., [18]. This power structure is defined in a way similar to that one over the ring $K_0(\operatorname{Var}_{\mathbb{C}})$. Namely, if A_i , $i = 1, 2, \ldots$, and M are finite G-sets, the same equation (2) defines the series $(1 + [A_1]t + [A_2]t^2 + \cdots)^{[M]}$, where the action of the group G on the summands is the natural (diagonal) one. In particular, for a finite G-set X one has

$$(1-t)^{-[X]} = 1 + [X]t + [S^2X]t^2 + [S^3X]t^3 + \cdots,$$

where $S^k X = X^k / S_k$ is the *k*th symmetric power of the *G*-set *X* with the diagonal action of *G*.

Let X be a G-space, i.e., a space with a left action of the group G. For a point $x \in X$, let $G_x = \{g \in G : g \cdot x = x\}$ be the isotropy subgroup of the point x. For a subgroup $H \subset G$, let $X^H = \{x \in X : Hx = x\}$ be the fixed point set of H $(X^H = \{x \in X : H \subset G_x\})$ and let $X^{(H)} = \{x \in X : G_x = H\}$ be the set of points with the isotropy group H. Let Conjsub G be the set of the conjugacy classes of subgroups of G. For a conjugacy class $\mathfrak{h} \in \text{Conjsub } G$, let $X^{\mathfrak{h}} = \{x \in X : x \in X^H \text{ for a subgroup } H \in \mathfrak{h}\}, X^{(\mathfrak{h})} = \{x \in X : G_x \in \mathfrak{h}\}.$

The equivariant Euler characteristic of a (good enough) G-space X is defined by

$$\chi^G(X) := \sum_{\mathfrak{h} \in \text{Conjsub}\,G} \chi(X^{(\mathfrak{h})}/G)[G/H] \in A(G),\tag{3}$$

where H is a representative of the conjugacy class \mathfrak{h} (see, e.g., [29]).

3. Higher Order Euler Characteristics & MacDonald Type Equations

The orbifold Euler characteristic $\chi^{\text{orb}}(X, G)$ of the G-space X is defined, e.g., in [1], [20]:

$$\chi^{\text{orb}}(X, G) = \frac{1}{|G|} \sum_{\substack{(g_0, g_1) \in G \times G: \\ g_0 g_1 = g_1 g_0}} \chi(X^{\langle g_0, g_1 \rangle}) = \sum_{[g] \in G_*} \chi(X^{\langle g \rangle} / C_G(g)) \in \mathbb{Z}, \quad (4)$$

where G_* is the set of the conjugacy classes of elements of G, g is a representative of the class [g], $C_G(g) = \{h \in G : h^{-1}gh = g\}$ is the centralizer of g, and $\langle g \rangle$ and $\langle g_0, g_1 \rangle$ are the subgroups of G generated by the corresponding elements. check the reference

The higher order Euler characteristics of (X, G) are defined by:

$$\chi^{(k)}(X, G) = \frac{1}{|G|} \sum_{\substack{\mathbf{g} \in G^{k+1}:\\ g_i g_j = g_j g_i}} \chi(X^{\langle \mathbf{g} \rangle}) = \sum_{[g] \in G_*} \chi^{(k-1)}(X^{\langle g \rangle}, C_G(g)),$$
(5)

where $\mathbf{g} = (g_0, g_1, \ldots, g_k)$, $\langle \mathbf{g} \rangle$ is the subgroup generated by g_0, g_1, \ldots, g_k , and (for the second, recurrent, definition) $\chi^{(0)}(X, G)$ is defined as the usual Euler characteristic $\chi(X/G)$ of the quotient. The orbifold Euler characteristic $\chi^{\text{orb}}(X, G)$ is the Euler characteristic $\chi^{(1)}(X, G)$ of order 1.

For a G-space X, the cartesian power X^n carries the natural action of the wreath product $G_n = G^n \rtimes S_n$ generated by the natural action of the symmetric group S_n (permuting the factors) and by the natural (component-wise) action of the cartesian power G^n . The pair (X_n, G_n) should be (or can be) considered as an analogue of the symmetric power of the pair (X, G). Let

$$\mathbb{X}^{(k)}_{(X,G)}(t) := 1 + \sum_{n \ge 1} \chi^{(k)}(X^n, G_n) \cdot t^n.$$

One has the following Macdonald type equation (see [28, Theorem A])

$$\sum_{n \ge 0} \chi^{(k)}(X^n, G_n) \cdot t^n = \left(\prod_{r_1, \dots, r_k \ge 1} (1 - t^{r_1 r_2 \dots r_k})^{r_2 r_3^2 \dots r_k^{k-1}}\right)^{-\chi^{(n)}(X, G)}.$$
 (6)

(k) (\mathbf{r} \mathbf{c})

When k = 0, one gets the equation (1) for the quotient X/G.

The equation (6) can be interpreted in the following way. For the natural (left) action of the group G on G (by the multiplication) one has $\chi^{(k)}(G, G) = 1$ for any $k \ge 1$. (This follows from the fact that the only element $g \in G$ with a non-empty fixed point set $G^{\langle g \rangle}$ is g = e and $G^{\langle e \rangle} = G$.) Therefore

$$\mathbb{X}_{(G,G)}^{(k)}(t) = \left(\prod_{r_1,\dots,r_k \ge 1} (1 - t^{r_1 r_2 \cdots r_k})^{r_2 r_3^2 \cdots r_k^{k-1}}\right)^{-1}$$

and thus

$$\mathbb{X}_{(X,G)}^{(k)}(t) = \left(\mathbb{X}_{(G,G)}^{(k)}(t)\right)^{\chi^{(k)}(X,G)}.$$

Let $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]_{s\in\mathbb{Q}}$ (or $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ for short) be the modification of the Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$ obtained by adding all rational powers of \mathbb{L} . The elements of $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ are in one-to-one correspondence with the finite sums of the form $\sum_i c_i \mathbb{L}^{r_i}$, where c_i are elements of the localization $K_0(\operatorname{Var}_{\mathbb{C}})_{(\mathbb{L})}$ of the ring $K_0(\operatorname{Var}_{\mathbb{C}})$ by the class \mathbb{L} , r_i are different rational numbers inbetween 0 and 1: $0 \leq r_i < 1$. Thus the ring $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ contains the localization $K_0(\operatorname{Var}_{\mathbb{C}})_{(\mathbb{L})}$. It was shown that \mathbb{L} is a zero divisor in $K_0(\operatorname{Var}_{\mathbb{C}})$ ([6]) and therefore the natural map $K_0(\operatorname{Var}_{\mathbb{C}}) \to K_0(\operatorname{Var}_{\mathbb{C}})_{(\mathbb{L})}$ is not injective. Therefore the natural map $K_0(\operatorname{Var}_{\mathbb{C}}) \to K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ is not injective as well.

A power structure on the ring $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ can be defined through a λ -structure on it using the equation:

$$\zeta_{\mathbb{L}^s[M]}(t) = \zeta_{[M]}(\mathbb{L}^s t),\tag{7}$$

which holds for the Kapranov zeta function on the Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$. This equation (used for integer values of s) defines a λ -structure on the localization $K_0(\operatorname{Var}_{\mathbb{C}})_{(\mathbb{L})}$: see [15]. For an element $c = \sum_i c_i \mathbb{L}^{r_i} \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ one defines $\zeta_c(t)$ by

$$\zeta_c(t) := \prod_i \zeta_{c_i}(\mathbb{L}^{r_i} t).$$

The λ -structure on the ring $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ defines a power structure over it in the standard way [15]:-a series $A(t) \in 1+tK_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s][[t]]$ has a unique representation in the form $A(t) = \prod_{i=1}^{\infty} \zeta_{b_i}(t^i)$ with $b_i \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ and one defines $(A(t))^m$, $m \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$, by

$$(A(t))^m := \prod_{i=1}^{\infty} \zeta_{mb_i}(t^i).$$

This definition together with the equation (7) implies that

$$(A(\mathbb{L}^s t))^m = (A(t))^m|_{t \mapsto \mathbb{L}^s t}.$$

Remark. If, in the considerations below, one uses only non-negative weights φ_i , one can work with the ring $K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]_{s\in\mathbb{Q}_{\geq 0}}$. The natural map $K_0(\operatorname{Var}_{\mathbb{C}}) \to K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]_{s\in\mathbb{Q}_{\geq 0}}$ is injective.

Now let X be a smooth quasi-projective variety of dimension d with an (algebraic) action of the group G. To define the higher order generalized (orbifold) Euler characteristics of the pair (X, G), one has to use the so called age (or fermion shift) F_x^g of an element $g \in G$ at a fixed point x of g defined in [32], [21]. The element g acts on the tangent space $T_x X$ as an automorphism of finite order. This action on $T_x X$ can be represented by a diagonal matrix diag $(\exp(2\pi i\theta_1), \ldots, \exp(2\pi i\theta_d))$ with $0 \leq \theta_j < 1$ for $j = 1, 2, \ldots, d$ (θ_j are rational numbers). The age of the element g at the point x is defined by $F_x^g = \sum_{j=1}^d \theta_j \in \mathbb{Q}_{\geq 0}$.

For $g \in G$, let the number of $C_G(g)$ -orbits in the the set of connected components of the fixed point set $X^{\langle g \rangle}$ be equal to N_g , and let $X_1^{\langle g \rangle}$, $X_2^{\langle g \rangle}$, ..., $X_{N_g}^{\langle g \rangle}$ be the unions of the components of each of the orbits. For $1 \leq \alpha_g \leq N_g$, let $F_{\alpha_g}^g$ be the age F_x^g at a point of $X_{\alpha_g}^{\langle g \rangle}$ (this age does not depend on the point $x \in X_{\alpha_g}^{\langle g \rangle}$).

For a rational number $\varphi_1 \in \mathbb{Q}$, the generalized orbifold Euler characteristic of weight φ_1 of the pair (X, G) is defined by

$$[X, G]_{\varphi_1} := \sum_{[g]\in G_*} \sum_{\alpha_g=1}^{N_g} [X_{\alpha_g}^{\langle g \rangle} / C_G(g)] \cdot \mathbb{L}^{\varphi_1 F_{\alpha_g}^{\langle g \rangle}} \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s].$$
(8)

For $\varphi_1 = 1$ one gets the definition of the generalized orbifold Euler characteristic from [17] inspired by the definition of the orbifold Hodge–Deligne polynomial from [2]. (This generalized orbifold Euler characteristic maps to the orbifold Hodge– Deligne polynomial by the natural ring homomorphism $e: K_0(\operatorname{Var}_{\mathbb{C}}) \to \mathbb{Z}[u, v]$.) For $\varphi_1 = 0$ one gets the so called inertia stack class: see, e.g., [13].

For a subgroup $H \subset G$, and for an H-invariant submanifold $Y \subset X$, let us define the relative generalized orbifold Euler characteristic of the pair (Y, H) by

$$[Y, H]_{X,\varphi_1} := \sum_{[g]\in H_*} \sum_{\alpha_g=1}^{N_g} [Y_{\alpha_g}^{\langle g \rangle} / C_H(g)] \cdot \mathbb{L}^{\varphi_1 F_{\alpha_g}^{\langle g \rangle}} \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s], \tag{9}$$

where $Y_{\alpha_g}^{\langle g \rangle} := X_{\alpha_g}^{\langle g \rangle} \cap Y$, $F_{\alpha_g}^{\langle g \rangle}$ is the age of g at a point of $X_{\alpha_g}^{\langle g \rangle}$. (Pay attention that the age of an element g is determined from its action on $T_x X$.)

Let $\varphi = (\varphi_1, \varphi_2, ...)$ be a fixed sequence of rational numbers. The *relative* generalized (orbifold) Euler characteristics of order k of weight φ of the pair (Y, H)is defined recursively by

$$[Y, H]_{X, \underline{\varphi}}^{k} := \sum_{[g] \in H_*} \sum_{\alpha_g = 1}^{N_g} [Y_{\alpha_g}^{\langle g \rangle}, C_H(g)]_{X, \underline{\varphi}}^{k-1} \cdot \mathbb{L}^{\varphi_k F_{\alpha_g}^{\langle g \rangle}} \in K_0(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s],$$
(10)

where $[Y, H]^1_{X,\varphi} := [Y, H]_{X,\varphi_1}$ is defined by (9). (Alternatively one can start from k = 0 using the definition $[Y, H]^0_{X, \underline{\varphi}} = [Y, H]^0 = [Y/H]$.) The generalized (orbifold) Euler characteristics of order k of weight φ of the pair (X, G) is

$$[X, G]^k_{\underline{\varphi}} := [X, G]^k_{X, \underline{\varphi}}$$

One has the following Macdonald type equations ([19])

$$\sum_{n \ge 0} [X^n, G_n]_{\underline{\varphi}}^k \cdot t^n = \left(\prod_{r_1, \dots, r_k \ge 1} \left(1 - \mathbb{L}^{\Phi_k(\underline{r})d/2} \cdot t^{r_1 r_2 \cdots r_k}\right)^{r_2 r_3^2 \cdots r_k^{k-1}}\right)^{-[X, G]_{\underline{\varphi}}^k}, \quad (11)$$

where

$$\Phi_k(r_1, \ldots, r_k) = \varphi_1(r_1 - 1) + \varphi_2 r_1(r_2 - 1) + \cdots + \varphi_k r_1 r_2 \cdots r_{k-1}(r_k - 1).$$

4. Equivariant Higher Order Euler Characteristics

Assume that X is a (good enough) topological space with *commuting* actions of two finite groups G_O and G_B (or equivalently with an action of the direct product $G_O \times G_B$). The quotient X/G_O carries the natural G_B -action and thus one can define $\chi^{(0,G_B)}(X; G_O, G_B)$ as $\chi^{G_B}(X/G_O) \in A(G_B)$. For an element $g \in G_O$, the fixed point set $X^{\langle g \rangle}$ is G_B -invariant and the quotient

 $X^{\langle g \rangle}/C_{G_O}(g)$ by the centralizer $C_{G_O}(g)$ carries the natural G_B -action.

Definition. The equivariant orbifold Euler characteristics of $(X; G_O, G_B)$ is

$$\chi^{(1,G_B)}(X; G_O, G_B) := \sum_{[g] \in G_{O*}} \chi^{G_B}(X^{\langle g \rangle} / C_{G_O}(g))$$
$$= \sum_{[g] \in G_{O*}} \chi^{(0,G_B)}(X^{\langle g \rangle}; C_{G_O}(g), G_B) \in A(G_B).$$
(12)

The equivariant higher order Euler characteristics are defined recursively in the same way as the non-equivariant one.

Definition. The equivariant Euler characteristics of order k of $(X; G_O, G_B)$ is

$$\chi^{(k,G_B)}(X; G_O, G_B) := \sum_{[g] \in G_{O_*}} \chi^{(k-1,G_B)}(X^{\langle g \rangle}; C_{G_O}(g), G_B) \in A(G_B).$$
(13)

Definition (13) is convenient for the proof of Theorem 1. Iterating (13) one gets the following equation for the equivariant higher order Euler characteristics:

$$\chi^{(k,G_B)}(X; G_O, G_B) = \sum_{[\phi] \in \operatorname{Hom}\left(\mathbb{Z}^k, G_O\right)/G_O} \chi^{G_B}(X^{\langle \phi \rangle}/C_{G_O}(\phi)), \qquad (14)$$

where the group G_O acts on Hom (\mathbb{Z}^k, G_O) by conjugation, $X^{\langle \phi \rangle}$ is the fixed point set of the image of ϕ , $C_{G_O}(\phi) = \{g \in G_O : g^{-1}\phi g = \phi\}.$

Examples. 1. Let $X = G_O$ be endowed with the left action of the group G_O defined by the multiplication and with the trivial action of the group G_B . Since the only element $g \in G_O$ with a non-empty fixed point set $G_O^{(g)}$ is the unit g = e and $G_O^{(e)} = G_O$, one has $\chi^{(k,G_B)}(G_O; G_O, G_B) = 1$ for any $k \ge 0$. 2. Let $X = G_B$ and let G_O be a subgroup of G_B . Define the action of the group

2. Let $X = G_B$ and let G_O be a subgroup of G_B . Define the action of the group G_B (respectively of the group G_O) on X by $ga = g \cdot a$ ($ag = a \cdot g^{-1}$ respectively) for $a \in X$, $g \in G_B$ ($g \in G_O$ respectively). Since the only element $g \in G_O$ with a non-empty fixed point set $G_O^{\langle g \rangle}$ is the unit g = e and $G_B^{\langle e \rangle} = G_B$, one has $\chi^{(k,G_B)}(G_B; G_O, G_B) = [G_B/G_O] \in A(G)$ for any $k \ge 0$.

$$\mathbb{X}_{(X;G_O,G_B)}^{(k)}(t) := 1 + \sum_{n \ge 1} \chi^{(k,G_B)}(X^n; G_{O_n}, G_B) \cdot t^n.$$

Theorem 1. One has

$$\mathbb{X}_{(X;G_O,G_B)}^{(k)}(t) = \left(\prod_{r_1,\dots,r_k \ge 1} (1 - t^{r_1 r_2 \cdots r_k})^{r_2 r_3^2 \cdots r_k^{k-1}}\right)^{-\chi^{(k,G_B)}(X;G_O,G_B)}, \quad (15)$$

where the exponent in the right hand side is defined by the power structure over the Burnside ring $A(G_B)$.

Proof. The proof essentially repeats, e.g., the one in [28] (see also[19]). One has to pay attention to two facts. First, all subsets participating in the course of the proof in [28] are *G*-invariant. Second, one has to use the Macdonald type equation for the equivariant Euler characteristic $\chi^G(\cdot)$: see (16) below.

The Macdonald type equation for the equivariant Euler characteristic must be known. However, we have not found it in the literature. Therefore we put its proof here.

Lemma 1. For a G-space X one has

$$1 + \sum_{n=1}^{\infty} \chi^G(S^n X) \cdot t^n = (1-t)^{-\chi^G(X)} \in 1 + tA(G)[[t]],$$
(16)

where the right hand side is defined by the power structure over the Burnside ring A(G) (or by the λ -structure on it).

Remark. Note that, for a finite G-set X, the equation (16) is just the definition of its right hand side through the power structure over the Burnside ring described in Section 2. (In this case $\chi^G(X) = [X] \in A(G)$.)

Proof. Let us denote the left hand side of (16) by $\chi^G \zeta_{X,G}(t)$. If $X = X_1 \sqcup X_2$ is a decomposition of X into two G-subspaces, one has

$$\chi^G \zeta_{X,G}(t) = \chi^G \zeta_{X_1,G}(t) \cdot \chi^G \zeta_{X_2,G}(t)$$

(This follows from the identities $S^n X = \prod_{m=0}^n (S^m X_1) \times (S^{n-m} X_2), \ \chi^G (X \times Y) =$

 $\chi^G(X)\chi^G(Y)$.) Therefore it is sufficient to prove (16) for the elements of a decomposition of X into G-invariant subspaces. A "good enough" G-space X (say, a real subanalytic space with a subanalytic action of the group G) can be represented as the disjoint union of subspaces of the form $\sigma^d \times (G/H)$, where H is a subgroup of G, G/H is the corresponding G-set (the quotient), and σ^d is the open cell of dimension d with the trivial G-action. (This follows, e.g., from the fact that the quotient X/G has a triangulation coordinated by the stratification of this quotient by the types of the G-orbits.) Let $\bar{\sigma}^d \supset \sigma^d$ be the closed d-dimensional ball. Since $\bar{\sigma}^d$ can be contracted to a point, $S^k(\bar{\sigma}^d \times (G/H))$ can be G-equivariantly contracted to $S^k(G/H)$. Therefore

$$\chi^G(S^k(\bar{\sigma}^d \times (G/H))) = \chi^G(S^k(G/H)) = [S^k(G/H)]$$

(see, e.g., [29], where this is formulated for finite G-CW-complexes) and thus

)

$$\chi^G \zeta_{S^k(\bar{\sigma}^d \times (G/H))}(t) = 1 + \sum_{i=1}^{\infty} [S^i(G/H)]t^i = (1-t)^{-[G/H]} = (1-t)^{-\chi^G(\bar{\sigma}^d \times (G/H))}.$$

The equation (16) obviously holds for $X = \sigma^d \times (G/H)$ with d = 0 (when σ^d is a point). Assume that it holds for $X = \sigma^d \times (G/H)$ with $d < d_0$. One has a decomposition $\sigma^{d_0} = \sigma^{d_0} \sqcup \sigma^{d_0} \sqcup \sigma^{d_0-1}$. Therefore

$$\chi^{G}\zeta_{\sigma^{d_{0}}\times(G/H)}(t) = \left(\chi^{G}\zeta_{\sigma^{d_{0}}\times(G/H)}(t)\right)^{2} \cdot \chi^{G}\zeta_{\sigma^{(d_{0}-1)}\times(G/H)}(t),$$

$$\chi^{G}\zeta_{\sigma^{d_{0}}\times(G/H)}(t) = \left(\chi^{G}\zeta_{\sigma^{(d_{0}-1)}\times(G/H)}(t)\right)^{-1} = \left((1-t)^{-(-1)^{(d_{0}-1)}[G/H]}\right)^{-1}$$

$$= (1-t)^{-(-1)^{d_{0}}[G/H]} = (1-t)^{-\chi^{G}(\sigma^{d_{0}}\times(G/H))}.$$

The equation (15) has an interpretation similar to the one for "usual" (nonequivariant) higher order Euler characteristics in Section 3. Set

Set what?

Let G_{O} (regarded as a zero-dimensional finite space) be endowed with the natural action of the group G_O and with the trivial action of the group G_B . Then one has $\chi^{(k,G_B)}(G_O; G_O, G_B) = 1$. Let $(G_O; G_O, G_B)$ be the set G_O with the actions described in Example 1 after the definition of higher order equivariant Euler characteristics. One has

$$\mathbb{X}_{(G_O;G_O,G_B)}^{(k)}(t) = \left(\prod_{r_1,\dots,r_k \geqslant 1} (1 - t^{r_1 r_2 \dots r_k})^{r_2 r_3^2 \dots r_k^{k-1}}\right)^{-1}$$
$$\mathbb{X}_{(X;G_O,G_B)}^{(k)}(t) = \left(\mathbb{X}_{(G_O;G_O,G_D)}^{(k)}(t)\right)^{\chi^{(k,G_B)}(X;G_O,G_B)}.$$

and thus

$$\mathbb{X}_{(X;G_O,G_B)}^{(k)}(t) = \left(\mathbb{X}_{(G_O;G_O,G_B)}^{(k)}(t)\right)^{\chi^{(k,G_B)}(X;G_O,G_B)}.$$

10

5. Equivariant Generalized Higher Order Euler Characteristics

For a finite group G, let $K_0^G(\operatorname{Var}_{\mathbb{C}})$ be the Grothendieck ring of complex quasiprojective G-varieties (i.e., varieties with left algebraic actions of the group G). By that we mean the free abelian group generated by the G-isomorphism classes [X, G](or [X] for short) of complex quasi-projective varieties X with G-actions modulo the relation: $[X, G] = [Y, G] + [X \setminus Y, G]$ for a Zariski closed G-invariant subvariety Y of X. The multiplication in $K_0^G(\operatorname{Var}_{\mathbb{C}})$ is defined by the cartesian product with the diagonal G-action. Let $\mathbb{L} \in K_0^G(\operatorname{Var}_{\mathbb{C}})$ be the class of the affine line $\mathbb{A}^1_{\mathbb{C}}$ with the trivial G-action.

Remark. Usually, in the definition of the Grothendieck ring of complex quasiprojective *G*-varieties, one adds one more relation: if $E \to X$ is a *G*-equivariant vector bundle of rank *n*, then $[E] = [\mathbb{A}^n_{\mathbb{C}} \times X]$. We do not need this relation for the construction. One can say that we use the Grothendieck ring denoted by $K_0^{',G}(\operatorname{Var}_{\mathbb{C}})$ in [5]. The same definition was used in [25]. An equation which holds in the equivariant Grothendieck ring $K_0^G(\operatorname{Var}_{\mathbb{C}})$ defined here, holds in the "traditional" one as well.

There is a natural power structure over the (equivariant) Grothendieck ring $K_0^G(\operatorname{Var}_{\mathbb{C}})$. Its geometric definition is given in the same way as the usual power structure over the (non-equivariant) Grothendieck ring $K_0(\operatorname{Var}_{\mathbb{C}})$ in [15]: for complex quasi-projective *G*-varieties A_i , $i = 1, 2, \ldots$, and *M* one has the same equation (2). One has to take into account that all summands in the right hand side of the equation (2) are *G*-invariant spaces. The proof of the necessary properties of the power structure is the same as in [15]. This power structure is induced by the λ -structure on $K_0^G(\operatorname{Var}_{\mathbb{C}})$ defined by the Kapranov zeta-function. For a quasi-projective *G*-variety *X*, the series $(1 - t)^{-[X]}$ is the equivariant Kapranov zeta-function of $X: \zeta_{[X]}(t) := 1 + [X] \cdot t + [S^2X] \cdot t^2 + [S^3X] \cdot t^3 + \cdots = (1 - t)^{-[X]}$, where $S^kX = X^k/S_k$ is the *k*th symmetric power of the *G*-variety *X* with the natural *G*-action (see, e.g., [22], [25] for the non-equivariant case). The map $\chi^G: K_0^G(\operatorname{Var}_{\mathbb{C}}) \to A(G)$ is a λ -ring homomorphism.

In what follows we need the following statement.

Lemma 2. Let $p: E \to X$ be a *G*-equivariant vector bundle of rank *n* such that for each $x \in X$ the action of the isotropy subgroup G_x on the fibre $E_x = p^{-1}(x)$ is trivial. Then $[E] = \mathbb{L}^n[X]$.

Proof. Factorizing by the action of G one gets the map $\check{p}: E/G \to X/G$, which is a vector bundle (due to the triviality of the action of G_x on E_x). According to [26] the quotient X/G can be covered by Zariski open subsets U_i such that over each U_i the fibre bundle \check{p} is trivial. Therefore

$$\check{p}^{-1}(U_i) \cong U_i \times \mathbb{A}^n_{\mathbb{C}}.$$
(17)

If $V_i = \pi^{-1}(U_i)$, where $\pi \colon X \to X/G$ is the canonical factorization map, then the trivialization (17) gives a trivialization of the bundle p over V_i , i.e., an isomorphism between $p^{-1}(V_i)$ and $V_i \times \mathbb{A}^n_{\mathbb{C}}$ with the trivial G-action on $\mathbb{A}^n_{\mathbb{C}}$. This gives the statement.

In what follows we need the following properties of the power structure over the equivariant Grothendieck ring $K_0^G(\operatorname{Var}_{\mathbb{C}})$.

Proposition 1. Let A_i and M be G-varieties, and let $A(t) := 1 + [A_1]t + [A_2]t^2 + \cdots \in K_0^G(\operatorname{Var}_{\mathbb{C}})$. Then, for $s \ge 0$,

$$(A(\mathbb{L}^{s}t))^{[M]} = (A(t))^{[M]}|_{t \mapsto \mathbb{L}^{s}t}.$$
(18)

Proof. The coefficient at the monomial t^k in the power series $(A(t))^{[M]}$ is a sum of the classes of the varieties of the form

$$V = \left(\left(\prod_{i} M^{k_{i}}\right) \setminus \Delta \right) \times \prod_{i} A_{i}^{k_{i}} / \prod_{i} S_{k_{i}}$$

with $\sum i k_i = k$. The corresponding summand \widetilde{V} in the coefficient at the monomial t^k in the power series $(A(\mathbb{L}^s t))^M$ has the form

$$\widetilde{V} = \left(\left(\prod_{i} M^{k_i}\right) \setminus \Delta \right) \times \prod_{i} \left(\mathbb{L}^{si} A_i \right)^{k_i} / \prod_{i} S_{k_i}.$$

There exists a natural map $\widetilde{V} \to V$ which is a *G*-equivariant vector bundle of rank sk satisfying the conditions of Lemma 2 (just because the action of the group $\prod_i S_{k_i}$ is free). By (2) one has $[\widetilde{V}] = \mathbb{L}^{sk}[V]$, what implies (18).

Proposition 1 and the multiplicative property $(A(t))^{[M][N]} = (A(t)^{[M]})^{[N]}$ of the power structure imply the following statement.

Proposition 2. For a complex quasi-projective G-variety X one has

$$\zeta_{\mathbb{L}[X]}(t) = \zeta_{[X]}(\mathbb{L}t). \tag{19}$$

Here, in a setting similar to the considered above, we suggest an equivariant version of the generalized higher order Euler characteristic with values in the modification $K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s] = K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]_{s \in \mathbb{Q}}$ of the equivariant Grothendieck ring of complex quasi-projective varieties.

Let X be a smooth quasi-projective variety of dimension d with commuting (algebraic) actions of two finite groups G_O and G_B (or equivalently with an action of the product $G_O \times G_B$).

Define the zero order equivariant generalized Euler characteristic of $(X; G_O, G_B)$ by

$$[X; G_O, G_B]^{0, G_B} := [X/G_O] \in K_0^{G_B}(\text{Var}_{\mathbb{C}}).$$
(20)

For $g \in G_O$, let the set of the connected components of the fixed point set $X^{\langle g \rangle}$ consist of N_g ($C_{G_O}(g) \times G_B$)-orbits and let $X_1^{\langle g \rangle}$, $X_2^{\langle g \rangle}$, ..., $X_{N_g}^{\langle g \rangle}$ be the unions of the components of each of the orbits. For $1 \leq \alpha_g \leq N_g$ let $F_{\alpha_g}^g$ be the age F_x^g at a point of $X_{\alpha_g}^{\langle g \rangle}$ (this age does not depend on the point $x \in X_{\alpha_g}^{\langle g \rangle}$).

12

Definition. For a rational number $\varphi_1 \in \mathbb{Q}$, the generalized orbifold Euler characteristic of weight φ_1 of $(X; G_O, G_B)$ is defined by

$$[X; G_O, G_B]_{\varphi_1}^{1, G_B} := \sum_{[g] \in (G_O)_*} \sum_{\alpha_g = 1}^{N_g} [X_{\alpha_g}^{\langle g \rangle} / C_{G_O}(g)] \cdot \mathbb{L}^{\varphi_1 F_{\alpha_g}^{\langle g \rangle}}$$
$$= \sum_{[g] \in (G_O)_*} \sum_{\alpha_g = 1}^{N_g} [X; G_O, G_B]^{0, G_B} \cdot \mathbb{L}^{\varphi_1 F_{\alpha_g}^{\langle g \rangle}} \in K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s].$$
(21)

For a subgroup $H_O \subset G_O$ and for an $H_O \times G_B$ -invariant submanifold $Y \subset X$, let us define the *relative generalized orbifold Euler characteristic* of $(Y; H_O, G_B)$ by

$$[Y; H_O, G_B]_{X,\varphi_1}^{1,G_B} := \sum_{[g]\in(H_O)_*} \sum_{\alpha_g=1}^{N_g} [Y_{\alpha_g}^{\langle g \rangle} / C_{H_O}(g)] \cdot \mathbb{L}^{\varphi_1 F_{\alpha_g}^{\langle g \rangle}} \in K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]_{s\in\mathbb{Q}},$$
(22)

where $Y_{\alpha_g}^{\langle g \rangle} := X_{\alpha_g}^{\langle g \rangle} \cap Y$, $F_{\alpha_g}^{\langle g \rangle}$ is the age of g at a point of $X_{\alpha_g}^{\langle g \rangle}$. Let $\underline{\varphi} = (\varphi_1, \varphi_2, \dots)$ be a fixed sequence of rational numbers.

Definition. The relative equivariant generalized Euler characteristics of order k of weight $\underline{\varphi}$ of $(Y; H_O, G_B)$, as an element of $K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$, is defined recursively by

$$[Y; H_O, G_B]_{X, \underline{\varphi}}^{k, G_B} := \sum_{[g] \in (G_O)_*} \sum_{\alpha_g = 1}^{N_g} [Y_{\alpha_g}^{\langle g \rangle}; C_{H_O}(g), G_B]_{X, \underline{\varphi}}^{(k-1), G_B} \cdot \mathbb{L}^{\varphi_k F_{\alpha_g}^{\langle g \rangle}},$$

where $[Y; H_O, G_B]_{X, \underline{\varphi}}^{0, G_B} = [Y; H_O, G_B]^{0, G_B}$ is the zero order equivariant generalized Euler characteristic given by (20).

The equivariant generalized Euler characteristics of order k of weight $\underline{\varphi}$ of the triple $(X; G_O, G_B)$ is

$$[X; G_O, G_B]_{\underline{\varphi}}^{k, G_B} := [X; G_O, G_B]_{X, \underline{\varphi}}^{k, G_B} \in K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s].$$

Remark. Iterating this definition one can write an equation for $[X; G_O, G_B]_{\underline{\varphi}}^{k, G_B}$ analogous to (14). For a homomorphism $\phi : \mathbb{Z}^k \to G_O$, let the set of the connected components of the fixed point set $X^{\langle \phi \rangle}$ consist of N_{ϕ} ($C_{G_O}(g) \times G_B$)-orbits and let $X_1^{\langle \phi \rangle}, X_2^{\langle \phi \rangle}, \ldots, X_{N_{\phi}}^{\langle \phi \rangle}$ be the unions of the components of each of the orbits. For $x \in X^{\langle \phi \rangle}$, let the shift F_x^{ϕ} be defined as $\sum_{i=1}^k \varphi_i F_x^{\phi(e_i)}$, where e_1, \ldots, e_k is the standard basis of \mathbb{Z}^k . For $1 \leq \alpha_{\phi} \leq N_g$ let $F_{\alpha_{\phi}}^{\phi}$ be the shift F_x^{ϕ} at a point of $X_{\alpha_{\phi}}^{\langle \phi \rangle}$. One has:

$$[X; G_O, G_B]_{\underline{\varphi}}^{k, G_B} = \sum_{[\phi] \in \operatorname{Hom}(\mathbb{Z}^k, G_O)/G_O} \sum_{\alpha_{\phi} = 1}^{N_{\phi}} [X^{\langle \phi \rangle}/C_{G_O}(\phi)] \cdot \mathbb{L}^{F_{\alpha_{\phi}}^{\phi}}.$$

Theorem 2. Let X be a (smooth) quasi-projective variety of dimension d with commuting actions of two finite groups G_O and G_B . Then

$$\sum_{n \ge 0} [X^n; (G_O)_n, G_B]_{\underline{\varphi}}^{k, G_B} \cdot t^n = \\ = \left(\prod_{r_1, \dots, r_k \ge 1} (1 - \mathbb{L}^{\Phi_k(\underline{r})d/2} \cdot t^{r_1 r_2 \cdots r_k})^{r_2 r_3^2 \cdots r_k^{k-1}}\right)^{-[X; G_0, G_B]_{\underline{\varphi}}^k}$$

where

 $\Phi_k(r_1, \ldots, r_k) = \varphi_1(r_1 - 1) + \varphi_2 r_1(r_2 - 1) + \cdots + \varphi_k r_1 r_2 \cdots r_{k-1}(r_k - 1).$

Proof. The proof in [19] was by induction started from k = 1, i.e., from the generalized orbifold case. The latter one was treated in [17]. One can easily see that the both proofs admit an action of an additional group G_B , i.e., all the subspaces are G_B -invariants. In particular, symmetric products participating in the proof of [17] carry the natural action of G_B . Using Propositions 1 and 2 the corresponding generating series can be written as exponents in terms of the power structure over the modification $K_0^{G_B}(\operatorname{Var}_{\mathbb{C}})[\mathbb{L}^s]$ of the equivariant Grothendieck ring of quasiprojective varieties.

References

- M. Atiyah and G. Segal, On equivariant Euler characteristics, J. Geom. Phys. 6 (1989), no. 4, 671–677. MR 1076708
- [2] V. V. Batyrev and D. I. Dais, Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry, Topology 35 (1996), no. 4, 901–929. MR 1404917
- [3] P. Berglund and M. Henningson, Landau-Ginzburg orbifolds, mirror symmetry and the elliptic genus, Nuclear Phys. B 433 (1995), no. 2, 311–332. MR 1310310
- [4] P. Berglund and T. Hübsch, A generalized construction of mirror manifolds, Nuclear Phys. B 393 (1993), no. 1–2, 377–391. MR 1214325
- [5] F. Bittner, The universal Euler characteristic for varieties of characteristic zero, Compos. Math. 140 (2004), no. 4, 1011–1032. MR 2059227
- [6] L. Borisov, Class of the affine line is a zero divisor in the Grothendieck ring, preprint arXiv:1412.6194 [math.AG].
- [7] S. Bouc, Burnside rings, Handbook of algebra, Vol. 2, North-Holland, Amsterdam, 2000, pp. 739–804. MR 1759611
- [8] J. Bryan and J. Fulman, Orbifold Euler characteristics and the number of commuting mtuples in the symmetric groups, Ann. Comb. 2 (1998), no. 1, 1–6. MR 1682916
- J. Bryan and A. Morrison, Motivic classes of commuting varieties via power structures, J. Algebraic Geom. 24 (2015), no. 1, 183–199. MR 3275657
- [10] L. Dixon, J. Harvey, C. Vafa, and E. Witten, *Strings on orbifolds*, Nuclear Phys. B 261 (1985), no. 4, 678–686. MR 818423
- [11] L. Dixon, J. Harvey, C. Vafa, and E. Witten, *Strings on orbifolds. II*, Nuclear Phys. B 274 (1986), no. 2, 285–314. MR 851703
- [12] W. Ebeling and S. M. Gusein-Zade, Orbifold zeta functions for dual invertible polynomials, preprint arXiv:1407.0154 [math.AG]; to appear in Proc. Edinb. Math. Soc.
- [13] T. de Fernex, E. Lupercio, T. Nevins, and B. Uribe, Stringy Chern classes of singular varieties, Adv. Math. 208 (2007), no. 2, 597–621. MR 2304329
- [14] E. Gorsky, Adams operations and power structures, Mosc. Math. J. 9 (2009), no. 2, 305–323, back matter. MR 2568440
- [15] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández, A power structure over the Grothendieck ring of varieties, Math. Res. Lett. 11 (2004), no. 1, 49–57. MR 2046199

14

- [16] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández, Power structure over the Grothendieck ring of varieties and generating series of Hilbert schemes of points, Michigan Math. J. 54 (2006), no. 2, 353–359. MR 2252764
- [17] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández, On the power structure over the Grothendieck ring of varieties and its applications, Tr. Mat. Inst. Steklova 258 (2007), Anal. i Osob. Ch. 1, 58–69 (Russian). MR 2400523. English translation: Proc. Steklov Inst. Math. 258 (2007), no. 1, 53–64.
- [18] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández, On an equivariant version of the zeta function of a transformation, Arnold Math. J. 1 (2015), no. 2, 127–140. MR 3370062
- [19] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández, Higher order generalized Euler characteristics and generating series, J. Geom. Phys. 95 (2015), 137–143. MR 3357827
- [20] F. Hirzebruch and T. Höfer, On the Euler number of an orbifold, Math. Ann. 286 (1990), no. 1–3, 255–260. MR 1032933
- [21] Y. Ito and M. Reid, The McKay correspondence for finite subgroups of SL(3, C), Higherdimensional complex varieties (Trento, 1994), de Gruyter, Berlin, 1996, pp. 221–240. MR 1463181
- [22] M. Kapranov, The elliptic curve in the S-duality theory and Eisenstein series for Kac-Moody groups, preprint arXiv:math/0001005 [math.AG].
- [23] D. Knutson, λ-rings and the representation theory of the symmetric group, Lecture Notes in Mathematics, Vol. 308, Springer-Verlag, Berlin-New York, 1973. MR 0364425
- [24] I. G. Macdonald, The Poincaré polynomial of a symmetric product, Proc. Cambridge Philos. Soc. 58 (1962), 563–568. MR 0143204
- [25] J. Mazur, Rationality of motivic zeta functions for curves with finite abelian group actions, J. Pure Appl. Algebra 217 (2013), no. 7, 1335–1349. MR 3019741
- [26] J.-P. Serre, Espaces fibrés algébriques, Séminaire Claude Chevalley, 3 (1958), Exposé No. 1, 37 pages.
- [27] R. P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999. MR 1676282
- [28] H. Tamanoi, Generalized orbifold Euler characteristic of symmetric products and equivariant Morava K-theory, Algebr. Geom. Topol. 1 (2001), 115–141 (electronic). MR 1805937
- [29] T. tom Dieck, Transformation groups and representation theory, Lecture Notes in Mathematics, vol. 766, Springer, Berlin, 1979. MR 551743
- [30] W. Wang, Equivariant K-theory, wreath products, and Heisenberg algebra, Duke Math. J. 103 (2000), no. 1, 1–23. MR 1758236
- [31] W. Wang and J. Zhou, Orbifold Hodge numbers of the wreath product orbifolds, J. Geom. Phys. 38 (2001), no. 2, 152–169. MR 1823666
- [32] E. Zaslow, Topological orbifold models and quantum cohomology rings, Comm. Math. Phys. 156 (1993), no. 2, 301–331. MR 1233848

Moscow State University, Faculty of Mathematics and Mechanics, GSP-1, Moscow, 119991, Russia

E-mail address: sabir@mccme.ru

ICMAT (CSIC-UAM-UC3M-UCM); COMPLUTENSE UNIVERSITY OF MADRID, DEPT. OF AL-GEBRA, MADRID, 28040, SPAIN.

E-mail address: iluengo@mat.ucm.es *E-mail address*: amelle@mat.ucm.es