

EXISTENCE AND ASYMPTOTIC BEHAVIOUR FOR A DEGENERATE KIRCHHOFF-CARRIER MODEL WITH VISCOSITY AND NONLINEAR BOUNDARY CONDITIONS

M.M. CAVALCANTI, V.N. DOMINGOS CAVALCANTI,
J.A. SORIANO and J.S. PRATES FILHO

Abstract

The present paper studies the existence and uniqueness of global solutions and decay rates to the nonlinear hyperbolic problem

$$(*) \quad \left\{ \begin{array}{l} \frac{\partial^2 y}{\partial t^2} - M \left(\int_{\Omega} |\nabla y|^2 dx \right) \Delta y - \frac{\partial}{\partial t} \Delta y = 0 \text{ in } Q = \Omega \times (0, \infty) \\ y = 0 \text{ on } \Sigma_1 = \Gamma_1 \times (0, \infty) \\ M \left(\int_{\Omega} |\nabla y|^2 dx \right) \frac{\partial y}{\partial \nu} + \frac{\partial}{\partial t} \left(\frac{\partial y}{\partial \nu} \right) + |y|^\gamma y = 0 \text{ on } \Sigma_0 = \Gamma_0 \times (0, \infty) \\ y(0) = y^0, \frac{\partial y}{\partial t}(0) = y^1 \text{ in } \Omega. \end{array} \right.$$

where M is a C^1 function; $M(\lambda) \geq 0; \forall \lambda \geq 0$.

1 Introduction

Let Ω be a bounded domain of \mathbf{R}^n and let Γ denote its C^2 boundary. Assume that Γ consists of two parts, Γ_0 and Γ_1 , with positive measure and such that Γ_0 and Γ_1 are closed and disjoint. Let ν be the unit normal vector pointing towards the exterior of Ω and let $\frac{\partial}{\partial \nu}$ denote the normal derivative.

Let $M \in C^1([0, \infty), \mathbf{R})$ be a function which verifies

$$M(\lambda) \geq 0; \quad \forall \lambda \geq 0 \quad . \quad (1.1)$$

2000 Mathematics Subject Classification: 35B40, 35L80.
Servicio de Publicaciones. Universidad Complutense. Madrid, 2001

The aim of this paper is to prove existence and uniqueness of strong and weak solutions and to show that these solutions decay to zero uniformly when t goes to infinity.

The motivation for this problem comes from the small vibrations of an elastic string whose mathematical model, c.f. J. L. Lions [12], is given by the following Kirchhoff-Carrier equation

$$\rho h \frac{\partial^2 y}{\partial t^2} = \left(P_0 + \frac{Eh}{2L} \int_0^L \left| \frac{\partial y}{\partial x}(x, t) \right|^2 dx \right) \frac{\partial^2 y}{\partial x^2}; \quad 0 \leq x \leq L, \quad t \geq 0.$$

Here y is the lateral displacement at the space coordinate x and the time t , ρ the mass density, h the cross-section area, L the length, E the Young's modulus and P_0 the initial axial tension.

When $n = 1$, for instance, problem (*) describes the nonlinear vibrations of an one dimensional string which is clamped in one end and is free in the other end in the sense that it can move up and down subject to nonlinear effects caused by the function $f(y) = |y|^\gamma y$.

Hypothesis $M(0) = 0$ and $M(\lambda) > 0$ in a neighbourhood of the origin (see assumption (2.5) below) raised from the fact that we are interested in the case $P_0 = 0$.

A lot of papers have been devoted to this context. In this direction we can cite, for instance, Menzala [17], Arosio-Spagnolo [1], Rivera [21] and Ebihara-Milla Miranda-Medeiros [5]. Let us also mention the results of Yamada [24, 25], Vasconcellos-Teixeira [23], Muñoz Rivera [18] and Medeiros-Milla Miranda [14] which are in connection with damped problems. It is interesting to observe that problems without viscosity, that is, when $\Delta y' = 0$ and assuming that $M = 1$ and a feedback occurs on the boundary were studied by many authors, see Quinn and Russel [20], Chen [4], Lagnese [8,9], Komornik and Zuazua [7], Lasiecka and Tataru [10] and Cavalcanti, Domingos Cavalcanti and Soriano [2].

In spite of the importance of the subject, there are relatively few mathematical results in the presence of nonlinear boundary conditions combined with the nonlinearity due to the function M . To deal with this kind of problem is the contribution of this paper. Comparing the present paper with the problems considered in [11] and [16], where the term Δy_t was not added, we observe that in both cases the function M is not supposed to degenerate which contrasts with assumption (1.1).

In order to obtain the existence of solutions in [3], the authors employ Galerkin's method and, as they are not allowed to use basis formed by eigenfunctions, make use of arguments which are only valid when the only nonlinearity is given by $M = M(t)$. However, in the present paper, we have to deal with a nonlinear boundary condition and the degeneracy of the function M , which do not permit us to repeat the same arguments used in [3]. To overcome these difficulties is the goal of our work.

To obtain the uniform decay rates of the energy (c.f. Theorem 2.1 below)

$$E(t) = \frac{1}{2} \left[\int_{\Omega} |y'(x, t)|^2 dx + \hat{M} \left(\int_{\Omega} |\nabla y(x, t)|^2 dx \right) \right] + \frac{1}{\gamma+2} \int_{\Gamma_0} |y(x, t)|^{\gamma+2} d\Gamma \quad (1.3)$$

where

$$\hat{M}(\lambda) = \int_0^\lambda M(s) ds \quad (1.4)$$

we use the perturbed energy method, see Zuazua [7,26], combined with techniques from Muñoz Rivera [18].

Our paper is organized as follows. In section 2 we give the notations and state the main result. In section 3 we study existence and uniqueness of strong and weak solutions while in section 4 we obtain the uniform decay rates for solutions obtained in section 3.

2 Notations and main results

In this section we present some notations which will be used throughout this paper and will state the main result.

Let

$$V = \{v \in H^1(\Omega); v = 0 \text{ on } \Gamma_1\}$$

which endowed with the topology given by the norm $|\nabla \cdot|_{L^2(\Omega)}$ is a Hilbert subspace of $H^1(\Omega)$.

We denote

$$(u, v) = \int_{\Omega} u(x)v(x) dx; \quad (u, v)_{\Gamma_0} = \int_{\Gamma_0} u(x)v(x) dx, \\ |u|^2 = \int_{\Omega} |u(x)|^2 dx, \quad |u|_{\Gamma_0}^2 = \int_{\Gamma_0} |u(x)|^2 d\Gamma, \quad \|u\|_{p, \Gamma_0}^p = \int_{\Gamma_0} |u(x)|^p d\Gamma$$

and assume that

$$0 < \gamma \leq \frac{1}{n-2}; \quad n \geq 3 \quad \text{or} \quad \gamma > 0 \quad \text{if} \quad n = 1, 2. \quad (2.1)$$

Now, we are in position to state our main results.

Theorem 2.1. *Assume that (1.1) holds and*

$$\{y^0, y^1\} \in V^2. \quad (2.2)$$

Then, the problem () possesses a unique strong solution satisfying*

$$y \in L^\infty(0, \infty; V), \quad y' \in L^\infty(0, \infty; V); \quad y'' \in L^2(0, \infty; L^2(\Omega)). \quad (2.3)$$

Besides, the following energy decay holds

$$E(t) \leq C(1+t)^{-1}, \quad \forall t \geq 0, \quad (2.4)$$

where C is a positive constant.

Assuming that M is non decreasing and, for some $\alpha > 0$, satisfies the assumptions

$$\liminf_{s \rightarrow 0^+} \frac{M(s)}{s^\alpha} > 0 \quad \text{and} \quad M(s) = O(s^\alpha), \quad s \rightarrow 0^+, \quad (2.5)$$

there exist positive constants C and N such that

$$E(t) + \frac{1}{2} |\nabla y(t)|^2 \leq C(1+t)^{-\frac{1}{\alpha}}, \quad \forall t \geq N \quad (2.6)$$

and

$$M(|\nabla y(t)|^2) \leq C(1+t)^{-1}, \quad t \geq N. \quad (2.7)$$

Moreover, under the hypothesis

$$M(\lambda) \geq \lambda_0 > 0 \quad \text{for all} \quad \lambda > 0 \quad (2.8)$$

we obtain the exponential decay, that is, there exist positive constants C, θ such that

$$E(t) \leq C \exp(-\theta t), \quad t \geq 0. \quad (2.9)$$

Remark. The assumption (2.5) means that the function $M(s) > 0$ and it is bounded by a polynomial $P(s) = k s^\alpha$ ($k > 0$) in a neighbourhood of

the origin. Consequently $M(0) = 0$ and the algebraic decay rates occur according to the exponent α of the polynomial $P(s)$ which dominates the function $M(s)$.

Theorem 2.2. *Let*

$$\{y^0, y^1\} \in V \times L^2(\Omega) \quad (2.10)$$

and assume that $M \in C^1(0, \infty)$. Then, problem () possesses a unique weak solution in the class*

$$y \in C^0([0, \infty); V); \quad y' \in C^0([0, \infty); L^2(\Omega)) \quad (2.11)$$

and verify the decay rate given in (2.4).

Moreover, if M is non decreasing and satisfies the assumptions given in (2.5), we have the decay rates obtained in (2.6) and (2.7) in the degenerate case and the decay given by (2.9) in the nondegenerate one.

3 Existence and uniqueness of solutions

Let us consider initially

$$\{y^0, y^1\} \in V^2. \quad (3.1)$$

The variational formulation associated with problem (*) is given by

$$\begin{aligned} (y''(t), w) + M(|\nabla y(t)|^2) (\nabla y(t), \nabla w) + (\nabla y'(t), \nabla w) \\ + (|y(t)|^\gamma y(t), w)_{\Gamma_0} = 0; \quad \text{for all } w \in V. \end{aligned} \quad (3.2)$$

We represent by $(\omega_j)_{j \in \mathbb{N}}$ a basis in V , which is orthonormal in $L^2(\Omega)$, and by V_m the subspace of V generate by the m -first vectors $\omega_1, \dots, \omega_m$. Let us define

$$y_m(t) = \sum_{i=1}^m g_{im}(t) \omega_i, \quad (3.3)$$

where $y_m(t)$ is the solution of the following Cauchy problem

$$\begin{aligned} (y_m''(t), \omega_j) + M(|\nabla y_m(t)|^2) (\nabla y_m(t), \nabla \omega_j) + (\nabla y_m'(t), \nabla \omega_j) \\ + (|y_m(t)|^\gamma y_m(t), \omega_j)_{\Gamma_0} = 0, \end{aligned} \quad (3.4)$$

$$y_m(0) = y_{0m} \rightarrow y^0 \text{ in } V \quad \text{and} \quad y_m'(0) = y_{1m} \rightarrow y^1 \text{ in } V. \quad (3.5)$$

The above approximate system is a normal one of ordinary differential equations and has a solution in $[0, t_m)$. The extension of the solution to the whole interval $[0, T]$ is a consequence of the first estimate which we are going to obtain below.

3.1 A priori estimates

The first estimate.

Multiplying (3.4) by $g'_{jm}(t)$ and summing over j from (1.4) we have

$$\frac{d}{dt} \left\{ \frac{1}{2} |y'_m(t)|^2 + \frac{1}{2} \hat{M} (|\nabla y_m(t)|^2) + \frac{1}{\gamma+2} \|y_m(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2} \right\} + |\nabla y'_m(t)|^2 = 0. \quad (3.6)$$

Integrating (3.6) over $(0, t)$ we get

$$\begin{aligned} \frac{1}{2} |y'_m(t)|^2 + \frac{1}{2} \hat{M} (|\nabla y_m(t)|^2) + \frac{1}{\gamma+2} \|y_m(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2} + \int_0^t |\nabla y'_m(s)|^2 ds \\ = \frac{1}{2} |y'_{1m}|^2 + \frac{1}{2} \hat{M} (|\nabla y_{0m}|^2) + \frac{1}{\gamma+2} \|y_{0m}\|_{\gamma+2, \Gamma_0}^{\gamma+2}. \end{aligned} \quad (3.7)$$

From (3.5) and (3.7) we obtain the first estimate

$$|y'_m(t)|^2 + \hat{M} (|\nabla y_m(t)|^2) + \|y_m(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2} + \int_0^t |\nabla y'_m(s)|^2 ds \leq L_1, \quad (3.9)$$

where L_1 is a positive constant independent of $m \in \mathbf{N}$ and $t \geq 0$.

The second estimate.

Multiplying (3.4) by $g_{jm}(t)$ and summing over j we have

$$\begin{aligned} \frac{d}{dt} (y'_m(t), y_m(t)) + M (|\nabla y_m(t)|^2) |\nabla y_m(t)|^2 + \frac{1}{2} \frac{d}{dt} |\nabla y_m(t)|^2 \\ + \|y_m(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2} = |y'_m(t)|^2. \end{aligned} \quad (3.10)$$

Integrating (3.10) over $(0, t)$, taking (1.1), (3.5) into account and observing that there exists $C_0 > 0$ such that $|v| \leq C_0 |\nabla v|, \forall v \in V$, it follows that

$$\frac{1}{2} |\nabla y_m(t)|^2 + \int_0^t \|y_m(s)\|_{\gamma+2, \Gamma_0}^{\gamma+2} ds$$

$$\leq k_0 + C_0 |y'_m(t)| |\nabla y_m(t)| + C_0^2 \int_0^t |\nabla y'_m(s)|^2 ds,$$

where k_0 is a positive constant, that is,

$$\begin{aligned} & \frac{1}{4} |\nabla y_m(t)|^2 + \int_0^t \|y_m(s)\|_{\gamma+2, \Gamma_0}^{\gamma+2} ds \\ & \leq k_0 + \frac{C_0^2}{2} |y'_m(t)|^2 + C_0^2 \int_0^t |\nabla y'_m(s)|^2 ds. \end{aligned} \quad (3.11)$$

From (3.11) and considering the first estimate we obtain the second one

$$|\nabla y_m(t)|^2 + \int_0^t \|y_m(s)\|_{\gamma+2, \Gamma_0}^{\gamma+2} ds \leq L_2, \quad (3.12)$$

where L_2 is a positive constant independent of $m \in \mathbf{N}$ and $t \geq 0$.

The third estimate.

Multiplying (3.4) by $g''_{jm}(t)$ and summing over $(0, t)$ we have

$$\begin{aligned} & |y''_m(t)|^2 + M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y''_m(t)) + \frac{1}{2} \frac{d}{dt} |\nabla y'_m(t)|^2 \\ & + (|y_m(t)|^\gamma y_m(t), y''_m(t))_{\Gamma_0} = 0. \end{aligned} \quad (3.13)$$

Estimate for $I_1 := M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y''_m(t))$.

We have

$$\begin{aligned} I_1 &= \frac{d}{dt} \left[M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y'_m(t)) \right] \\ & - 2M' \left(|\nabla y_m(t)|^2 \right) [(\nabla y_m(t), \nabla y'_m(t))]^2 - M \left(|\nabla y_m(t)|^2 \right) |\nabla y'_m(t)|^2. \end{aligned}$$

Considering that $M \in C^1(\mathbf{R}_+)$ and the inequality (3.12), we obtain

$$\begin{aligned} & \left| 2M' \left(|\nabla y_m(t)|^2 \right) [(\nabla y_m(t), \nabla y'_m(t))]^2 \right| \\ & \leq 2 \left| M' \left(|\nabla y_m(t)|^2 \right) \right| |\nabla y_m(t)|^2 |\nabla y'_m(t)|^2 \\ & \leq C_1 |\nabla y'_m(t)|^2 \end{aligned}$$

and

$$\left| M \left(|\nabla y_m(t)|^2 \right) |\nabla y'_m(t)|^2 \right| \leq C_2 |\nabla y'_m(t)|^2.$$

Then,

$$\begin{aligned} & |y''_m(t)|^2 + \frac{d}{dt} \left[M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y'_m(t)) \right] \\ & + \frac{1}{2} \frac{d}{dt} |\nabla y'_m(t)|^2 + \frac{d}{dt} (|y_m(t)|^\gamma y_m(t), y'_m(t))_{\Gamma_0} \\ & - (\gamma + 1) (|y_m(t)|^\gamma y'_m(t), y'_m(t))_{\Gamma_0} \leq C_3 |\nabla y'_m(t)|^2, \end{aligned} \quad (3.14)$$

where C_3 is the positive constant $C_1 + C_2$.

Estimate for I_2 : $I_2 := (\gamma + 1) (|y_m(t)|^\gamma y'_m(t), y'_m(t))_{\Gamma_0}$.

Noting that $\frac{\gamma}{2(\gamma+1)} + \frac{1}{2(\gamma+1)} + \frac{1}{2} = 1$, the generalized Hölder inequality yields

$$|I_2| \leq (\gamma + 1) \| |y_m(t)|^\gamma \|_{2(\gamma+1), \Gamma_0} \| y'_m(t) \|_{2(\gamma+1), \Gamma_0} \| y'_m(t) \|_{\Gamma_0}.$$

Now, since $V \hookrightarrow L^{2(\gamma+1)}(\Gamma_0)$ then, from the last inequality and considering the second estimate we conclude

$$|I_3| \leq (\gamma + 1) |\nabla y_m(t)|^\gamma |\nabla y'_m(t)| |\nabla y'_m(t)| \leq C_4 |\nabla y'_m(t)|^2, \quad (3.15)$$

where C_4 is a positive constant.

Combining (3.14) and (3.15) and integrating the obtained result over $(0, t)$ we deduce

$$\begin{aligned} & \int_0^t |y''_m(s)|^2 ds + 2M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y'_m(t)) \\ & - 2M \left(|\nabla y_{0m}|^2 \right) (\nabla y_{0m}, \nabla y_{1m}) + |\nabla y'_m(t)|^2 - |\nabla y_{1m}|^2 \\ & \leq 2C_5 \int_0^t |\nabla y'_m(s)|^2 ds - 2 (|y_m(t)|^\gamma y_m(t), y'_m(t))_{\Gamma_0} + 2 (|y_{0m}|^\gamma y_{0m}, y_{1m})_{\Gamma_0} \end{aligned} \quad (3.16)$$

where C_5 is a positive constant.

Estimate for I_3 : $I_3 := -2 (|y_m(t)|^\gamma y_m(t), y'_m(t))_{\Gamma_0}$.

From Cauchy-Schwarz inequality, making use of the inequality $ab \leq \frac{1}{4\eta}a^2 + \eta b^2$ ($\eta > 0$ arbitrary) and considering the second estimate we obtain k_1 and $C_6(\eta)$ positive constants such that

$$\begin{aligned} |I_3| &\leq 2 \int_{\Gamma_0} |y_m|^{\gamma+1} |y'_m| d\Gamma & (3.17) \\ &\leq 2 \|y_m(t)\|_{2(\gamma+1), \Gamma_0}^{\gamma+1} \|y'_m(t)\|_{\Gamma_0} \\ &\leq k_1 |\nabla y_m(t)|^{\gamma+1} |\nabla y'_m(t)| \\ &\leq C_6(\eta) + \eta |\nabla y'_m(t)|^2. \end{aligned}$$

Thus, from (3.5), (3.16) and (3.17) there exists a positive constant $C_7(\eta)$ such that

$$\begin{aligned} \int_0^t |y''_m(s)|^2 ds + 2M \left(|\nabla y_m(t)|^2 \right) (\nabla y_m(t), \nabla y'_m(t)) + (1-\eta) |\nabla y'_m(t)|^2 \\ \leq C_7(\eta) + 2C_5 \int_0^t |\nabla y'_m(s)|^2 ds. \end{aligned}$$

Consequently, from the second estimate and from the last inequality we conclude

$$\begin{aligned} \int_0^t |y''_m(s)|^2 ds + (1-2\eta) |\nabla y'_m(t)|^2 & (3.18) \\ \leq C_8(\eta) + 2C_5 \int_0^t |\nabla y'_m(s)|^2 ds. \end{aligned}$$

where C_8 is a positive constant.

Finally from (3.18), choosing η sufficiently small and making use of Gronwall's lemma we obtain the third estimate

$$\int_0^t |y''_m(s)|^2 ds + |\nabla y'_m(t)|^2 \leq L_3, \quad (3.19)$$

where L_3 is a positive constant independent of $m \in \mathbf{N}$ and $t \in [0, T]$.

From the estimate (3.19) we can extract a subsequence (y_μ) of (y_m) such that

$$y''_\mu \rightharpoonup y'' \quad \text{weakly in } L^2(0, T; L^2(\Omega)) \quad (3.20)$$

and

$$y'_\mu \rightharpoonup y' \quad \text{weakly star in } L^\infty(0, T; V). \quad (3.21)$$

3.2 Analysis of the nonlinear terms

From now on we are interested in the convergence of the nonlinear terms.

We define:

$$\varphi_m(t) = |\nabla y_m(t)|^2; \quad t \in [0, T]. \quad (3.22)$$

From the second estimate we have

$$|\varphi_m(t)| \leq C; \quad \forall m \in \mathbf{N}, \quad \forall t \in [0, T], \quad (3.23)$$

where C is a positive constant.

Now, if $t_1, t_2 \in [0, T]$ we obtain

$$|\varphi_m(t_1) - \varphi_m(t_2)| \leq \int_{t_1}^{t_2} |\varphi'_m(s)| ds. \quad (3.24)$$

On the other hand, from the second and third estimates we deduce

$$\varphi'_m(s) = 2 (\nabla y_m(s), \nabla y'_m(s)) \leq C; \quad \forall m \in \mathbf{N}, \quad \forall s \in [0, T]; \quad (3.25)$$

where C is a positive constant independent of m .

Combining (3.24) and (3.25) it follows that

$$|\varphi_m(t_1) - \varphi_m(t_2)| \leq C |t_1 - t_2|; \quad \forall m \in \mathbf{N}. \quad (3.26)$$

Then, from (3.23) and (3.26) by Arzela-Ascoli's theorem there exists a continuous function $\varphi : [0, T] \rightarrow \mathbf{R}$ such that

$$\varphi_m(t) \rightarrow \varphi(t) \quad (3.27)$$

uniformly in $[0, T]$ and, since $M \in C^1([0, \infty), \mathbf{R})$, we obtain

$$M(|\nabla y_m(t)|^2) \rightarrow M(\varphi(t)) \quad (3.28)$$

uniformly in $[0, T]$.

From the second estimate and noting that $V \hookrightarrow L^{2(\gamma+1)}(\Gamma_0)$ we deduce

$$\{|y_m|^\gamma y_m\} \text{ is bounded in } L^\infty(0, T; L^2(\Gamma_0)). \quad (3.29)$$

Also, we note that from the second and third estimates and noting that $\|v\|_{H^{1/2}(\Gamma_0)} \leq C |\nabla v|$, for all $v \in V$, we infer

$$\{y_m\} \text{ is bounded in } L^2(0, T; H^{1/2}(\Gamma_0)), \quad (3.30)$$

$$\{y'_m\} \text{ is bounded in } L^2(0, T; L^2(\Gamma_0)). \quad (3.31)$$

Since the injection $H^{1/2}(\Gamma_0) \hookrightarrow L^2(\Gamma_0)$ is continuous and compact, making use of Aubin-Lions theorem, see J. L. Lions [13, Theo. 5.1], there exist $\{y_\mu\}$ a subsequence of $\{y_m\}$ such that

$$y_\mu \rightarrow y \text{ strongly in } L^2(0, T; L^2(\Gamma_0)). \quad (3.32)$$

From now on all the eventual subsequences are denoted by the same notation.

Therefore,

$$y_\mu \rightarrow y \text{ a.e. in } \Gamma_0 \times (0, T)$$

and consequently

$$|y_\mu|^\gamma y_\mu \rightarrow |y|^\gamma y \text{ a.e. in } \Gamma_0 \times (0, T). \quad (3.33)$$

Combining (3.29) and (3.33) we deduce, thanks to Lions' lemma, see J. L. Lions [13, lemma 1.3],

$$|y_\mu|^\gamma y_\mu \rightharpoonup |y|^\gamma y \text{ weakly in } L^2(0, T; L^2(\Gamma_0)). \quad (3.34)$$

Considering the above convergences we can pass to the limit in the approximate system given by (3.4) using standard arguments in order to obtain

$$y'' - M(\varphi(t)) \Delta y - \Delta y' = 0 \quad \text{in } L^2(0, T; L^2(\Omega)), \quad (3.35)$$

$$y(0) = y^0; \quad y'(0) = y^1.$$

Moreover, making use of the generalized Green formula we deduce

$$M(\varphi(t)) \frac{\partial y}{\partial \nu} + \frac{\partial y_t}{\partial \nu} + |y|^\gamma y = 0 \quad \text{in } L^2(0, T; L^2(\Gamma_0)). \quad (3.36)$$

Our aim is to show that

$$\varphi(t) = |\nabla y(t)|^2; \quad t \in [0, T]. \quad (3.37)$$

Indeed, multiplying (3.4) by g_{jm} , summing over j and integrating over $[0, t]$ we obtain

$$\int_0^t (y''_m(s), y_m(s)) ds + \int_0^t M(|\nabla y_m(s)|^2) |\nabla y_m(s)|^2 ds \quad (3.38)$$

$$+ \int_0^t (|y_m(s)|^\gamma y_m(s), y_m(s))_{\Gamma_0} ds + \int_0^t (\nabla y'_m(s), \nabla y_m(s)) ds = 0.$$

On the other hand, from the first and second estimates and making use of Aubin-Lions theorem one has

$$y_m \rightarrow y \quad \text{strongly in } L^2(0, T; L^2(\Omega)). \quad (3.39)$$

Considering the convergences given by (3.20), (3.27), (3.28), (3.32), (3.34), (3.39) we can pass to the limit in (3.38) to obtain

$$\begin{aligned} & \lim_{m \rightarrow \infty} \int_0^t (\nabla y'_m(s), \nabla y_m(s)) ds \quad (3.40) \\ &= - \int_0^t (y''(s), y(s)) ds - \int_0^t M(\varphi(s)) \varphi(s) ds - \int_0^t (|y(s)|^\gamma y(s), y(s))_{\Gamma_0} ds. \end{aligned}$$

Combining (3.35), (3.36) and (3.40) and taking the generalized Green formula into account we deduce

$$\begin{aligned} & \lim_{m \rightarrow \infty} \int_0^t (\nabla y'_m(s), \nabla y_m(s)) ds \\ &= \int_0^t M(\varphi(s)) \left[|\nabla y(s)|^2 - \varphi(s) \right] ds + \int_0^t (\nabla y'(s), \nabla y(s)) ds. \end{aligned}$$

The last equality yields

$$\begin{aligned} & \lim_{m \rightarrow \infty} \left[\frac{1}{2} |\nabla y_m(t)|^2 - \frac{1}{2} |\nabla y_{0m}|^2 \right] \quad (3.41) \\ &= \int_0^t M(\varphi(s)) \left[|\nabla y(s)|^2 - \varphi(s) \right] ds + \frac{1}{2} |\nabla y(t)|^2 - \frac{1}{2} |\nabla y(0)|^2. \end{aligned}$$

Then, from (3.41) and considering the convergences in (3.5) and (3.27) we conclude

$$\begin{aligned} \left| \varphi(t) - |\nabla y(t)|^2 \right| &\leq 2 \int_0^t M(\varphi(s)) \left| |\nabla y(s)|^2 - \varphi(s) \right| ds \\ &\leq C \int_0^t \left| |\nabla y(s)|^2 - \varphi(s) \right| ds, \end{aligned}$$

where C is a positive constant. Employing Gronwall's lemma, from the last inequality we obtain the desired in (3.37).

We observe that for each fixed t in $[0, \infty)$, we have that $y(t)$ is the weak solution of the Dirichlet-Neumann problem

$$\begin{cases} -\Delta \left[M \left(|\nabla y|^2 \right) y + y' \right] = -y'' & \text{in } L^2(0, \infty; L^2(\Omega)) \\ M \left(|\nabla y|^2 \right) y + y' = 0 & \text{on } \Sigma_1 \\ \frac{\partial}{\partial \nu} \left[M \left(|\nabla y|^2 \right) y + y' \right] + |y|^\gamma y = 0 & \text{in } L^2(0, \infty; L^2(\Gamma_0)). \end{cases}$$

Then, the theory of elliptic problems gives

$$M \left(|\nabla y|^2 \right) y + y' \in L^2(0, \infty; V \cap H^{3/2}(\Omega)). \tag{3.42}$$

3.3 Uniqueness

Let y and \hat{y} be two solutions of problem (*). Denoting $z = y - \hat{y}$, it comes from (3.35) and (3.36) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} |z'(t)|^2 + M \left(|\nabla y(t)|^2 \right) (\nabla y(t), \nabla z'(t)) \\ & - M \left(|\nabla \hat{y}(t)|^2 \right) (\nabla \hat{y}(t), \nabla z'(t)) + |\nabla z'(t)|^2 \\ & + (|y(t)|^\gamma y(t), z'(t))_{\Gamma_0} - (|\hat{y}(t)|^\gamma \hat{y}(t), z'(t))_{\Gamma_0} = 0. \end{aligned} \tag{3.43}$$

Summing and subtracting the term $M \left(|\nabla y(t)|^2 \right) (\nabla \hat{y}(t), \nabla z'(t))$ in (3.43) we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} |z'(t)|^2 + M \left(|\nabla y(t)|^2 \right) (\nabla z(t), \nabla z'(t)) \\ & + |\nabla z'(t)|^2 = \left(M \left(|\nabla \hat{y}(t)|^2 \right) - M \left(|\nabla y(t)|^2 \right) \right) (\nabla \hat{y}(t), \nabla z'(t)) \\ & + (|\hat{y}(t)|^\gamma \hat{y}(t) - |y(t)|^\gamma y(t), z'(t))_{\Gamma_0}. \end{aligned} \tag{3.44}$$

On the other hand, since M is C^1 we get

$$\left| M \left(|\nabla \hat{y}(t)|^2 \right) - M \left(|\nabla y(t)|^2 \right) \right| \tag{3.45}$$

$$\begin{aligned}
&\leq k_1 \left| |\nabla y(t)|^2 - |\nabla \hat{y}(t)|^2 \right| \\
&\leq k_1 [|\nabla y(t)| + |\nabla \hat{y}(t)|] |\nabla y(t) - \nabla \hat{y}(t)| \\
&\leq k_2 |\nabla z(t)|
\end{aligned}$$

where k_1 and k_2 are positive constants.

Combining (3.44) and (3.45) and using analogous considerations like those ones used in the third estimate (see estimate for I_3) it follows that

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} |z'(t)|^2 + |\nabla z'(t)|^2 \tag{3.46} \\
&\leq C_1 |\nabla z(t)| |\nabla z'(t)| + C_2(\gamma) \int_{\Gamma_0} (|\hat{y}|^\gamma + |y|^\gamma) |z| |z'| d\Gamma \\
&\leq C_3 |\nabla z(t)| |\nabla z'(t)| \\
&\leq \frac{C_3^2}{2} |\nabla z(t)|^2 + \frac{1}{2} |\nabla z'(t)|^2.
\end{aligned}$$

Integrating (3.46) over $(0, t)$ we obtain

$$\begin{aligned}
&\frac{1}{2} |z'(t)|^2 + \frac{1}{2} \int_0^t |\nabla z'(s)|^2 ds \tag{3.47} \\
&\leq \frac{C_3^2}{2} \int_0^t |\nabla z(s)|^2 ds.
\end{aligned}$$

Since

$$|\nabla z(t)|^2 \leq T \int_0^t |\nabla z'(s)|^2 ds \tag{3.48}$$

combining (3.47) and (3.48) and applying Gronwall's lemma we obtain $|\nabla z(t)| = |z'(t)| = 0$. This concludes the uniqueness of strong solutions.

3.4 Solvability of weak solutions

We have just proved the existence of solutions to problem (*) when $\{y^0, y^1\} \in V^2$. However, when $\{y^0, y^1\} \in V \times L^2(\Omega)$, by density arguments and using analogous considerations like those ones used in the first and second estimates and in the uniqueness, we can find a sequence $\{y_\mu\}$ of solutions to problem (*) and a function $y : \Omega \times (0, T) \rightarrow \mathbf{R}$ such that

$$y_\mu \in L^\infty(0, \infty; V), \quad y'_\mu \in L^\infty(0, \infty; V), \quad y''_\mu \in L^2(0, \infty; L^2(\Omega)),$$

$$y_\mu \rightarrow y \text{ strongly in } C^0([0, T]; V) \quad (3.49)$$

$$y'_\mu \rightarrow y' \text{ strongly in } C^0([0, T]; L^2(\Omega)) \quad (3.50)$$

$$y'_\mu \rightarrow y' \text{ strongly in } L^2(0, T; V) \quad (3.51)$$

$$y'_\mu \rightharpoonup y' \text{ weakly star in } L^\infty(0, T; V) \quad (3.52)$$

and

$$y''_\mu - M \left(|\nabla y_\mu|^2 \right) \Delta y_\mu - \Delta y'_\mu = 0 \text{ in } L^2(0, T; L^2(\Omega)).$$

Consider $\theta \in D(0, T)$ and $\phi \in D(\Omega)$. Then,

$$\begin{aligned} - \int_0^T (y'_\mu(t), \phi) \theta'(t) dt + \int_0^T M \left(|\nabla y_\mu(t)|^2 \right) (\nabla y_\mu(t), \nabla \phi) \theta(t) dt \\ + \int_0^T (\nabla y'_\mu(t), \nabla \phi) \theta(t) dt = 0. \end{aligned}$$

The above convergences are sufficient to pass to the limit in the above equality in order to obtain

$$\begin{aligned} - \int_0^T (y'(t), \phi) \theta'(t) dt + \int_0^T M \left(|\nabla y(t)|^2 \right) (\nabla y(t), \nabla \phi) \theta(t) dt \\ + \int_0^T (\nabla y'(t), \nabla \phi) \theta(t) dt = 0, \end{aligned}$$

for all $\theta \in D(0, T)$ and $\phi \in D(\Omega)$.

Consequently,

$$\langle -\langle y', \theta' \rangle, \phi \rangle - \left\langle \left\langle M \left(|\nabla y|^2 \right) \Delta y, \theta \right\rangle, \phi \right\rangle - \langle \langle \Delta y', \theta \rangle, \phi \rangle = 0,$$

for all $\theta \in D(0, T)$, $\phi \in D(\Omega)$; where $\langle \cdot, \theta \rangle$ and $\langle \cdot, \phi \rangle$ represent the duality $D'(0, T) \times D(0, T)$ and $D'(\Omega) \times D(\Omega)$, respectively.

Then,

$$\langle y'', \theta \phi \rangle - \left\langle M \left(|\nabla y|^2 \right) \Delta y, \theta \phi \right\rangle - \langle \Delta y', \theta \phi \rangle = 0,$$

for all $\theta \in D(0, T)$, $\phi \in D(\Omega)$; that is,

$$y'' - M(|\nabla y|^2) \Delta y - \Delta y' = 0 \quad \text{in } D'(\Omega \times]0, T[),$$

since $\{\theta\phi; \theta \in D(0, T), \phi \in D(\Omega)\}$ is dense in $D(\Omega \times]0, T[)$.

However, $\Delta y, \Delta y' \in L^2(0, T; V')$, which implies that

$$y'' \in L^2(0, T; V'),$$

provided that $y'' = M(|\nabla y|^2) \Delta y + \Delta y'$ in $D'(\Omega \times]0, T[)$. Then, we obtain a weak solution y to (*) in the class

$$\begin{aligned} y &\in C^0([0, \infty); V); y' \in C^0([0, \infty); \\ &L^2(\Omega)) \cap L^2(0, \infty; V); y'' \in L^2(0, \infty; V') \end{aligned}$$

verifying

$$y'' - M(|\nabla y|^2) \Delta y - \Delta y' = 0 \quad \text{in } L^2(0, \infty; V'). \quad (3.53)$$

3.5 Characterization of the boundary condition

Let us consider the elliptic problem

$$\left\{ \begin{array}{l} -\Delta p = y' \quad \text{in } \Omega \\ p = 0 \quad \text{on } \Gamma_1 \\ \frac{\partial p}{\partial \nu} = \int_0^t |y(s)|^\gamma y(s) ds \quad \text{on } \Gamma_0 \end{array} \right. \quad (3.54)$$

where y is the weak solution of (*) verifying (3.53). Taking the regularity of Γ into account, we have

$$p \in L^2(0, \infty; \mathcal{H}) \quad (3.55)$$

where

$$\mathcal{H} = \{u \in V; \Delta u \in L^2(\Omega)\}.$$

Next, we are going to prove that

$$M(|\nabla y|^2) y + y' = -p' \quad \text{in } H^{-1}(0, \infty; \mathcal{H}). \quad (3.56)$$

Indeed, from (3.53) we can write

$$-\Delta \left[M(|\nabla y|^2) y + y' \right] = -y'' \quad \text{in } L^2_{loc}(0, \infty; V')$$

and from (3.56) we obtain

$$-\Delta \left[M \left(|\nabla y|^2 \right) y + y' \right] = \Delta p' \text{ in } D'_{loc}(0, \infty; V').$$

Therefore

$$\left\langle -\Delta \left[M \left(|\nabla y|^2 \right) y + y' \right], \theta \right\rangle = \langle -\Delta p, \theta' \rangle \text{ in } V', \text{ for all } \theta \in D(0, \infty).$$

Hence,

$$\int_0^T (-\Delta) \left(M \left(|\nabla y|^2 \right) y + y' \right) (t) \theta(t) dt = \int_0^T (-\Delta) p(t) \theta'(t) dt \text{ in } V'$$

and consequently

$$\Delta \int_0^T \left(M \left(|\nabla y|^2 \right) y + y' \right) \theta(t) dt = \Delta \int_0^T p(t) \theta'(t) dt \text{ in } V'. \quad (3.57)$$

In order to prove (3.56) it remains to prove that the operator $-\Delta : V \rightarrow V'$ is an injective map. In fact, if this affirmative is true we conclude from (3.57) that

$$\int_0^T \left(M \left(|\nabla y|^2 \right) y + y' \right) \theta(t) dt = \int_0^T p(t) \theta(t) dt \text{ in } V,$$

for all $\theta \in D(0, \infty)$, and, consequently, (3.56) follows from the above inequality and from (3.55). To prove the desired result, let us define the following

$$a(u, v) = \int \nabla u \cdot \nabla v dx; \text{ for all } u, v \in V$$

$$L(v) = \int_{\Gamma_0} g v d\Gamma + \int_{\Omega} f v dx; \text{ for all } v \in V,$$

where $g \in L^2(\Gamma_0)$ and $f \in L^2(\Omega)$. From the above definitions we conclude from Lax-Milgran's lemma that there exists a unique $u \in V$ such that

$$a(u, v) = L(v), \text{ for all } v \in V.$$

In other words, there exists a unique $u \in V$ verifying

$$\left\{ \begin{array}{l} -\Delta u = f \quad \text{in } \Omega \\ u = 0 \quad \text{on } \Gamma_1 \\ \frac{\partial u}{\partial \nu} = g \quad \text{on } \Gamma_0. \end{array} \right.$$

Then, given $\{f, g\} \in L^2(\Omega) \times L^2(\Gamma_0)$, the solution of the above elliptic problem is uniquely determined, which proves that the operator $-\Delta : V \rightarrow V'$ is an injective map.

In what follows we are going to consider the dual trace operators

$$\hat{\gamma}_0 : H_{loc}^{-1}(0, \infty; V) \rightarrow H_{loc}^{-1}(0, \infty; L^2(\Gamma_0)), \tag{3.58}$$

$$\hat{\gamma}_1 : H_{loc}^{-1}(0, \infty; \mathcal{H}) \rightarrow H_{loc}^{-1}(0, \infty; H^{-1/2}(\Gamma_0)), \tag{3.59}$$

whose constructions and properties can be found in Milla Miranda [15].

From (3.56) we obtain

$$(\hat{\gamma}_1|_{\Sigma_0}) \left(M \left(|\nabla y|^2 \right) y + y' \right) = -(\hat{\gamma}_1|_{\Sigma_0}) (p') = -(\bar{\gamma}_1(p))', \tag{3.60}$$

where

$$\bar{\gamma}_1 : L_{loc}^2(0, \infty; \mathcal{H}) \rightarrow L_{loc}^2(0, \infty; H^{-1/2}(\Gamma_0))$$

is the trace operator given by

$$(\bar{\gamma}_1 u)(t) = \gamma_1(u(t))$$

and $\gamma_1 : \mathcal{H} \rightarrow H^{-1/2}(\Gamma_0)$ is the usual one related to the normal derivative.

Thus, from (3.54) and (3.60) we conclude

$$\begin{aligned} (\hat{\gamma}_1|_{\Sigma_0}) \left(M \left(|\nabla y|^2 \right) y + y' \right) &= - \left(\bar{\gamma}_0 \left(\int_0^t |y(s)|^\gamma y(s) ds \right) \right)' \\ &= -(\hat{\gamma}_0|_{\Sigma_0}) (|y|^\gamma y) \end{aligned} \tag{3.61}$$

where, analogously,

$$\bar{\gamma}_0 : L_{loc}^2(0, \infty; V) \rightarrow L_{loc}^2(0, \infty; L^2(\Gamma_0))$$

is the trace operator defined by

$$(\bar{\gamma}_0 u)(t) = \gamma_0(u(t)).$$

Hence, from (3.61) we can give meaning to the boundary condition. More precisely we have

$$\frac{\partial}{\partial \nu} \left(M \left(|\nabla y|^2 \right) y + y' \right) + |y|^\gamma y = 0 \quad \text{in } H_{loc}^{-1}(0, \infty; L^2(\Gamma_0)). \quad (3.62)$$

Since $|y|^\gamma y \in L^2(0, \infty; L^2(\Gamma_0))$ we have the boundary characterization valid a.e. on Σ_0 .

3.6 Uniqueness

According to (3.53) we have

$$y'' - M \left(|\nabla y|^2 \right) \Delta y - \Delta y' = 0 \quad \text{in } L^2(0, T; V').$$

Let y and \hat{y} be two weak solutions of problem (*). Defining $z = y - \hat{y}$ we have that $z \in H^1(0, T, V)$ and z satisfies

$$z'' - \Delta \left(M |\nabla y|^2 y - y' \right) + \Delta \left(M |\nabla \hat{y}|^2 \hat{y} - \hat{y}' \right) = 0 \quad \text{in } L^2(0, T; V').$$

Then, we are able to compose the above equation with z' , and following step by step the arguments done in the case of strong solutions, we conclude that $|\nabla z(t)| = |z'(t)| = 0$; which proves the uniqueness.

4 Asymptotic behaviour

In this section we are going to obtain the algebraic decay for strong solutions of (*). Using density arguments we obtain the same result for weak solutions.

The derivative of the energy defined in (1.3) is given by

$$E'(t) = - |\nabla y'(t)|^2. \quad (4.1)$$

Let us consider the perturbed energy

$$E_\varepsilon(t) = E(t) + \varepsilon \psi(t), \quad (4.2)$$

where

$$\psi(t) = (y'(t), y(t)) + \frac{1}{2} |\nabla y(t)|^2. \quad (4.3)$$

Taking the derivative of (4.2) with respect to t , replacing

$$(y''(t), y(t)) = -M \left(|\nabla y(t)|^2 \right) |\nabla y(t)|^2 - (\nabla y'(t), \nabla y(t)) - (|y(t)|^\gamma y(t), y(t))_{\Gamma_0}$$

in the obtained result and considering (4.1), it follows that

$$E'_\varepsilon(t) = E'(t) + \varepsilon \psi'(t) \quad (4.4)$$

$$= -|\nabla y'(t)|^2 - \varepsilon M \left(|\nabla y(t)|^2 \right) |\nabla y(t)|^2 - \varepsilon \|y(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2} + \varepsilon |y'(t)|^2.$$

Let μ be a positive constant such that

$$|v|^2 \leq \mu |\nabla v|^2, \quad \forall v \in V. \quad (4.5)$$

From (4.4) and (4.5) we obtain

$$E'_\varepsilon(t) \leq \left(-1 + \frac{3}{2}\varepsilon\mu\right) |\nabla y'(t)|^2 - \varepsilon M \left(|\nabla y(t)|^2 \right) |\nabla y(t)|^2 - \varepsilon \|y(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2}. \quad (4.6)$$

Besides, taking into account that M is a non decreasing function we have

$$\hat{M} \left(|\nabla y(t)|^2 \right) \leq M \left(|\nabla y(t)|^2 \right) |\nabla y(t)|^2$$

and therefore (4.6) yields

$$E'_\varepsilon(t) \leq -\left(1 - \frac{3}{2}\varepsilon\mu\right) |\nabla y'(t)|^2 - \varepsilon E(t). \quad (4.7)$$

Considering $\varepsilon \in (0, \varepsilon_1]$, where $\varepsilon_1 = \frac{2}{3\mu}$, we deduce

$$E'_\varepsilon(t) \leq -\varepsilon E(t) \quad (4.8)$$

and, consequently, integrating the last inequality over $[0, t]$ it follows that

$$\varepsilon \int_0^t E(s) ds \leq E_\varepsilon(0) - E_\varepsilon(t) \leq L, \quad (4.9)$$

where L is a positive constant which depends only on the initial data $\{y^0, y^1\}$.

Thus, from (4.9) we have

$$\int_0^t E(s)ds \leq \frac{L}{\varepsilon},$$

which implies that

$$E \in L^1(0, \infty). \quad (4.10)$$

On the other hand, since $E'(t) \leq 0$, we have

$$(1+t)E'(t) \leq 0, \quad \forall t \geq 0,$$

then

$$\frac{d}{dt} [(1+t)E(t)] \leq E(t). \quad (4.11)$$

Integrating (4.11) and taking (4.10) into account we obtain

$$E(t) \leq \left[E(0) + \int_0^\infty E(s)ds \right] (1+t)^{-1} \quad (4.12)$$

which concludes the desired in (2.4).

From now on we assume that hypothesis (2.5) hold. Then, we conclude the following result

Lemma 4.1. *We have*

$$\lim_{t \rightarrow \infty} |\nabla y(t)|^2 = 0. \quad (4.13)$$

Proof. Assume that (4.13) does not hold. Then, there exists $\varepsilon_0 > 0$ such that for every $n \in \mathbf{N}$ there exists t_n , with $t_n > n$ and

$$|\nabla y(t_n)|^2 \geq \varepsilon_0.$$

From the last inequality and since \hat{M} is a non decreasing function we have

$$\hat{M} \left(|\nabla y(t_n)|^2 \right) \geq \hat{M}(\varepsilon_0) \geq 0; \quad t_n > n. \quad (4.14)$$

Consequently from (4.12) and (4.14) we obtain

$$\hat{M}(\varepsilon_0) = 0. \quad (4.15)$$

On the other hand, from assumption (2.5) we have

$$\liminf_{s \rightarrow 0^+} \frac{M(s)}{s^\alpha} = l > 0,$$

which implies the existence of a real number $\delta_1 > 0$ such that

$$M(s) > \frac{l}{2} s^\alpha; \quad \forall s \in (0, \delta_1) \quad (4.16)$$

and

$$M(s) > 0, \quad \forall s \in (0, \delta_1). \quad (4.17)$$

Defining

$$\delta = \min\{\delta_1, \varepsilon_0\}$$

we have from (4.15) that

$$0 \leq \int_0^\delta M(\lambda) d\lambda \leq \int_0^{\varepsilon_0} M(\lambda) d\lambda = \hat{M}(\varepsilon_0) = 0,$$

that is,

$$\int_0^\delta M(\lambda) d\lambda = 0. \quad (4.18)$$

But (4.18) it is a contradiction with (4.17). This proves the lemma. ■

Next, we are going to prove that there exist C_1 and N positive constant such that

$$|\nabla y(t)|^{2(\alpha+1)} \leq C_1 \hat{M}(|\nabla y(t)|^2); \quad t > N. \quad (4.19)$$

Indeed, from (4.13) there exists $N > 0$ such that

$$|\nabla y(t)|^2 \leq \delta_1; \quad t > N. \quad (4.20)$$

On the other hand, integrating (4.16) over $(0, t)$ it follows that

$$\frac{l}{2} t^{\alpha+1} \leq \hat{M}(t), \quad \forall t \in (0, \delta_1). \quad (4.21)$$

Combining (4.20) and (4.21) we conclude the desired in (4.19). ■

From (4.2), (4.3) and (4.5) we can write

$$E_\varepsilon(t) \geq \left(\frac{1}{2} - \mu\varepsilon\right) |y'(t)|^2 + \frac{\varepsilon}{4} |\nabla y(t)|^2 + \frac{1}{2} \hat{M} \left(|\nabla y(t)|^2\right) \quad (4.22)$$

$$+ \frac{\varepsilon}{\gamma + 2} \|y(t)\|_{\gamma+2, \Gamma_0}^{\gamma+2}.$$

Then,

$$E_\varepsilon(t) \geq 0, \quad \forall \varepsilon \in (0, 1/2\mu]. \quad (4.23)$$

From (4.2), (4.3), (4.5), (4.12) and (4.19) we have for all $\varepsilon \leq \left\{\varepsilon_1, \frac{1}{2\mu}, 1\right\}$ and $t > N$ that

$$E_\varepsilon(t) \leq E(t) + \frac{\varepsilon}{2} |y'(t)|^2 + \frac{C_1\varepsilon}{2} (\mu + 1) \left[\hat{M} \left(|\nabla y(t)|^2\right)\right]^{\frac{1}{\alpha+1}}$$

$$\leq E(t) + \varepsilon E(t) + C_1\varepsilon(\mu + 1) [E(t)]^{\frac{1}{\alpha+1}}$$

$$\leq C_2 [E(t)]^{\frac{1}{\alpha+1}}$$

and therefore

$$[E_\varepsilon(t)]^{\alpha+1} \leq CE(t), \quad t > N. \quad (4.24)$$

Combining (4.8) and (4.24), it follows that

$$E'_\varepsilon(t) + \frac{\varepsilon}{C} E_\varepsilon^{\alpha+1}(t) \leq 0, \quad t > N. \quad (4.25)$$

Taking (4.23) into account and considering in (4.25) the change of variables

$$z_\varepsilon(t) = E_\varepsilon^{-\alpha}(t),$$

for a fixed and sufficiently small ε , we obtain

$$z'_\varepsilon(t) \geq \frac{\varepsilon\alpha}{C_2}.$$

Integrating the last inequality over $[0, t]$ we get

$$z_\varepsilon(t) \geq z_\varepsilon(0) + \frac{\varepsilon\alpha}{C_2} t$$

and, consequently,

$$E_\varepsilon^{-\alpha}(t) \geq C_3(1 + t). \quad (4.26)$$

From the last inequality we conclude that there exists a positive constant C , which depends on the initial data, such that

$$E_\varepsilon(t) \leq C(1+t)^{-\frac{1}{\alpha}}, \quad t > N. \quad (4.27)$$

This result combined with (4.22) proves (2.6). Finally, since $M(s) = O(s^\alpha)$, $s \rightarrow 0$ we deduce (2.7) from (2.6).

Finally, if we have $M(\lambda) \geq \lambda_0 > 0$, then from (4.2) and (4.3), there exists $\theta_1 > 0$ such that

$$|E_\varepsilon(t) - E(t)| \leq \varepsilon\theta_1 E(t). \quad (4.28)$$

Then, combining (4.8) and (4.28) we obtain the exponential decay mentioned in (2.9). The proof is now completed. ■

Further remarks

(1) Let $-\Delta$ the operator defined by the triple $\{V, L^2(\Omega), a(u, v)\}$ where

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v \, dx, \quad u, v \in V$$

and

$$D(-\Delta) = \left\{ u \in V \cap H^2(\Omega); \frac{\partial u}{\partial \nu} = 0 \text{ on } \Gamma_0 \right\}.$$

We recall that the Spectral Theorem for self-adjoint operators guarantees the existence of a complete orthonormal system (ω_ν) of $L^2(\Omega)$ given by the eigenfunctions of $-\Delta$. If (λ_ν) are the eigenvalues of $-\Delta$, then $\lambda_\nu \rightarrow +\infty$ as $\nu \rightarrow +\infty$. Now, since $-\Delta$ is positive, given $\alpha > 0$ one has

$$D[(-\Delta)^\alpha] = \left\{ u \in L^2(\Omega); \sum_{\nu=1}^{\infty} \lambda_\nu^{2\alpha} |(u, \omega_\nu)|^2 < \infty \right\}$$

and

$$(-\Delta)^\alpha u = \sum_{\nu=1}^{\infty} \lambda_\nu^\alpha (u, \omega_\nu) \omega_\nu, \quad \text{for all } u \in D[(-\Delta)^\alpha].$$

In $D[(-\Delta)^\alpha]$ we consider the topology given by

$$\|u\|_{D[(-\Delta)^\alpha]} = \|(-\Delta)^\alpha u\|_{L^2(\Omega)}.$$

We observe that such operators are self-adjoints, that is,

$$((-\Delta)^\alpha u, v) = (u, (-\Delta)^\alpha v); \quad \text{for all } u, v \in D [(-\Delta)^\alpha],$$

and, moreover, $D [(-\Delta)^{1/2}] = V$.

Choosing smooth initial data, considering a basis in $D [(-\Delta)^\alpha]$, for $\alpha \geq 1$, and noting that the injection $D [(-\Delta)^{\frac{\alpha}{2}}] \hookrightarrow D [(-\Delta)^{\frac{1}{2}}]$ is continuous we can repeat all the considerations used in the above estimates in order to extend our results for a general class of damped problems

$$\left\{ \begin{array}{l} y_{tt} - M \left(\int_{\Omega} |\nabla y|^2 dx \right) \Delta y + (-\Delta)^\alpha y_t = 0 \quad \text{in } Q \\ y = 0 \quad \text{on } \Sigma_1 \\ M \left(\int_{\Omega} |\nabla y|^2 dx \right) \frac{\partial y}{\partial \nu} + \frac{\partial}{\partial \nu} [(-\Delta)^{\alpha-1} y_t] + |y|^\gamma y = 0 \quad \text{on } \Sigma_0 \\ y(0) = y^0; \quad y_t(0) = y^1 \quad \text{in } \Omega. \end{array} \right.$$

(2) We observe that we could avoid the characterization of the boundary condition if we use Semigroup theory. The linear part of the system generates an analytic semigroup and the nonlinear terms could be treated as perturbations, and it is enough to show solvability for small times. In spite of all these facilities we need the a priori estimates to show global solvability.

Acknowledgement.

The authors are grateful to the referee for the constructive comments and useful suggestions.

References

- [1] A. Arosio - S. Spagnolo, *Global solutions of the Cauchy problem for a nonlinear hyperbolic equation*, Nonlinear Differential Equations and Their Applications (edited by H. Brezis and J. L. Lions), Collège de France Seminar, **6**, Pitman, London (1984).
- [2] M. M. Cavalcanti, V. N. Domingos Cavalcanti & J. A. Soriano, *Existence and Boundary Stabilization of a Nonlinear Hyperbolic Equation with Time-Dependent Coefficients*, EJDE, **1998**(8), (1998), 1-21.

- [3] M. M. Cavalcanti, V. N. Domingos Cavalcanti, J. S. Prates Filho & J. A. Soriano, *Existence and Exponential Decay for a Kirchhoff-Carrier Model with Viscosity*, J. Math. Anal. Appl., **226**(1998), 20-40.
- [4] G. Chen, *A Note on the Boundary Stabilization of the Wave Equation*, Siam J. Control and Optimization, **19**(1), (1981), 106-113.
- [5] Y. Ebihara, M. Milla Miranda & L. A. Medeiros, *Local solutions for a nonlinear degenerated hyperbolic equation*, Nonlinear Anal. Theory, Meth. and Appl., **10**(1), (1986), 27-40.
- [6] A. Friedman, *Variational Principles and Free-Boundary Problems*, John Wiley & Sons, (1982).
- [7] V. Komornik & E. Zuazua, *A Direct Method for Boundary Stabilization of the Wave Equation*, J. Math. Pures et Appl. **69**, (1990), 33-54.
- [8] J. E. Lagnese, *Decay of Solution of the Wave Equations in a Bounded Region with Boundary Dissipation*, Journal of Differential Equation, **50**, (1983), 163-182.
- [9] J. E. Lagnese, *Note on Boundary Stabilization of the Wave Equations*, Siam J. Control and Optimization, **26**, N.5, (1988), 1250-1257.
- [10] I. Lasiecka & D. Tataru, *Uniform Boundary Stabilization of Semilinear Wave Equations with Nonlinear Boundary Damping*, Diff. and Integral Equations, **6**(2), (1993), 507-533.
- [11] I. Lasiecka & J. Ong, *Global Solvability and Uniform Decays of Solutions to Quasilinear Equation with Nonlinear Boundary Conditions*, Communications in PDE's, **24**, (1999), 2069-2109.
- [12] J. L. Lions, *On some questions in boundary value problems of Mathematical Physics, in Contemporary Developments in Continuum Mechanics and Partial Differential Equations*(edited by G. M. de La Penha and L. A. Medeiros), North-Holland, Amsterdam,(1978).
- [13] J. L. Lions - *Quelques Méthodes de Résolution des Problemes aux Limites Non-Lineares*, Dunod, paris, 1969.
- [14] L. A. Medeiros & M. Milla Miranda, *On a nonlinear wave equation with damping*, Revista Matemática de la Universidad Complutense de Madrid, **3** (2), (1990), 613-627.
- [15] M. Milla Miranda, *Traço para o dual dos Espacos de Sobolev*, Bol. Soc. Paran. Mat. (2^a série), **11**(2) , (1990), 131-157.
- [16] M. Milla Miranda & P. San Gil Jutuca, *Existence and Boundary Stabilization of Solutions for the Kirchhoff Equation*, Communications in PDE's, **24**, (1999), 1759-1801.

- [17] G. P. Menzala, *On classical solutions of a quasi-linear hyperbolic equation*, Nonlinear Anal. T. M. A. **3**, (1979), 613-627.
- [18] J. E. Muñoz Rivera, *Smooth effect and decay on a class of non linear evolution equation*, Ann. de la Fac. des Sci. de Toulouse, **1**(2), (1992), 237-260.
- [19] K. Nishihara & Y. Yamada, *On global solutions of some degenerate quasi-linear hyperbolic equations with dissipative terms*, Funkcial. Ekvac. **33**, (1990), 151-159.
- [20] J. Quinn & D. L. Russel, *Asymptotic Stability and Energy Decay for Solutions of Hyperbolic Equations, with Boundary Damping*, Proceedings of the Royal Society of Edinburg, **77A**, (1977), 97-127.
- [21] P. H. Rivera, *On local strong solutions of a nonlinear partial differential equation*, Applicable Analysis, **10**, (1980), 93-104.
- [22] M. Tuskack, *Boundary stabilization for the stretched string equation*, Diff. and Int. Equations, **6**(4), (1993), 925-935.
- [23] C. F. Vasconcellos & L. M. Teixeira, *Strong solutions and exponential decay for a nonlinear hyperbolic equation*, Applicable Analysis, **51**, (1993), 155-173.
- [24] Y. Yamada, *On the decay of solutions for some nonlinear evolution equations of second order*, Nagoya Math. Journal **73**, (1970), 69-98.
- [25] Y. Yamada, *Some nonlinear degenerate wave equations*, NonLinear Analysis, The. Meth. and App., **1** (10), (1987), 1155-1168.
- [26] E. Zuazua, *Stability and decay for a class of nonlinear hyperbolic problems*, Asymptotic Analysis, **1**(1988), 161-185.

Departamento de Matemática
Universidade Estadual de Maringá
87020-900 Maringá-PR, Brasil

Recibido: 26 de Octubre de 1999
Revisado: 3 de Agosto de 2000