

# DELTA LINK-HOMOTOPY ON SPATIAL GRAPHS

Ryo NIKKUNI


## Abstract

We study new equivalence relations in spatial graph theory. We consider natural generalizations of delta link-homotopy on links, which is an equivalence relation generated by delta moves on the same component and ambient isotopies. They are stronger than edge-homotopy and vertex-homotopy on spatial graphs which are natural generalizations of link-homotopy on links. Relationship to existing familiar equivalence relations on spatial graphs are stated, and several invariants are defined by using the second coefficient of the Conway polynomial and the third derivative at 1 of the Jones polynomial of a knot.

## 1 Introduction and results

Throughout this paper we work in the piecewise linear category. Let  $G$  be a finite simple graph, namely it has no loops and multiedges. We denote the set of all edges of  $G$  by  $E(G)$ . We consider  $G$  as a topological space in the usual way and study an embedding of  $G$  into the 3-sphere  $S^3$ , called a *spatial embedding* of  $G$  or simply a *spatial graph*.

In [28], eight equivalence relations (1) *ambient isotopy*, (2) *cobordism*, (3) *isotopy*, (4) *I-equivalence*, (5) *edge-homotopy*, (6) *vertex-homotopy*, (7) *homology*, (8)  $\mathbf{Z}_2$ -*homology* on spatial graphs are introduced and the following implication between them are stated [28, FUNDAMENTAL THEOREM]:



Moreover these eight equivalence relations are different equivalence relations. We refer the reader to [28] for precise definitions. Specially, (5) and (6) (In [28], (5) and (6) was called *homotopy* and *weak homotopy*, respectively) were introduced as natural generalizations of *link-homotopy* [12] on links. Two spatial embeddings  $f, g : G \rightarrow S^3$  are said to be *edge-homotopic* if  $f$  and  $g$  can be transformed into one another by *self-crossing changes* and ambient isotopies, where a self-crossing change is a crossing change on an edge, and *vertex-homotopic* if  $f$  and  $g$  can be transformed into one another by crossing changes on adjacent edges and ambient isotopies.

In this paper we define new equivalence relations and discuss them. A *delta move* is a local move as illustrated in Fig. 1.1. It is known that this move is an *unknotting operation*, namely any knot can be transformed into a trivial one by delta moves and ambient isotopies [11] [16].





Fig. 1.1.

We say that two spatial embeddings  $f, g : G \rightarrow S^3$  are (DEH) *delta edge-homotopic* if  $f$  and  $g$  can be transformed into one another by *self-delta moves* and ambient isotopies, where a self-delta move is a delta move on an edge, and (DVH) *delta vertex-homotopic* if  $f$  and  $g$  can be transformed into one another by *quasi-adjacent delta moves* and ambient isotopies, where a quasi-adjacent delta move is a delta move on exactly two adjacent edges. We note that these equivalence relations coincides

with *delta link-homotopy* (or *self delta-equivalence*) on links [25] if  $G$  is homeomorphic to a mutually disjoint union of 1-spheres. Namely these are natural generalizations of delta link-homotopy on links. We refer the reader to [26], [27], [20], [17], [18] and [19] for works related to delta link-homotopy on links.

It is natural to ask how strong are the above-mentioned equivalence relations. We show the following relations between (1),(2),..., (8) and (DEH), (DVH).

**Theorem 1.1.**



Moreover these ten equivalence relations are different equivalence relations.

A graph  $H$  is called a *minor* of  $G$  if  $H$  is obtained from  $G$  by a finite sequence of the following two operations: (1) edge contraction, (2) taking a subgraph. We note that any subgraph of  $G$  is a minor of  $G$ . For delta vertex-homotopy, we have the following as a corollary of Theorem 1.1 and [28, Theorem B].

**Corollary 1.2.** For a graph  $G$ , the following conditions are mutually equivalent:

- (i) Any two spatial embeddings  $f, g : G \rightarrow S^3$  are delta vertex-homotopic.
- (ii) None of  $G_1, G_2$  and  $G_3$  as illustrated in Fig. 1.2 is a minor of  $G$ . ■

A graph satisfying with the condition of Corollary 1.2 is called a *generalized bouquet* [28]. Therefore Corollary 1.2 means that a quasi-adjacent delta move is an “unknotting operation” for spatial embeddings of a generalized bouquet.

To detect the equivalence class, we construct some delta edge (resp. vertex)-homotopy invariants. A *cycle* is a subgraph of  $G$  which is homeomorphic to the 1-sphere  $S^1$ , and a *k-cycle* is a cycle which contains exactly  $k$  vertices. We regard  $f(\gamma)$  as a knot for a cycle  $\gamma$  and a spatial




Fig. 1.2.

embedding  $f : G \rightarrow S^3$ . We denote the set of all cycles of  $G$ , the set of all cycles containing an edge  $e$  of  $G$  and the set of all cycles containing edges  $e_1, e_2$  of  $G$  by  $\Gamma(G)$ ,  $\Gamma_e(G)$  and  $\Gamma_{e_1, e_2}(G)$ , respectively. Let  $\mathbf{Z}_r = \{0, 1, \dots, r - 1\}$  for a positive integer  $r$  and  $\mathbf{Z}_0 = \mathbf{Z}$ , where  $\mathbf{Z}$  is the integers (We regard  $\mathbf{Z}_r$  ( $r \geq 1$ ) as the cyclic group of order  $r$  whenever we consider  $\mathbf{Z}_r$  a group). Then a map  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  is called a *weight* on  $\Gamma(G)$ . For a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  and a spatial embedding  $f : G \rightarrow S^3$ , we set

$$\tilde{\alpha}_\omega(f) \equiv \sum_{\gamma \in \Gamma(G)} \omega(\gamma) a_2(f(\gamma)) \pmod{r},$$

where  $a_2(J)$  denotes the second coefficient of the *Conway polynomial* [2] of a knot  $J$ . Then we have the following.

- Theorem 1.3.** (1) *If a weight  $\omega$  is weakly balanced on each of edges of  $G$ , then  $\tilde{\alpha}_\omega$  is a delta edge-homotopy invariant.*  
 (2) *If a weight  $\omega$  is weakly balanced on each pair of adjacent edges of  $G$ , then  $\tilde{\alpha}_\omega$  is a delta vertex-homotopy invariant.*

We give the definition of “weakly balanced” in section 3. Let  $d_s$  be the greatest common divisor of  $\{\#\Gamma_e(G) \mid e \in E(G)\}$ , where  $\#\{\cdot\}$  means the number of elements of the set. If  $d_s \geq 2$ , we define a weight  $\omega_s : \Gamma(G) \rightarrow \mathbf{Z}_{d_s}$  by  $\omega_s(\gamma) = 1$  for any  $\gamma \in \Gamma(G)$ . Let  $d_{ad}$  be the greatest common divisor of  $\{\#\Gamma_{e_1, e_2}(G) \mid \text{adjacent edges } e_1, e_2 \in E(G)\}$ . If  $d_{ad} \geq 2$ , we define a weight  $\omega_{ad} : \Gamma(G) \rightarrow \mathbf{Z}_{d_{ad}}$  by  $\omega_{ad}(\gamma) = 1$  for any  $\gamma \in \Gamma(G)$ . Then by Theorem 1.3 we have the following.

- Corollary 1.4.** (1)  *$\tilde{\alpha}_{\omega_s}$  is a delta edge-homotopy invariant.*  
 (2)  *$\tilde{\alpha}_{\omega_{ad}}$  is a delta vertex-homotopy invariant.*

By using  $\tilde{\alpha}_{\omega_s}$ , we show that there exists a non-trivial  $\theta$ -curve up to delta edge-homotopy (Example 4.1).

For a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  and a spatial embedding  $f : G \rightarrow S^3$ , we set

$$n_\omega(f) \equiv \frac{1}{18} \sum_{\gamma \in \Gamma(G)} \omega(\gamma) V_{f(\gamma)}^{(3)}(1) \pmod{r},$$

where  $V_J^{(k)}(1)$  denotes the  $k$ -th derivative at 1 of the Jones polynomial  $V_J(t)$  [6] of a knot  $J$ . We note that  $(1/18) \sum_{\gamma \in \Gamma(G)} \omega(\gamma) V_{f(\gamma)}^{(3)}(1)$  is integer-valued (cf. Remark 3.4 (1)). Then we have the following.

**Theorem 1.5.** (1) *If a weight  $\omega$  is balanced [29] on each of edges of  $G$ , then  $n_\omega$  is a delta edge-homotopy invariant.*

(2) *If a weight  $\omega$  is balanced on each pair of adjacent edges of  $G$ , then  $n_\omega$  is a delta vertex-homotopy invariant.* We give the definition of

“balanced” in section 3. By Theorem 1.1, we have the following immediately.

**Corollary 1.6.** (1) *If a weight  $\omega$  is weakly balanced on each pair of adjacent edges of  $G$ , then  $\tilde{\alpha}_\omega$  is an isotopy invariant.*

(2) *If a weight  $\omega$  is balanced on each pair of adjacent edges of  $G$ , then  $n_\omega$  is an isotopy invariant.* ■

**Remark 1.7.** If a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  is balanced on each of edges (resp. each pair of adjacent edges) of  $G$ , then our  $\tilde{\alpha}_\omega$  coincides with the Taniyama’s edge (resp. vertex)-homotopy invariants  $\alpha_\omega$  [29].

In the next section we prove Theorem 1.1. The proofs of Theorems 1.3 and 1.5 are given in section 3. We give specific examples in section 4. In this paper we calculate the Jones polynomial of a knot by the skein relation  $tV_{J_+}(t) - t^{-1}V_{J_-}(t) = (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{J_0}(t)$ .

## 2 Proof of Theorem 1.1

**Lemma 2.1.** (1)  $\Rightarrow$  (DEH)  $\Rightarrow$  (DVH)  $\Rightarrow$  (5). *Moreover these four equivalence relations are different equivalence relations.*

**Proof.** By the definition,  $(1) \Rightarrow (\text{DEH})$  is clear. Since a self-delta move is realized by quasi-adjacent delta moves (see Fig. 2.1) and a quasi-adjacent delta move is realized by two self-crossing changes (see Fig. 2.2), we have that  $(\text{DEH}) \Rightarrow (\text{DVH}) \Rightarrow (5)$ .




Fig. 2.1.

(1) and (DEH) are different because a trefoil knot is delta-edge homotopic to a trivial knot but they are not ambient isotopic. In section 4, we show that there are delta vertex-homotopic spatial graphs which are not delta edge-homotopic (Example 4.1 and Example 4.2), and edge-homotopic spatial graphs which are not delta vertex-homotopic (Example 4.3). Therefore (1), (DEH), (DVH) and (5) are different equivalence relations. ■

In the following we investigate more about isotopy and delta vertex-homotopy. For a spatial embedding  $f : G \rightarrow S^3$  and a 3-ball  $B$  in  $S^3$ , we say that the pair  $(B, B \cap f(G))$  is a *ball-star pair* (cf. [5]) if  $B \cap f(G)$  is either a proper arc or a star of degree  $n$ , where  $n$  is a natural number, that is,  $\text{int}B$  contains only one vertex  $f(v)$ , and  $B \cap f(G)$  consists of  $n$  edges  $f(e_i)$  that are incident to  $f(v)$  and  $f(\partial e_i) - \{f(v)\} \subset \partial B$  (when  $B \cap f(G)$  is a proper arc, it is regarded as a star of degree 2 even if




Fig. 2.2.

it does not contain vertices of  $f(G)$ ). A ball-star pair  $(B, B \cap f(G))$  is said to be *standard* if there exists a properly embedded 2-disk  $D$  in  $B$  with  $D \supset B \cap f(G)$ . We set  $J = G - f^{-1}(\text{int}B)$ . Then a spatial embedding  $g : G \rightarrow S^3$  is said to be obtained from  $f$  by a *blowing-down* in  $B$  if  $g|_J = f|_J$  and  $(B, B \cap g(G))$  is standard. Conversely  $f$  is said to be obtained from  $g$  by a *blowing-up* occurring in  $B$ . It is known that spatial embeddings  $f, g : G \rightarrow S^3$  are isotopic if and only if they can be transformed into one another by blowing-downs, blowing-ups and ambient isotopies [28] (see also [24] for links). The following lemma shows that (3) implies (5).

**Lemma 2.2.** ([28, Lemma 2.1]) *A blowing down is realized by self-crossing changes.* ■

Moreover we have the following.

**Lemma 2.3.** *Isotopy implies delta vertex-homotopy.*

**Proof.** It is sufficient to show that a blowing-down is realized by quasi-adjacent delta moves. Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a blowing-down in  $B$ . By Lemma 2.2,  $f$  is obtained from  $g$  by self-crossing changes on  $B \cap g(G)$ , where  $(B, B \cap g(G))$  is a standard ball-star pair. For each of crossing points, we deform  $f$  up to ambient isotopy as illustrated in Fig. 2.3. Therefore we can deform  $f$  up to ambient isotopy so that a “band sum of Hopf links and  $g$ ” (cf. [31] and [30]). We note that the deformations as illustrated in Fig. 2.4




Fig. 2.3.

are realized by delta moves [30]. For (1) and (2), see Fig. 2.5 (1) and (2), respectively. (3) is realized by ambient isotopies only. (4) is clear by (2). For (5), see Fig. 2.5 (5). By using the deformation of Fig. 2.4 (1), we can undo the linking between Hopf bands whose attaching edges are different (see Fig. 2.6). Moreover Hopf bands attached to the same edge are gone by self-delta moves by the deformations of Fig. 2.4 (see Fig. 2.7), thus by quasi-adjacent delta moves. Therefore we can remove all Hopf bands by quasi-adjacent delta moves, namely we can obtain  $g$  from  $f$  up to delta vertex-homotopy. This completes the proof. ■

**Lemma 2.4.** (i) *Delta edge-homotopy and delta vertex-homotopy do not imply cobordism, isotopy and I-equivalence.*

(ii) *Cobordism and I-equivalence do not imply delta edge-homotopy and delta vertex-homotopy.*

(iii) *Isotopy does not imply delta edge-homotopy.*

**Proof.** (i) Let  $M$  be a 2-component link as illustrated in Fig. 2.8. We have that  $M$  is delta edge-homotopic to a trivial 2-component link (by using the deformation of Fig. 2.4 (1)) but not  $I$ -equivalent to a trivial 2-component link because they have different Milnor  $\bar{\mu}$ -invariant [13] (see also [1]). Thus we also have that they are not cobordant and isotopic.






Fig. 2.4.




Fig. 2.5.




Fig. 2.6.




Fig. 2.7.

Therefore delta edge-homotopy and delta vertex-homotopy do not imply cobordism, isotopy and  $I$ -equivalence.

(ii) Let  $L$  be a 2-component link as illustrated in Fig. 2.8. It is known that  $L$  and a Hopf link are cobordant but not delta vertex-homotopic [20, Claim 4.5]. Thus a cobordism does not imply delta edge-homotopy and delta vertex-homotopy for links. Thus we also have that  $I$ -equivalence does not imply delta edge-homotopy and delta vertex-homotopy.




Fig. 2.8.

(iii) In Example 4.1, we show that there exist  $\theta$ -curves which are not delta edge-homotopic. Since any  $\theta$ -curves are isotopic (see [28, Theorem B]), we have the desired conclusion. ■

**Proof of Theorem 1.1.** By Lemmas 2.1, 2.3 and 2.4, we have the result. ■

We have the following immediately as well.

**Corollary 2.5.** *Isotopy implies delta link-homotopy on links.* ■

### 3 Delta edge and vertex-homotopy invariants

In this section we prove Theorems 1.3, 1.5 and their corollaries. Let  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  be a weight. Let  $e$  be an edge of  $G$ . Then we say that  $\omega$  is *weakly balanced on  $e$*  if

$$\sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \equiv 0 \pmod{r}.$$

Let  $e_1$  and  $e_2$  be adjacent edges of  $G$ . Then we say that  $\omega$  is *weakly balanced on  $e_1, e_2$*  if

$$\sum_{\gamma \in \Gamma_{e_1, e_2}(G)} \omega(\gamma) \equiv 0 \pmod{r}.$$

**Lemma 3.1.** (1) *Let  $e$  be an edge of  $G$  and  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  a weight which is weakly balanced on  $e$ . Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a self-delta move on  $f(e)$ . Then  $\tilde{\alpha}_\omega(f) \equiv \tilde{\alpha}_\omega(g) \pmod{r}$ .*

(2) *Let  $e_1$  and  $e_2$  be adjacent edges of  $G$  and  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  a weight which is weakly balanced on  $e_1, e_2$ . Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a quasi-adjacent delta move on  $f(e_1)$  and  $f(e_2)$ . Then  $\tilde{\alpha}_\omega(f) \equiv \tilde{\alpha}_\omega(g) \pmod{r}$ .*

To prove Lemma 3.1, we recall the M. Okada's work. She showed that the variation of  $a_2$  of knots which differed by a single delta move is  $\pm 1$  [22, Theorem 1.1]. We can rewrite the fact above as the following:

$$a_2(K_+) - a_2(K_-) = 1, \tag{3.1}$$

where  $K_+, K_-$  are knots as illustrated in Fig. 3.1. We remark here




Fig. 3.1.

that each of the following two local moves (i) and (ii) got from the one as illustrated in Fig. 3.1: (i) the orientation of one string is changed, (ii) the upper-lower relations at all crossings are reversed, is obtained

by using the move as illustrated in Fig. 3.1 (cf. [16]). Therefore any application of a delta move can be regarded just as the one in Fig. 3.1.

**Proof of Lemma 3.1.** (1) Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a single self-delta move on  $f(e)$ . It is sufficient to show that  $\tilde{\alpha}_\omega(f) \equiv \tilde{\alpha}_\omega(g) \pmod r$ . For  $\gamma \in \Gamma_e(G)$ , we may assume that  $f(\gamma)$  and  $g(\gamma)$  are as illustrated in Fig. 3.2 without loss of generality. Then by (3.1) we have that




Fig. 3.2.

$$\begin{aligned}
 \tilde{\alpha}_\omega(f) - \tilde{\alpha}_\omega(g) &\equiv \sum_{\gamma \in \Gamma(G)} \omega(\gamma) a_2(f(\gamma)) - \sum_{\gamma \in \Gamma(G)} \omega(\gamma) a_2(g(\gamma)) \\
 &= \sum_{\gamma \in \Gamma(G)} \omega(\gamma) \{a_2(f(\gamma)) - a_2(g(\gamma))\} \\
 &= \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \{a_2(f(\gamma)) - a_2(g(\gamma))\} \\
 &= \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \\
 &\equiv 0 \pmod r.
 \end{aligned}$$

Therefore we have that  $\tilde{\alpha}_\omega(f) \equiv \tilde{\alpha}_\omega(g) \pmod r$ .

(2) We can prove in the same way as (1) (see Fig. 3.3). ■

**Proof of Theorem 1.3.** It is clear by Lemma 3.1. ■




Fig. 3.3.

**Proof of Corollary 1.4.** It is clear that the weight  $\omega_s : \Gamma(G) \rightarrow \mathbf{Z}_{d_s}$  is weakly balanced on each of edges of  $G$  and the weight  $\omega_{ad} : \Gamma(G) \rightarrow \mathbf{Z}_{d_{ad}}$  is weakly balanced on each pair of adjacent edges of  $G$ . Therefore by Theorem 1.3 we have the desired conclusion. ■

In [29], Taniyama defined edge (resp. vertex)-homotopy invariants  $\alpha_\omega$  of spatial graphs by taking advantage of the following fact which is well-known in knot theory (cf. [8]):

$$a_2(J_+) - a_2(J_-) = lk(J_0), \tag{3.2}$$

where  $J_+, J_-$  and  $J_0$  are knots and a 2-component link which are identical except inside the depicted regions as illustrated in Fig. 3.4, and  $lk$  denotes the *linking number*. We note that  $\alpha_\omega$  is a delta edge (vertex)-




Fig. 3.4.

homotopy invariant by Theorem 1.1. The essential advantage of (3.2)

was that the variation of  $a_2$  of knots which differed by a single crossing change is determined by the linking number which is a homological invariant (cf. [23]). On the other hand, we have the following.

**Theorem 3.2.**

$$\frac{1}{18}V_{K_+}^{(3)}(1) - \frac{1}{18}V_{K_-}^{(3)}(1) = 2Lk(K_0) - 1,$$

where  $K_+$ ,  $K_-$  and  $K_0$  are knots and a 3-component link which are identical except inside the depicted regions as illustrated in Fig. 3.5 and  $Lk$  denotes the total linking number (namely the summation of all linking numbers of 2-component sublinks). ■




Fig. 3.5.

Therefore the variation of  $V^{(3)}(1)$  of knots which differed by a single delta move is also determined by the linking number. We define some delta edge (resp. vertex)-homotopy invariants by taking advantage of Theorem 3.2 in a similar way as Taniyama’s construction of edge (resp. vertex)-homotopy invariants.

To prove Theorem 3.2, we show the following relation between a single delta move on a knot and the Jones polynomial.

**Lemma 3.3.**

$$V_{K_+}(t) - V_{K_-}(t) = (t - 2t^2 + t^3) \{V_{K_\infty}(t) - V_{K_0}(t)\}, \quad (3.3)$$

where  $K_+$ ,  $K_-$ ,  $K_\infty$  and  $K_0$  are knots and a 3-component link which are identical except inside the depicted regions as illustrated in Fig. 3.5 and Fig. 3.6.






Fig. 3.6.




Fig. 3.7.

**Proof.** Consider a skein tree as illustrated in Fig. 3.7. For  $K_2$ ,  $K_4$  and  $K_\infty$  we have that

$$tV_{K_4}(t) - t^{-1}V_{K_2}(t) = (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_\infty}(t). \tag{3.4}$$

For  $K_+$ ,  $K_1$  and  $K_2$  we have that

$$tV_{K_1}(t) - t^{-1}V_{K_+}(t) = (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_2}(t). \tag{3.5}$$

By (3.4) and (3.5) we have that

$$\begin{aligned} & V_{K_1}(t) - t^{-2}V_{K_+}(t) \\ &= t(t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_4}(t) - (t^{-\frac{1}{2}} - t^{\frac{1}{2}})^2V_{K_\infty}(t). \end{aligned} \tag{3.6}$$

On the other hand, for  $K_3$ ,  $K_4$  and  $K_0$  we have that

$$tV_{K_4}(t) - t^{-1}V_{K_3}(t) = (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_0}(t). \tag{3.7}$$

For  $K_1$ ,  $K_-$  and  $K_3$  we have that

$$tV_{K_1}(t) - t^{-1}V_{K_-}(t) = (t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_3}(t). \tag{3.8}$$

By (3.7) and (3.8) we have that

$$\begin{aligned} & V_{K_1}(t) - t^{-2}V_{K_-}(t) \\ &= t(t^{-\frac{1}{2}} - t^{\frac{1}{2}})V_{K_4}(t) - (t^{-\frac{1}{2}} - t^{\frac{1}{2}})^2V_{K_0}(t). \end{aligned} \tag{3.9}$$

By (3.6) and (3.9), we have the desired conclusion. ■

**Proof of Theorem 3.2.** We set  $f(t) = t - 2t^2 + t^3$  and denote the  $k$ -th derivative of  $f(t)$  by  $f^{(k)}(t)$ . By differentiating both sides of (3.3), we have that

$$\begin{aligned} V_{K_+}^{(3)}(t) - V_{K_-}^{(3)}(t) &= f^{(3)}(t) \{V_{K_\infty}(t) - V_{K_0}(t)\} \\ &\quad + 3f^{(2)}(t) \{V_{K_\infty}^{(1)}(t) - V_{K_0}^{(1)}(t)\} \\ &\quad + 3f^{(1)}(t) \{V_{K_\infty}^{(2)}(t) - V_{K_0}^{(2)}(t)\} \\ &\quad + f(t) \{V_{K_\infty}^{(3)}(t) - V_{K_0}^{(3)}(t)\}. \end{aligned}$$

Thus it is clear that

$$\begin{aligned}
 V_{K_+}^{(3)}(1) - V_{K_-}^{(3)}(1) &= 6 \{V_{K_\infty}(1) - V_{K_0}(1)\} \\
 &\quad + 6 \{V_{K_\infty}^{(1)}(1) - V_{K_0}^{(1)}(1)\}. \tag{3.10}
 \end{aligned}$$

Since it is known that

$$V_L(1) = (-2)^{n-1} \quad ([6, \text{Theorem 15}]) \tag{3.11}$$

and

$$\begin{aligned}
 V_L^{(1)}(1) &= \begin{cases} 0 & \text{if } n = 1, \\ 3(-2)^{n-2}Lk(L) & \text{if } n \geq 2, \end{cases} \tag{3.12} \\
 & \quad ([6, \text{Theorem 16}], [15, \text{Theorem 1}])
 \end{aligned}$$

for an  $n$ -component link  $L$ , therefore by (3.11), (3.12) and (3.10) we have that

$$\begin{aligned}
 V_{K_+}^{(3)}(1) - V_{K_-}^{(3)}(1) &= 6 \{1 - (-2)^2\} + 6 \{-3(-2)Lk(K_0)\} \\
 &= 36Lk(K_0) - 18.
 \end{aligned}$$

This completes the proof. ■

**Remark 3.4.** (1) Since the delta move is an unknotting operation for knots, Theorem 3.2 implies that  $(1/18)V_J^{(3)}(1)$  is an integer for any knot  $J$ .

(2) The essential idea of the proof of Lemma 3.3 has already appeared in [22] (namely in the proof of (3.1)) and extended to a self-delta move on links in terms of the Conway polynomial [17]. Moreover it can be applied to the *HOMFLY polynomial* [4]. We are due to mention it in the future paper [7].

Let  $e$  be an edge of  $G$ . We give an orientation to  $e$ . We give an orientation to each  $\gamma \in \Gamma_e(G)$  by the orientation of  $e$ . Since  $H_1(G; \mathbf{Z}_r) = \text{Ker}\partial_1$ , we denote  $[\gamma] \in H_1(G; \mathbf{Z}_r)$  by  $\gamma$  for a cycle  $\gamma$ . Then a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  is said to be *balanced on  $e$*  if

$$\sum_{\gamma \in \Gamma_e(G)} \omega(\gamma)\gamma = 0 \in H_1(G; \mathbf{Z}_r).$$

Note that this definition does not depend on the choice of the orientation of  $e$ .

Let  $e_1, e_2$  be adjacent edges of  $G$ . We give an orientation to  $e_1$ . We give an orientation to each  $\gamma \in \Gamma_{e_1, e_2}(G)$  by the orientation of  $e_1$ . Then a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  is said to be *balanced on  $e_1, e_2$*  if

$$\sum_{\gamma \in \Gamma_{e_1, e_2}(G)} \omega(\gamma)\gamma = 0 \in H_1(G; \mathbf{Z}_r).$$

We also note that this definition does not depend on the choice of the orientation of  $e_1$ . We remark here that a weight  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  which is balanced on an edge  $e$  (resp. adjacent edges  $e_1, e_2$ ) is weakly balanced on an edge  $e$  (resp. adjacent edges  $e_1, e_2$ ), namely it is easy to see that if  $\sum_{\gamma \in \Gamma_e(G)} \omega(\gamma)\gamma = 0$  (resp.  $\sum_{\gamma \in \Gamma_{e_1, e_2}(G)} \omega(\gamma)\gamma = 0$ ) in  $H_1(G; \mathbf{Z}_r)$  then  $\sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \equiv 0 \pmod{r}$  (resp.  $\sum_{\gamma \in \Gamma_{e_1, e_2}(G)} \omega(\gamma) \equiv 0 \pmod{r}$ ).

**Lemma 3.5.** (1) *Let  $e$  be an edge of  $G$  and  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  a weight which is balanced on  $e$ . Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a self-delta move on  $f(e)$ . Then  $n_\omega(f) \equiv n_\omega(g) \pmod{r}$ .*

(2) *Let  $e_1$  and  $e_2$  be adjacent edges of  $G$  and  $\omega : \Gamma(G) \rightarrow \mathbf{Z}_r$  a weight which is balanced on  $e_1, e_2$ . Let  $f, g : G \rightarrow S^3$  be spatial embeddings such that  $g$  is obtained from  $f$  by a quasi-adjacent delta move on  $f(e_1)$  and  $f(e_2)$ . Then  $n_\omega(f) \equiv n_\omega(g) \pmod{r}$ .*

**Proof.** (1) Note that the orientation of  $\gamma \in \Gamma_e(G)$  is given by the one of  $e$ . We may assume that  $f(\gamma)$  and  $g(\gamma)$  are as illustrated in Fig. 3.8 without loss of generality. Let  $L_{f,g}(\gamma) = l_{f,g}^{(1)}(\gamma) \cup l_{f,g}^{(2)} \cup l_{f,g}^{(3)}$  be a 3-component link as the right-hand figure in Fig. 3.8. Note that  $l_{f,g}^{(2)}$  and  $l_{f,g}^{(3)}$  are common to all  $\gamma \in \Gamma_e(G)$ . Then we have

$$\begin{aligned} & n_\omega(f) - n_\omega(g) \\ & \equiv \frac{1}{18} \sum_{\gamma \in \Gamma(G)} \omega(\gamma)V_{f(\gamma)}^{(3)}(1) - \frac{1}{18} \sum_{\gamma \in \Gamma(G)} \omega(\gamma)V_{g(\gamma)}^{(3)}(1) \\ & = \sum_{\gamma \in \Gamma(G)} \omega(\gamma) \left\{ \frac{1}{18}V_{f(\gamma)}^{(3)}(1) - \frac{1}{18}V_{g(\gamma)}^{(3)}(1) \right\} \end{aligned}$$




Fig. 3.8.

$$\begin{aligned}
 &= \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \left\{ \frac{1}{18} V_{f(\gamma)}^{(3)}(1) - \frac{1}{18} V_{g(\gamma)}^{(3)}(1) \right\} \\
 &= \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \left\{ 2lk(l_{f,g}^{(1)}(\gamma), l_{f,g}^{(2)}) + 2lk(l_{f,g}^{(1)}(\gamma), l_{f,g}^{(3)}) + 2lk(l_{f,g}^{(2)}, l_{f,g}^{(3)}) - 1 \right\} \\
 &= 2 \sum_{k=2,3} \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) lk(l_{f,g}^{(1)}(\gamma), l_{f,g}^{(k)}) + \left\{ 2lk(l_{f,g}^{(2)}, l_{f,g}^{(3)}) - 1 \right\} \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) \\
 &\equiv 2 \sum_{k=2,3} lk \left( \sum_{\gamma \in \Gamma_e(G)} \omega(\gamma) l_{f,g}^{(1)}(\gamma), l_{f,g}^{(k)} \right) \\
 &\equiv 2 \sum_{k=2,3} lk(0, l_{f,g}^{(k)}) \\
 &\equiv 0 \pmod{r}.
 \end{aligned}$$

Thus we have the desired conclusion.

(2) We can complete the proof in a similar way as (1) (see Fig. 3.9). ■

**Proof of Theorem 1.5.** It is clear by Lemma 3.5. ■

### 4 Examples

**Example 4.1.** Let  $f_m$  be a  $\theta$ -curve as illustrated in Fig. 4.1, where  $m$  is a non-negative integer. Note that  $f_i$  and  $f_j$  are delta vertex-homotopic for any  $i, j$ . It is easy to see that  $f_m$  contains a unique non-trivial knot  $J$  which is a connected sum of  $m$  trefoil knots for  $m \neq 0$ . Since  $\sharp\Gamma_e(\theta) = 2$




Fig. 3.9.




Fig. 4.1.

for any edge  $e$ , we have that  $d_s = 2$ . By a calculation we have that  $a_2(J) = m$ . Therefore we have that  $\tilde{\alpha}_{\omega_s}(f_m) \equiv 0 \pmod{2}$  if  $m$  is even and  $1 \pmod{2}$  if  $m$  is odd. Thus by Corollary 1.4 (1) we have that  $f_i$  and  $f_j$  are not delta edge-homotopic if the one of  $i, j$  is odd and the other is even. Especially  $f_m$  is not delta edge-homotopic to the trivial embedding  $f_0$  if  $m$  is odd.

**Example 4.2.** (cf. [29, Example 3.1]) Let  $K_4$  be the complete graph on 4 vertices (namely the graph  $G_2$  as illustrated in Fig. 1.2). Let  $f_m$  and  $g$  be spatial embeddings of  $K_4$  as illustrated in Fig. 4.2, where  $m$  is an integer. Note that  $f_m$  and  $g$  are delta vertex-homotopic (Undo the local Borromean ring in  $f(K_4)$  by a quasi-adjacent delta move). It




Fig. 4.2.

is easy to see that  $f_m(K_4)$  contains two non-trivial knots  $J_1$  and  $J_2$  as illustrated in Fig. 4.3. Since  $\sharp\Gamma_e(K_4) = 4$  for any edge  $e \in E(K_4)$ , we have that  $d_s = 4$ . By a calculation we have that  $a_2(J_1) = a_2(J_2) = 1$ . Therefore we have that  $\tilde{\alpha}_{\omega_s}(f_m) \equiv 2 \pmod{4}$  for any integer  $m$ . Since  $\tilde{\alpha}_{\omega_s}(g) \equiv 0 \pmod{4}$ , by Theorem 1.4 (1) we have that  $f_m$  and  $g$  are not delta edge-homotopic for any integer  $m$ .

Let  $\omega : \Gamma(K_4) \rightarrow \mathbf{Z}$  be a weight defined by

$$\omega(\gamma) = \begin{cases} 1 & \text{if } \gamma \text{ is a 4-cycle,} \\ -1 & \text{if } \gamma \text{ is a 3-cycle.} \end{cases}$$

Then we can see that  $\omega$  is balanced on each of edges of  $K_4$ . Thus by Theorem 1.5 (1),  $n_\omega$  is a delta edge-homotopy invariant. By Theorem 3.2 we have that  $V_{J_1}^{(3)}(1) = 36m - 18$  and  $V_{J_2}^{(3)}(1) = -18$ . Therefore

we have that  $n_\omega(f_m) = 2m$ . This shows that  $f_i$  and  $f_j$  are not delta edge-homotopic for  $i \neq j$ .




Fig. 4.3.

**Example 4.3.** (cf. [29, Example 3.2]) Let  $K_5$  be the complete graph on 5 vertices. Let  $f_m$  be a spatial embedding of  $K_5$  as illustrated in Fig. 4.4, where  $m$  is a non-negative integer. Note that  $f_i$  and  $f_j$  are edge-homotopic for any  $i, j$  (Undo the local Borromean ring in  $f_m(K_5)$  by two self-crossing changes). It is easy to see that  $f_m(K_5)$  contains six




Fig. 4.4.

non-trivial knots  $J_k$  ( $k = 1, 2, \dots, 6$ ) for  $m \neq 0$  as illustrated in Fig. 4.5. Note that  $J_k$  is the image of a 4-cycle if  $k = 1, 2$  and the image of a 5-cycle if  $k = 3, 4, 5, 6$ . Since  $\#\Gamma_{e_1, e_2}(K_5) = 5$  for any adjacent edges  $e_1, e_2 \in E(K_5)$ , we have that  $d_{ad} = 5$ . By a calculation we have that  $a_2(J_1) = a_2(J_3) = a_2(J_5) = m$  and  $a_2(J_2) = a_2(J_4) = a_2(J_6) = -m$ . Therefore we have that  $\tilde{\alpha}_{\omega_{ad}}(f_m) \equiv 0 \pmod{5}$ . Since  $\tilde{\alpha}_{\omega_{ad}}(g) \equiv 0$



(mod 5), We can not distinguish  $f_i$  and  $f_j$  for  $i \neq j$  up to delta vertex-homotopy by using  $\tilde{\alpha}_{\omega_{ad}}$ .

Let  $\omega : \Gamma(K_5) \rightarrow \mathbf{Z}$  be a weight defined by

$$\omega(\gamma) = \begin{cases} 1 & \text{if } \gamma \text{ is a 5-cycle,} \\ -1 & \text{if } \gamma \text{ is a 4-cycle and} \\ 0 & \text{if } \gamma \text{ is a 3-cycle.} \end{cases}$$

Then we can see that  $\omega$  is balanced on each pair of adjacent edges of  $K_5$ . Thus by Theorem 1.5 (2),  $n_\omega$  is a delta vertex-homotopy invariant. By Theorem 3.2 we have that  $V_{J_k}^{(3)}(1) = -18m$  ( $k = 1, 2, \dots, 6$ ). Therefore we have that  $n_\omega(f_m) = -2m_k$ . This shows that  $f_i$  and  $f_j$  are not delta vertex-homotopic for  $i \neq j$ .




Fig. 4.5.

**Acknowledgements.** The author is most grateful to Professors T. Asoh and K. Shimokawa for their invaluable comments and encouragement. The author is grateful to the referee for his (or her) comments.

## References

- [1] Cochran, Tim D., *Concordance invariance of coefficients of Conway's link polynomial*, Invent. Math., 82 (1985) 527-541.
- [2] Conway, J.H., *An enumeration of knots and links, and some of their algebraic properties*, In: Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967) (1970) 329-358.
- [3] Conway, J.H. and Gordon, McA., *Knots and links in spatial graphs*, J. Graph Theory, 7 (1983) 445-453.
- [4] Freyd, P., Yetter, D., Hoste, J., Lickorish, W.B.R., Millett, K. and Ocneanu, A., *A new polynomial invariant of knots and links*, Bull. Amer. Math. Soc. (N.S.), 12 (1985) 239-246.
- [5] Inaba, H. and Soma, T., *On spatial graphs isotopic to planar embeddings*, In: S. Suzuki (ed.) Proceedings of Knots '96 Tokyo (World Scientific Publ. Co., 1997) 1-22.
- [6] Jones, V.F.R., *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc., 12 (1985) 103-111.
- [7] Kanenobu, T. and Nikkuni, R., *Delta move and polynomial invariants of links*, preprint.
- [8] Kauffman, L., *Formal Knot Theory* (Mathematical Notes 30, Princeton Univ. Press, 1983).
- [9] Kauffman, L., *On Knots* (Ann. of Math. Studies, 115, Princeton Univ. Press, Princeton, N.J, 1987)
- [10] Kawachi, A., *A Survey of Knot Theory* (Birkhäuser, 1996).
- [11] Matveev, S., *Generalized surgeries of three-dimensional manifolds and representations of homology sphere* (Russian), Mat. Zametki, 42 (1987) 268-278, 345 (English translation: Math. Notes 42 (1987) 651-656.)
- [12] Milnor, J., *Link groups*, Ann. of Math. 59 (1954) 177-195.
- [13] Milnor, J., *Isotopy of links. Algebraic geometry and topology*, In: A symposium in honor of S. Lefschetz, (Princeton University Press, Princeton, N. J., 1957) 280-306.
- [14] Motohashi, T. and Taniyama, K., *Delta unknotting operation and vertex homotopy of spatial graphs*, In: S. Suzuki (ed.) Proceedings of Knots '96 Tokyo (World Scientific Publ. Co., 1997) 185-200.
- [15] Murakami, H., *On derivatives of the Jones polynomial*, Kobe J. Math., 3 (1986) 61-64.

- [16] Murakami, H. and Nakanishi, Y., *On a certain move generating link-homology*, Math. Ann., 284 (1989) 75-89.
- [17] Nakanishi, Y., *Delta link homotopy for two component links*, to appear in Topology and its Applications. (Proceedings of Mexico-Japan first joint meeting for Topology and its Applications.)
- [18] Nakanishi, Y. and Ohyama, Y., *Delta link homotopy for two component links, II*, to appear in Journal of Knot Theory and Its Ramifications
- [19] Nakanishi, Y. and Ohyama, Y., *Delta link homotopy for two component links, III*, preprint
- [20] Nakanishi, Y. and Shibuya, T., *Link homotopy and quasi self delta-equivalence for links*, J. Knot Theory Ramifications., 9 (2000), 683-691.
- [21] Ohyama, Y. and Taniyama, K., *Vassiliev invariants of knots in a spatial graph*, to appear in Pacific Journal of Mathematics
- [22] Okada, M., *Delta-unknotting operation and the second coefficient of the Conway polynomial*, J. Math. Soc. Japan., 42 (1990) 713-717.
- [23] Rolfsen, D., *Knots and Links*, (Mathematics Lecture Series, 7, Publish or Perish, Inc., Berkeley, Calif., 1976).
- [24] Rolfsen, D., *Isotopy of links in codimension two*, J. Indian Math. Soc. (N.S.), 36 (1972) 263-278.
- [25] Shibuya, T., *Self  $\Delta$ -equivalence of ribbon links*, Osaka J. Math., 33 (1996) 751-760.
- [26] Shibuya, T., *Self delta-equivalence of  $F$ -isotopic links*, Mem. Osaka Inst. Tech. Ser. A, 43 (1998) 21-25.
- [27] Shibuya, T., *On self  $\Delta$ -equivalence of boundary links*, Osaka J. Math., 37 (2000) 37-55.
- [28] Taniyama, K., *Cobordism, homotopy and homology of graphs in  $\mathbf{R}^3$* , Topology, 33 (1994) 509-523.
- [29] Taniyama, K., *Link homotopy invariants of graphs in  $\mathbf{R}^3$* , Rev. Mat. Univ. Complut. Madrid., 7 (1994) 129-144.
- [30] Taniyama, K. and Yasuhara, A., *Clasp-pass moves on knots, links and spatial graphs*, to appear in Topology and its Applications
- [31] Yamamoto, M., *Knots in spatial embeddings of the complete graph on four vertices*, Topology Appl., 36 (1990) 291-298.

Division of Mathematics,  
Graduate School of Information Sciences,  
Tohoku University,  
Aramaki aza Aoba 09, Aoba-ku, Sendai 980-8579, Japan  
*E-mail*: nick@ims.is.tohoku.ac.jp

Recibido: 18 de Abril de 2001  
Revisado: 7 de Marzo de 2002