

A capacity approach to the Poincaré inequality and Sobolev imbeddings in variable exponent Sobolev spaces

Petteri HARJULEHTO and Peter HÄSTÖ

Department of Mathematics
P.O. Box 4 (Yliopistonkatu 5)
FIN-00014 University of Helsinki, Finland
petteri.harjulehto@helsinki.fi,
peter.hasto@helsinki.fi

Recibido: 7 de Mayo de 2003
Aceptado: 10 de Septiembre de 2003

ABSTRACT

We study the Poincaré inequality in Sobolev spaces with variable exponent. Under a rather mild and sharp condition on the exponent p we show that the inequality holds. This condition is satisfied e. g. if the exponent p is continuous in the closure of a convex domain. We also give an essentially sharp condition for the exponent p as to when there exists an imbedding from the Sobolev space to the space of bounded functions.

Key words: Sobolev spaces, variable exponent, Poincaré inequality, Sobolev imbedding, continuity

2000 Mathematics Subject Classification: 46E35

1. Introduction

There has recently been a surge of interest in Sobolev spaces with variable exponent, cf. [4–7, 9–11, 17, 22]. These spaces, introduced in [17], are the natural generalization of Sobolev spaces to the non-homogeneous situation; they have been used e. g. in modeling electrorheological fluids, see the book of M. Růžička, [22]. Lebesgue and Sobolev spaces with variable exponent share many properties with their classical equivalents, but there is also some crucial differences. For instance the Hardy-Littlewood maximal

The second author was supported financially by the Academy of Finland.

operator is bounded on $L^{p(\cdot)}$ if the exponent is 0-Hölder continuous (i. e. satisfies (10)) and $1 < \text{ess inf } p \leq \text{ess sup } p < \infty$, [5]. If the exponent is not 0-Hölder continuous, then the maximal operator need not be bounded on $L^{p(\cdot)}$, [21].

The Poincaré inequality, although of great importance in classical non-linear potential theory (especially in metric spaces) has not been previously studied in the case of variable exponent Sobolev spaces. Our first result, Theorem 2.2, is the following: If $D \subset \mathbb{R}^n$ is smooth domain, say a John domain, and the essential supremum of p is less than the Sobolev conjugate of the essential infimum of p then the Poincaré inequality

$$\|u - u_B\|_{L^{p(\cdot)}(D)} \leq C \|\nabla u\|_{L^{p(\cdot)}(D)}$$

holds for every $u \in W^{1,p(\cdot)}(D)$, where $u_B = \int u(x)dx$. Here the constant C depends on $n, p, \text{diam}(D)$ and the John constant of D . We give an example which shows that the condition for p is sharp even in a ball. It follows from this that if p is continuous in the closure of a convex domain then the Poincaré inequality holds (Corollary 2.7).

In classical theory the constant of the Poincaré inequality is $C \text{diam}(D)$. It is possible to achieve this also for variable exponent Sobolev spaces, as we prove in Corollary 2.10. The price we have to pay is that the exponent p has to be 0-Hölder continuous.

Sobolev imbeddings in variable exponent Sobolev spaces have been studied by many authors in the case when p is less than the dimension, see [6, 9–11]. We give two results in the case when p is greater than the dimension. We prove a result for continuity of the Sobolev functions, namely that every Sobolev function is continuous if the exponent is locally bounded away from the dimension. We show that if a domain satisfies a uniform interior cone condition and $p(x) \geq n + f(d(x, \partial G))$ for every x and a certain increasing function f then there exists an imbedding from the variable exponent Sobolev space to L^∞ . Our condition is essentially sharp.

Notation

We denote by \mathbb{R}^n the Euclidean space of dimension $n \geq 2$. For $x \in \mathbb{R}^n$ and $r > 0$ we denote an open ball with center x and radius r by $B(x, r)$.

Let $A \subset \mathbb{R}^n$ and $p: A \rightarrow [1, \infty)$ be a measurable function (called a *variable exponent* on A). We define $p_A^+ = \text{ess sup}_{x \in A} p(x)$ and $p_A^- = \text{ess inf}_{x \in A} p(x)$. If $A = \mathbb{R}^n$ we write $p^+ = p_{\mathbb{R}^n}^+$ and $p^- = p_{\mathbb{R}^n}^-$.

Let $\Omega \subset \mathbb{R}^n$ be an open set. We define the *generalized Lebesgue space* $L^{p(\cdot)}(\Omega)$ to consist of all measurable functions $u: \Omega \rightarrow \mathbb{R}$ such that

$$\varrho_{p(\cdot)}(\lambda u) = \int_{\Omega} |\lambda u(x)|^{p(x)} dx < \infty$$

for some $\lambda > 0$. The function $\varrho_{p(\cdot)}: L^{p(\cdot)}(\Omega) \rightarrow [0, \infty)$ is called the *modular* of the space $L^{p(\cdot)}(\Omega)$. One can define a norm, the so-called *Luxemburg norm*, on this space by the formula $\|u\|_{p(\cdot)} = \inf\{\lambda > 0 : \varrho_{p(\cdot)}(u/\lambda) \leq 1\}$. Notice that if $p \equiv p_0$ then

$L^{p(\cdot)}(\Omega)$ is the classical Lebesgue space, so there is no danger of confusion with the new notation.

The *generalized Sobolev space* $W^{1,p(\cdot)}(\Omega)$ is the space of measurable functions $u: \Omega \rightarrow \mathbb{R}$ such that u and the absolute value of the distributional gradient $\nabla u = (\partial_1 u, \dots, \partial_n u)$ are in $L^{p(\cdot)}(\Omega)$. The function $\varrho_{1,p(\cdot)}: W^{1,p(\cdot)}(\Omega) \rightarrow [0, \infty)$ is defined as $\varrho_{1,p(\cdot)}(u) = \varrho_{p(\cdot)}(u) + \varrho_{p(\cdot)}(|\nabla u|)$. The norm $\|u\|_{1,p(\cdot)} = \|u\|_{p(\cdot)} + \|\nabla u\|_{p(\cdot)}$ makes $W^{1,p(\cdot)}(\mathbb{R}^n)$ a Banach space.

See [17] for basic properties of variable exponent Lebesgue and Sobolev spaces.

2. The Poincaré inequality

In this section we give a relatively mild condition on the exponent for the Poincaré inequality to hold. We also show that this condition is, in a certain sense, the best possible. For Sobolev functions with zero boundary values the Poincaré inequality was given in [10, Lemma 3.1] and considerably generalized in [14].

Recall the following well known Sobolev-Poincaré inequality. By q^* we denote the Sobolev conjugate of $q < n$, $q^* = nq/(n - q)$.

Lemma 2.1. *Let $D \subset \mathbb{R}^n$ be a bounded John domain. Let $1 \leq p < n$ and $p \leq q \leq p^*$ be fixed exponents. Then*

$$\|u - u_D\|_q \leq C(n, p, \lambda) |D|^{1/n+1/q-1/p} \|\nabla u\|_p$$

for all functions $u \in W^{1,p}(D)$, where λ is the John constant.

If $p \geq n$ and $q < \infty$ then

$$\|u - u_D\|_q \leq C(n, q, \lambda) |D|^{1/n+1/q-1/p} \|\nabla u\|_p$$

for all functions $u \in W^{1,p}(D)$.

Proof. The case $p < n$ and $q = p^*$ is by B. Bojarski [3, (6.6)]. The case $q < p^*$ follows from this by standard arguments: we choose $s \in [1, n)$ such that $s^* = q$ (or $s = 1$ if $q < 1^*$). By Hölder's inequality and Bojarski's result we obtain

$$\begin{aligned} \left(\int_D |u - u_D|^q dx \right)^{\frac{1}{q}} &\leq |D|^{-\frac{1}{s^*}} \left(\int_D |u - u_D|^{s^*} dx \right)^{\frac{1}{s^*}} \leq C |D|^{-\frac{1}{s^*}} \left(\int_D |\nabla u|^s dx \right)^{\frac{1}{s}} \\ &= C |D|^{\frac{1}{s} - \frac{1}{s^*}} \left(\int_D |\nabla u|^s dx \right)^{\frac{1}{s}} \leq C |D| \left(\int_D |\nabla u|^p dx \right)^{\frac{1}{p}}, \end{aligned}$$

which is clearly equivalent to the inequalities in the theorem. □

Theorem 2.2. *Let $D \subset \mathbb{R}^n$ be a bounded John domain, with constant λ . If $p_D^+ \leq (p_D^-)^*$ or $p_D^- \geq n$ and $p_D^+ < \infty$ then there exists a constant $C = C(n, p_D^-, p_D^+, \lambda)$ such that*

$$\|u - u_D\|_{p(\cdot)} \leq C(1 + |D|)^2 |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p(\cdot)} \tag{1}$$

for every $u \in W^{1,p(\cdot)}(D)$.

Proof. Assume first that $p_D^+ \leq (p_D^-)^*$. Since $p(x) \leq p_D^+ \leq (p_D^-)^*$ we obtain by [17, Theorem 2.8] and Lemma 2.1 that

$$\begin{aligned} \|u - u_D\|_{p(\cdot)} &\leq (1 + |D|) \|u - u_D\|_{p_D^+} \\ &\leq C(n, p_D^-, \lambda) (1 + |D|) |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p_D^-} \\ &\leq C(n, p_D^-, \lambda) (1 + |D|)^2 |D|^{\frac{1}{n} + \frac{1}{p_D^+} - \frac{1}{p_D^-}} \|\nabla u\|_{p(\cdot)}. \end{aligned}$$

The case $p_D^- \geq n$ is similar, the only difference is that the constant in the second inequality in the above chain of inequalities is $C(n, p_D^+, \lambda)$. □

Remark 2.3. John domains are almost the right class of irregular domains for the classical Sobolev-Poincaré inequality, see [3], [1] and [2, Theorem 4.1].

Previous results on Sobolev imbeddings in the variable exponent setting have been derived in domains whose boundary is locally a graph of a Lipschitz continuous function, see [9–11]. It is therefore of interest to note that every domain, whose boundary is locally the graph of a Lipschitz continuous function, is a John domain, see [19]. In particular every ball is a John domain.

If D is a ball in Theorem 2.2, then the constant in inequality (1) is the classical Sobolev-Poincaré inequality in a ball, see for example [18, Corollary 1.64, p. 38].

The next example shows that if $p_D^- < n$ and $p_D^+ > (p_D^-)^*$ then there need not exist a constant $C > 0$ such that inequality (1) holds for every $u \in W^{1,p(\cdot)}(D)$.

Recall that the *variational capacity* for fixed p , $\text{cap}_p(E, F; D)$, is defined for sets E, F and open D by

$$\text{cap}_p(E, F; D) = \inf_{u \in L(E, F; D)} \int_D |\nabla u|^p dx,$$

where $L(E, F; D)$ is the set of continuous functions u that satisfy $u|_{E \cap D} = 1$, $u|_{F \cap D} = 0$ and $|\nabla u| \in L^{p(\cdot)}(D)$. We use the short-hand notation $\text{cap}(E, F)$ for $\text{cap}(E, F; \mathbb{R}^n)$, similarly for $L(E, F)$. For more information on capacities see [15, Chapter 2] or [20]. The following lemma will be used several times to estimate the gradient of variable exponent functions.

Lemma 2.4 ([15, Example 2.12, p. 35]). *For fixed $p \neq 1, n$, arbitrary $x \in \mathbb{R}^n$ and $R > r > 0$ we have*

$$\text{cap}_p(\mathbb{R}^n \setminus B(x, R), B(x, r)) = \omega_{n-1} \left| \frac{p-n}{p-1} \right|^{p-1} \left| R^{(p-n)/(p-1)} - r^{(p-n)/(p-1)} \right|^{1-p}.$$

Example 2.5. Our aim is construct a sequence of functions in $B = B(0, 1) \subset \mathbb{R}^2$ for which the constant in the Poincaré inequality (1) goes to infinity. Let $B_i = B(2^{-i}e_1, \frac{1}{4}2^{-i}) \subset \mathbb{R}^2$ and $B'_i = B(2^{-i}e_1, \frac{1}{8}2^{-i^2}) \subset \mathbb{R}^2$ for every $i = 1, 2, \dots$ and let $1 < p_1 < 2$. Choose a function $u_i \in C_0^\infty(B_i)$ with $u_i|_{B'_i} = 1$ be such that

$$\left(2 \text{cap}_{p_1}(B'_i, \mathbb{R}^2 \setminus B_i) \right)^{\frac{1}{p_1}} \geq \|\nabla u_i\|_{L^{p_1}(B_i)}. \tag{2}$$

Let $p_2 > 2$ and define $p(x) = p_1 \chi_{B_i \setminus B'_i}(x) + p_2 \chi_{B'_i}(x)$ for $x \in B$ with positive first coordinate. Since $\nabla u_i = 0$ in B'_i we obtain

$$\|\nabla u_i\|_{L^{p(\cdot)}(B_i)} = \|\nabla u_i\|_{L^{p_1}(B_i)}. \tag{3}$$

Let $\tilde{B}_i = B(-2^{-i}e_1, \frac{1}{4}2^{-i})$. We extend u_i to B as an odd function of the first coordinate in \tilde{B}_i and by zero elsewhere. We also extend p to B as an even function of the first coordinate. We denote the extensions by \tilde{u}_i and \tilde{p} . By (2) and (3) we obtain

$$2^{1+\frac{1}{p_1}} \left(\text{cap}_{p_1}(B'_i, \mathbb{R}^2 \setminus B_i) \right)^{\frac{1}{p_1}} \geq \|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)}.$$

By Lemma 2.4 this yields

$$\|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)} \leq C(p_1) \left| \frac{1}{4} 2^{-i \frac{p_1-2}{p_1-1}} - \frac{1}{8} 2^{-i^2 \frac{p_1-2}{p_1-1}} \right|^{\frac{1-p_1}{p_1}}. \tag{4}$$

For large i the right hand side is approximately equal to $C(p_1) 2^{-i^2 \frac{2-p_1}{p_1}}$.

Since $(\tilde{u}_i)_B = 0$, we obtain

$$\|\tilde{u} - (\tilde{u}_i)_B\|_{L^{\tilde{p}(\cdot)}(B)} = \|\tilde{u}\|_{L^{\tilde{p}(\cdot)}(B)} \geq |B'_i|^{\frac{1}{p_2}} \approx 2^{-i^2 \frac{2}{p_2}}. \tag{5}$$

By inequalities (4) and (5) we find that

$$\frac{\|\tilde{u} - (\tilde{u}_i)_B\|_{L^{\tilde{p}(\cdot)}(B)}}{\|\nabla \tilde{u}_i\|_{L^{\tilde{p}(\cdot)}(B)}} \geq C(p_1) 2^{i^2 (\frac{2}{p_1} - 1 - \frac{2}{p_2})} \rightarrow \infty$$

as $i \rightarrow \infty$ if $\frac{2}{p_1} - 1 - \frac{2}{p_2} > 0$, that is, if $p_2 > \frac{2p_1}{2-p_1} = p_1^*$.

We next show that the condition $p_D^+ \leq (p_D^-)^*$ in Theorem 2.2 can be replaced by a set of local conditions.

Theorem 2.6. *Let $D \subset \mathbb{R}^n$ be a bounded John domain. Assume that there exist John domains G_i , $i = 1, \dots, j$, so that $G_i \subset D$ for every i , $D = \cup_{i=1}^j G_i$ and either $p_{G_i}^+ \leq (p_{G_i}^-)^*$ or $p_{G_i}^- \geq n$ for every i . Then there exists a constant $C > 0$ such that*

$$\|u - u_D\|_{p(\cdot)} \leq C \|\nabla u\|_{p(\cdot)} \tag{6}$$

for every $u \in W^{1,p(\cdot)}(D)$. The constant C depends on n , $\text{diam}(D)$, $|G_i|$, p and the John constants of D and G_i , $i = 1, \dots, j$.

Proof. Using the triangle inequality of the norm we obtain

$$\begin{aligned} \|u - u_D\|_{L^{p(\cdot)}(D)} &\leq \sum_{i=1}^j \|u - u_D\|_{L^{p(\cdot)}(G_i)} \\ &\leq \sum_{i=1}^j \|u - u_{G_i}\|_{L^{p(\cdot)}(G_i)} + \sum_{i=1}^j \|u_D - u_{G_i}\|_{L^{p(\cdot)}(G_i)}. \end{aligned} \tag{7}$$

We estimate the first part of the sum using Theorem 2.2. This yields

$$\begin{aligned} \|u - u_{G_i}\|_{L^{p(\cdot)}(G_i)} &\leq C(n, p_{G_i}, |G_i|, \lambda_i) \|\nabla u\|_{L^{p(\cdot)}(G_i)} \\ &\leq C(n, p_{G_i}, |G_i|, \lambda_i) \|\nabla u\|_{L^{p(\cdot)}(D)} \end{aligned} \tag{8}$$

for every $i = 1, \dots, j$. Here λ_i is the John constant of G_i . We next estimate the second part of the sum in (7) using the classical Poincaré inequality for the third inequality. We obtain

$$\begin{aligned} \|u_D - u_{G_i}\|_{L^{p(\cdot)}(G_i)} &\leq \|1\|_{L^{p(\cdot)}(G_i)} \int_{G_i} |u(x) - u_D| dx \\ &\leq \|1\|_{L^{p(\cdot)}(G_i)} |G_i|^{-1} \int_D |u(x) - u_D| dx \\ &\leq C(n, \text{diam}(D), \lambda) |G_i|^{-1} \|1\|_{L^{p(\cdot)}(G_i)} \|\nabla u\|_{L^1(D)} \\ &\leq C(n, \text{diam}(D), \lambda) (1 + |D|) |G_i|^{-1} \|1\|_{L^{p(\cdot)}(G_i)} \|\nabla u\|_{L^{p(\cdot)}(D)} \end{aligned} \tag{9}$$

for every $i = 1, \dots, j$. Here λ is the John constant of D . Now inequality (6) follows by inequalities (7), (8) and (9). □

Corollary 2.7. *Let $D \subset \mathbb{R}^n$ be a bounded convex domain and let $p: \bar{D} \rightarrow [1, \infty)$ be a continuous exponent. Then there exists a constant $C > 0$ such that*

$$\|u - u_D\|_{p(\cdot)} \leq C \|\nabla u\|_{p(\cdot)}$$

for every $u \in W^{1,p(\cdot)}(\mathbb{R}^n)$.

Proof. Since p is continuous we find for every $x \in \bar{D}$ a constant $r(x) > 0$ such that either

$$p_{B(x,r(x)) \cap D}^+ \leq (p_{B(x,r(x)) \cap D}^-)^* \quad \text{or} \quad p_{B(x,r(x)) \cap D}^- \geq n.$$

Since \bar{D} is compact it is possible to find finite covering of D with balls $B(x, r(x))$. It is easy to see that each $B(x, r(x)) \cap D$ is a John domain and hence the corollary follows by Theorem 2.6. \square

Sometimes it is useful to have better control over the constant in the Poincaré inequality as the domain D changes than we have in (1). In the fixed exponent case the constant of the Poincaré inequality is $C \text{diam}(D)$. We show that this kind of constant is also possible for variable exponent Sobolev spaces. The price we have to pay for this is that the exponent p has to satisfy a much stronger condition in Theorem 2.8 than in Theorem 2.2; in Theorem 2.2 the exponent p could be discontinuous even in every point, but in Theorem 2.8 the exponent is 0-Hölder continuous.

Theorem 2.8. *Let $D \subset \mathbb{R}^n$ be a bounded uniform domain. Let $p: D \rightarrow \mathbb{R}$ be such that $1 < p_D^- \leq p_D^+ < \infty$. Assume that there exists a constant $C > 0$ such that*

$$|p(x) - p(y)| \leq \frac{C}{-\log|x - y|} \tag{10}$$

for every $x, y \in D$ with $|x - y| \leq \frac{1}{2}$. Then the inequality

$$\|u - u_D\|_{p(\cdot)} \leq C \text{diam}(D) \left(1 + \max \left\{ |D|^{1/p_D^+ - 1/p_D^-}, |D|^{1/p_D^- - 1/p_D^+} \right\}\right) \|\nabla u\|_{p(\cdot)}, \tag{11}$$

holds for every $u \in W^{1,p(\cdot)}(D)$. Here the constant C depends on the dimension n , the uniform constant of D and p .

Proof. Since $W_0^{1,p(\cdot)}(D) \hookrightarrow W^{1,1}(D)$ we obtain as in the proof of [12, Theorem 11] for every $u \in W^{1,p(\cdot)}(D)$ that

$$|u(x) - u(y)| \leq C|x - y|(\mathbb{M}\nabla u(x) + \mathbb{M}\nabla u(y)) \tag{12}$$

for almost every $x, y \in D$. Here \mathbb{M} is the Hardy-Littlewood maximal operator:

$$\mathbb{M}\nabla u(x) = \sup_{r>0} \int_{B(x,r)} |\nabla u(y)| dy,$$

with the understanding that $\nabla u = 0$ outside D . The constant C depends on the dimension n and the uniform constants of D .

Integrating inequality (12) over y we obtain

$$\begin{aligned} \left| u(x) - \int_D u(y) dy \right| &\leq \int_D |u(x) - u(y)| dy \\ &\leq C \text{diam}(D) \left(\mathbb{M}\nabla u(x) + \int_D \mathbb{M}\nabla u(y) dy \right). \end{aligned}$$

By Hölder's inequality [17, Theorem 2.1] this yields

$$|u(x) - u_D| \leq C \operatorname{diam}(D) \left(\mathbb{M}\nabla u(x) + \frac{C(p)\|1\|_{L^{p'(\cdot)}(D)}}{|D|} \|\mathbb{M}\nabla u\|_{p(\cdot)} \right).$$

Since the previous inequality holds point-wise, it is clear that we have an inequality also for the Lebesgue norms of both sides:

$$\begin{aligned} \|u - u_D\|_{p(\cdot)} &\leq C \operatorname{diam}(D) \left(\|\mathbb{M}\nabla u\|_{p(\cdot)} + \frac{C}{|D|} \|1\|_{p'(\cdot)} \|1\|_{p(\cdot)} \|\mathbb{M}\nabla u\|_{p(\cdot)} \right) \\ &\leq C \operatorname{diam}(D) \left(1 + |D|^{-1} \max\{|D|^{1+1/p_D^+ - 1/p_D^-}, |D|^{1+1/p_D^- - 1/p_D^+}\} \right) \|\mathbb{M}\nabla u\|_{p(\cdot)} \end{aligned}$$

By [5, Theorem 3.5] (see also [7, Remark 2.2]) the Hardy-Littlewood maximal operator is bounded, and so we obtain

$$\|u - u_D\|_{p(\cdot)} \leq C \operatorname{diam}(D) \left(1 + \max\left\{ |D|^{1/p_D^+ - 1/p_D^-}, |D|^{1/p_D^- - 1/p_D^+} \right\} \right) \|\nabla u\|_{p(\cdot)},$$

where the constant C depends on the dimension n , the uniform constant of D and p . □

Remark 2.9. We refer to [19] for basic properties of uniform domains: Every uniform domain is a John domain. Every domain, whose boundary is locally a graph of a Lipschitz continuous function, is a uniform domain. In particular if D is a ball then the constant in (11) depends on the dimension n and p .

Corollary 2.10. *Let p be as in the previous theorem. If B is a ball with $|B| \leq 1$ then*

$$\|u - u_B\|_{p(\cdot)} \leq C \operatorname{diam}(B) \|\nabla u\|_{p(\cdot)},$$

where the constant C does not depend on B .

Proof. Since $|B| \leq 1$ we have

$$\max\left\{ |B|^{1/p_B^+ - 1/p_B^-}, |B|^{1/p_B^- - 1/p_B^+} \right\} = |B|^{1/p_B^+ - 1/p_B^-}.$$

Since p is 0-Hölder continuous, (10), we obtain by [5, Lemma 3.2] that there exists a constant $C > 0$, depending only on the dimension n and the constant in (10), such that $|B|^{1/p_B^+ - 1/p_B^-} \leq C$ for every ball B . Hence $|B| \leq 1$ implies that the constant in (11) is less than $C \operatorname{diam}(B)$. □

3. Continuity

The functions in the classical Sobolev space $W^{1,p}$ are continuous if $p > n$. In this section we consider when functions in variable exponent Sobolev space are continuous.

Theorem 3.1. *Suppose that $p > n$ is locally bounded away from n in D . Then $W^{1,p(\cdot)}(D) \subset C(D)$.*

Proof. Let $x \in D$ and consider the ball $B = B(x, \delta(x)/2)$. Define $q = \text{ess inf}_{y \in B} p(y)$. Then, by [17, Theorem 2.8],

$$W^{1,p(\cdot)}(B) \hookrightarrow W^{1,q}(B) \subset C(B).$$

Therefore every function in $W^{1,p(\cdot)}(D)$ is continuous at x , and since x was arbitrary, the claim follows. \square

The following corollary is immediate.

Corollary 3.2. *Suppose that p is continuous in D . Then $W^{1,p(\cdot)}(D) \subset C(D)$ if $p(x) > n$ for every $x \in D$.*

We next use a classical example to show that the assumption that p is locally bounded away from n in D is not superfluous when p is not continuous.

Example 3.3. Let $B = B(0, 1/16)$, $\varepsilon > 0$ and suppose that

$$p(x) \leq \bar{p}(|x|) = n + (n - 1 - \varepsilon) \frac{\log_2 \log_2(1/|x|)}{\log_2(1/|x|)}$$

for $x \in B \setminus \{0\}$ and $p(0) > n$. We show that then $W^{1,p(\cdot)}(B) \not\subset C(B)$.

Define $u(x) = \cos(\log_2 |\log_2 |x||)$ for $x \in B \setminus \{0\}$ and $u(0) = 0$. Clearly u is not continuous at the origin. So we have to show that $u \in W^{1,p(\cdot)}(B)$. It is clear that u has partial derivatives, except at the origin.

Since u is bounded it follows that $u \in L^{p(\cdot)}(B)$. We next estimate the gradient:

$$|\nabla u(x)| = \left| \sin(\log_2 |\log_2 |x||) \cdot \frac{1}{|x| \log_2 |x|} \right| \leq \left| \frac{1}{|x| \log_2 |x|} \right|.$$

We therefore find that

$$\begin{aligned} \int_B |\nabla u(x)|^{p(x)} dx &\leq \int_B \frac{dx}{(|x| |\log_2 |x||)^{p(x)}} \\ &= \omega_{n-1} \int_0^{1/16} \frac{r^{n-1} dr}{(r |\log_2 r|)^{\bar{p}(r)}} \\ &= \omega_{n-1} \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{r^{n-1} dr}{(r |\log_2 r|)^{\bar{p}(r)}}. \end{aligned}$$

Since $1/(r|\log_2 r|) > 1$ we may increase the exponent \bar{p} for an upper bound. In the annulus $B(0, 2^{-i}) \setminus B(0, 2^{-i-1})$ we have $i \leq \log_2(1/|x|) \leq i + 1$. Since $y \rightarrow \log_2(y)/y$ is decreasing we find that

$$\bar{p}(x) \leq n + (n - 1 - \varepsilon) \frac{\log_2 i}{i}$$

in the same annulus. We can therefore continue our previous estimate by

$$\begin{aligned} \int_B |\nabla u(x)|^{p(x)} dx &\leq \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{r^{n-1} dr}{(r|\log_2 r|)^{n+(n-1-\varepsilon)\log_2(i)/i}} \\ &\leq C \sum_{i=5}^{\infty} \int_{2^{-i-1}}^{2^{-i}} \frac{2^{-i(n-1)} dr}{(i2^{-i})^{n+(n-1-\varepsilon)\log_2(i)/i}} \\ &= C \sum_{i=5}^{\infty} 2^{(n-1-\varepsilon)\log_2(i)} i^{-n-(n-1-\varepsilon)\log_2(i)/i} \\ &= C \sum_{i=5}^{\infty} i^{-1-\varepsilon} i^{-(n-1-\varepsilon)\log_2(i)/i} \leq C \sum_{i=5}^{\infty} i^{-1-\varepsilon} < \infty. \end{aligned}$$

4. Sobolev imbedding theorems

We start by introducing a relative variational $p(\cdot)$ -pseudocapacity, and proving some basic properties for it. This capacity is quite similar to the Sobolev $p(\cdot)$ -capacity studied by P. Harjulehto, P. Hästö, M. Koskenoja and S. Varonen in [13].

Let $F, E \subset \mathbb{R}^n$ be closed disjoint sets and D be a domain in \mathbb{R}^n . The *variational $p(\cdot)$ -pseudocapacity* is defined as

$$\psi_{p(\cdot)}(F, E; D) = \inf_{u \in L(F, E; D)} \|\nabla u\|_{L^{p(\cdot)}(D)},$$

where $L(F, E; D)$ is as before (see Section 2). For $L(F, E; D) = \emptyset$ we define $\psi_{p(\cdot)}(F, E; D) = \infty$. We write $L(E, x; D)$ for $L(F, \{x\}; D)$ etc.

Remark 4.1. Including $C(D)$ in the definition of the capacity is somewhat strange in this context, since we do not, in general, know whether continuous functions are dense in $W^{1,p(\cdot)}(D)$, but see [8]. However, since we are interested in the case when $p > n$, the assumption makes sense, by Theorem 3.1.

The reason for calling the function $\psi_{p(\cdot)}(F, E; D)$ a pseudocapacity is that it is defined as a capacity but using the norm instead of the modular. This corresponds to introducing an exponent $1/p$ to the capacity in the fixed exponent case. Because of this we cannot expect the pseudocapacity to have all the usual properties of a capacity. It nevertheless has many of them:

Theorem 4.2. *Let $F, E \subset \mathbb{R}^n$ be closed sets and D be a domain in \mathbb{R}^n . Then the set function $(F, E) \mapsto \psi_{p(\cdot)}(F, E; D)$ has the following properties:*

- (i) $\psi_{p(\cdot)}(\emptyset, E; D) = 0$.
- (ii) $\psi_{p(\cdot)}(F, E; D) = \psi_{p(\cdot)}(E, F; D)$.
- (iii) *Outer regularity, i. e. $\psi_{p(\cdot)}(F, E_1; D) \leq \psi_{p(\cdot)}(F, E_2; D)$.*
- (iv) *If E is a subset of \mathbb{R}^n , then*

$$\psi_{p(\cdot)}(F, E; D) = \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D).$$

- (v) *If $K_1 \supset K_2 \supset \dots$ are compact, then*

$$\lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D) = \psi_{p(\cdot)}\left(F, \bigcap_{i=1}^{\infty} K_i; D\right).$$

- (vi) *Suppose that $p > n$ is locally bounded away from n . If $E_i \subset \mathbb{R}^n$ for every $i = 1, 2, \dots$, then*

$$\psi_{p(\cdot)}\left(F, \bigcup_{i=1}^{\infty} E_i; D\right) \leq \sum_{i=1}^{\infty} \psi_{p(\cdot)}(F, E_i; D).$$

Proof. Assertion (i) is clear since we may use a constant function. Assertion (ii) is clear since if $u \in L(F, E; D)$ then $1 - u \in L(E, F; D)$. Assertion (iii) follows since $L(F, E_2; D) \subset L(F, E_1; D)$.

Next we prove (iv). It is clear that

$$\psi_{p(\cdot)}(F, E; D) \leq \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D).$$

Let $\varepsilon > 0$. Assume that $u \in L(F, E; D)$ is such that

$$\|\nabla u\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, E; D) + \varepsilon.$$

Since u is continuous, $\{u > 1 - \varepsilon\}$ is an open set containing E . Hence we obtain

$$\begin{aligned} \inf_{\substack{E \subset U \\ U \text{ open}}} \psi_{p(\cdot)}(F, U; D) &\leq \psi_{p(\cdot)}(F, \{u > 1 - \varepsilon\}; D) \\ &\leq \left\| \nabla \min\left\{\frac{u}{1 - \varepsilon}, 1\right\} \right\|_{p(\cdot)} \leq (1 - \varepsilon)^{-1} \|\nabla u\|_{p(\cdot)} \\ &\leq (1 - \varepsilon)^{-1} \psi_{p(\cdot)}(F, E; D) + \frac{\varepsilon}{1 - \varepsilon}. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ yields assertion (iv).

We then prove (v). It is clear that

$$\psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) \leq \lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D)$$

Let $\varepsilon > 0$. Assume that $u \in L(F, \cap_{i=1}^{\infty} K_i; D)$ is such that

$$\|\nabla u\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) + \varepsilon.$$

When i is large the set K_i lies in the closed set $\{u \geq 1 - \varepsilon\}$; therefore

$$\begin{aligned} \lim_{i \rightarrow \infty} \psi_{p(\cdot)}(F, K_i; D) &\leq \psi_{p(\cdot)}(F, \{u \geq 1 - \varepsilon\}; D) \\ &\leq \left\| \nabla \min\left\{\frac{u}{1 - \varepsilon}, 1\right\} \right\|_{p(\cdot)} \leq (1 - \varepsilon)^{-1} \|\nabla u\|_{p(\cdot)} \\ &\leq (1 - \varepsilon)^{-1} \psi_{p(\cdot)}(F, \cap_{i=1}^{\infty} K_i; D) + \frac{\varepsilon}{1 - \varepsilon}. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ yields assertion (v).

To prove (vi) let $\varepsilon > 0$ and choose functions $u_i \in L(F, E_i; D)$ such that

$$\|\nabla u_i\|_{p(\cdot)} \leq \psi_{p(\cdot)}(F, E_i; D) + \varepsilon/2^i,$$

for $i = 1, \dots$. Let $v_i = u_1 + \dots + u_i$. Then (v_i) is a Cauchy sequence, and so it converges to a function $v \in W^{1,p(\cdot)}(D)$. Define $\tilde{v}(x) = \min\{v(x), 1\}$, so that $|\tilde{v}| \in L^{p(\cdot)}(D)$ by [13, Theorem 2.2]. It is clear that $\tilde{v}|_{F \cap D} = 0$ and $\tilde{v}|_{E \cap D} = 1$, where $E = \cup E_i$. Since $p > n$ is locally bounded away from n , it follows from Theorem 3.1 that every function in $W^{1,p(\cdot)}(D)$ is continuous, and so we have $\tilde{v} \in L(F, \cup E_i; D)$, from which the claim easily follows, since

$$\|\nabla \tilde{v}\|_{p(\cdot)} \leq \sum_{i=1}^{\infty} \|\nabla u_i\|_{p(\cdot)} \leq \sum_{i=1}^{\infty} \psi_{p(\cdot)}(F, E_i; D) + \varepsilon. \quad \square$$

Using the pseudocapacity we can start our study of Sobolev-type imbeddings. The following result is the direct generalization of [20, 5.1.1, Theorem 1].

Theorem 4.3. *If $p^+ < \infty$, then the following two conditions are equivalent:*

- (i) $W^{1,p(\cdot)}(D) \cap C(D) \hookrightarrow L^\infty(D)$.
- (ii) *There exist $r, k > 0$ such that $\psi_{p(\cdot)}(\overline{D} \setminus B(x, r), x; D) \geq k$ for every $x \in D$.*

Proof. Suppose that (2) holds, with constants $r, k > 0$. Let $u \in W^{1,p(\cdot)}(D) \cap C(D)$ and let $y \in D$ be a point with $u(y) \neq 0$. Fix a function $\eta \in C_0^\infty(B(0, 1))$ with $0 \leq \eta \leq 1$ and $\eta(0) = 1$. Define $v(x) = \eta((x - y)/r)u(x)/u(y)$. It is clear that $v \in W^{1,p(\cdot)}(D)$

and since $v(y) = 1$ and $v(x) = 0$ for $x \notin B(y, r)$ we see that $v \in L(\overline{D} \setminus B(y, r), y; D)$. It follows that

$$k \leq \psi_{p(\cdot)}(\overline{D} \setminus B(y, r), y; D) \leq \|\nabla v\|_{p(\cdot)}.$$

Then we calculate that

$$\begin{aligned} k|u(y)| &\leq \|\nabla(u(x)\eta((x-y)/r))\|_{p(x)} \\ &\leq \sup_{x \in D} \eta(x)\|\nabla u\|_{p(\cdot)} + \frac{1}{r} \sup_{x \in D} \nabla \eta(x)\|u\|_{p(\cdot)} \\ &\leq \max \left\{ \sup_{x \in D} \eta(x), \frac{1}{r} \sup_{x \in D} \nabla \eta(x) \right\} \|u\|_{1,p(\cdot)}, \end{aligned}$$

so that $|u(y)|$ is bounded by a constant independent of y .

Suppose conversely that (1) holds and let C be a constant such that $\|u\|_\infty \leq C\|u\|_{1,p(\cdot)}$ for all $u \in W^{1,p(\cdot)}(D)$. For functions in $v \in L(\overline{D} \setminus B(x, r), x; D)$ this gives

$$1 = \|v\|_\infty \leq C\|v\|_{1,p(\cdot)} \leq C(\|\chi_{B(x,r)}\|_{p(\cdot)} + \|\nabla v\|_{p(\cdot)}).$$

Since $p^+ < \infty$ we can choose r small enough that $\|\chi_{B(x,r)}\|_{p(\cdot)} \leq 1/(2C)$. For such r we have $\|\nabla v\|_{p(\cdot)} \geq 1/(2C)$. It follows that

$$\psi_{p(\cdot)}(\overline{D} \setminus B(x, r), x; D) = \inf_{u \in L(\overline{D} \setminus B(x,r),x;D)} \|\nabla u\|_{p(\cdot)} \geq 1/(2C)$$

for the same r . □

Remark 4.4. Since we do not know whether $C^\infty(D)$ is dense in $W^{1,p(\cdot)}(D)$ we have only proved the theorem for continuous functions in $W^{1,p(\cdot)}(D)$. If p is such that $C(D)$ is dense in $W^{1,p(\cdot)}(D)$, for instance if p is locally bounded above n , then we may replace condition (1) by $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$.

Define $D = B(1/16) \setminus \{0\}$ and let p be as in Example 3.3. Then the standard example $u(x) = \log|\log(x)|$ shows that $W^{1,p(\cdot)}(D) \not\hookrightarrow L^\infty$, the calculations being as in the theorem. We next show that the exponent p from the theorem is almost as good as possible. We need the following lemma.

Lemma 4.5. *Let $\{a_i\}$ be a partition of unity and $k > m - 1$. Then*

$$\sum_{i=0}^\infty a_i^m i^k \geq \left(\sum_{i=0}^\infty i^{-k/(m-1)} \right)^{1-m}.$$

Proof. Fix an integer i and consider the function

$$a \mapsto (a_i + a)^m i^k + (a_{i+1} - a)^m (i + 1)^k,$$

for $-a_i < a < a_{i+1}$. We find that this function has a minimum at $a = 0$ if and only if

$$\left(\frac{a_i}{a_{i+1}}\right)^{m-1} = \left(\frac{i+1}{i}\right)^k. \tag{13}$$

Let $\{a_i\}$ be a minimal sequence, so that (13) holds for every $i \geq 0$. This partition is given by $a_i = i^{-k/(m-1)}a_0$ for $i > 0$ and $a_0 = (\sum i^{-k/(m-1)})^{-1}$ and so we easily calculate the lower bound as given in the lemma. \square

We next give a simple sufficient condition for the imbedding $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$ to hold in a regular domain:

Theorem 4.6. *Suppose that D satisfies a uniform interior cone condition. If $p^+ < \infty$ and*

$$p(x) \geq n + (n - 1 + \varepsilon) \frac{\log_2 \log_2(c/\delta(x))}{\log_2(c/\delta(x))}$$

for some fixed $0 < \varepsilon < n - 1$ and constant $c > 0$ then $W^{1,p(\cdot)}(D) \hookrightarrow L^\infty(D)$. Here $\delta(x)$ denotes the distance of x from the boundary of D

Proof. Note first that the claim trivially holds in compact subsets of D which satisfy the cone condition, since p is bounded away from n in such sets. Therefore it suffices to prove the claim for $\delta(x)$ less than some constant.

By the uniform interior cone condition there exist real values $0 < \alpha < \pi/2$ and $r > 0$ and a unit vector field v_x such that for every $x \in D$ the cone

$$C_x = \{y \in B(x, r) : \langle x - y, v_x \rangle > |x - y| \cos \alpha\}$$

lies completely in D , where $\langle \cdot, \cdot \rangle$ denotes the usual inner product.

Fix $z \in D$. Consider the cone

$$C = \{y \in B(z, r/2) : \langle z - y, v_z \rangle > |z - y| \cos(\alpha/3)\}$$

and, for $i = 2, 3, \dots$, the annuli

$$A_i = (B(z, 2^{-i+1}r) \setminus B(z, 2^{-i}r)) \cap C.$$

To simplify notation let us assume that $z = 0$, $r = 1$ and $v_z = e_1$; the proof in the general case is essentially identical. Since $A_i \subset C \subset D$ we have $d(A_i, \partial D) \geq d(A_i, \partial C)$. We can estimate the latter distance as shown in Figure 1. This gives $d(A_i, \partial D) \geq 2^{-i} \sin(\alpha/3)$ so that

$$p(x) \geq n + (n - 1 + \varepsilon) \frac{\log_2(i + c)}{i + c}$$




Figure 1: The cone C and the distance to the boundary

for $x \in A_i$ and some c depending on α . Let us define $q_i = n + (n - 1 + \varepsilon) \frac{\log_2(i+c)}{i+c}$ and a new variable exponent by

$$q(x) = \begin{cases} q_i & \text{if } x \in A_i \text{ for some } i \\ p(x) & \text{otherwise} \end{cases}$$

By Theorem 4.3 we know that it suffices to find a lower bound for $\|\nabla u\|_{1,p(\cdot)}$ with $u \in L(\overline{D} \setminus B(0, r), 0; D)$ since, by Theorem 3.1, $W^{1,p(\cdot)}(D) \subset C(D)$. Since $\|u\|_{1,p(\cdot)} \geq c\|u\|_{1,q(\cdot)}$, we see that it suffices to estimate $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap D)$ for small R in order to prove the theorem. Moreover, by monotony, we need only consider $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C)$. For every function $u \in W^{1,q(\cdot)}(C)$ we have

$$\|u\|_{1,q(\cdot)} \geq \min\{1, \varrho_{1,q(\cdot)}(u)\},$$

by [17, Theorem 2.8]. Therefore we see that it suffices to show that $\varrho_{1,q(\cdot)}(u) > c$ for every $u \in L(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C)$ in order to get $\psi_{q(\cdot)}(\overline{D} \setminus B(0, R), 0; B(0, R) \cap C) \geq \min\{1, c\} > 0$, which will complete the proof.

It is clear that $|\nabla u| \geq |\partial u / \partial r|$, the radial component of the gradient, so that

$$\int_{A_i} |\nabla u|^{q_i} dx \geq \int_{A_i} \left| \frac{\partial u}{\partial r} \right|^{q_i} dx.$$

It is then easy to see that the function minimizing the sum over the integrals should depend only on the distance from the origin, not on the direction. For such a function let us denote the value at any point of distance 2^{-i} from the origin by v_i .

Consider then a function v which equals v_{i-1} on $S(0, 2^{-i+1})$ and v_i on $S(0, 2^{-i})$. Using Lemma 2.4 we find that

$$\begin{aligned} \int_{A_i} |\nabla v|^{q_i} dx &\geq (v_{i-1} - v_i)^{q_i} \text{cap}_{q_i}(\mathbb{R}^n \setminus B(0, 2^{-i+1}), B(0, 2^{-i})) \\ &= (v_{i-1} - v_i)^{q_i} \omega_{n-1} \left(\frac{q_i - n}{q_i - 1}\right)^{q_i - 1} (2^{(q_i - n)/(q_i - 1)} - 1)^{1 - q_i} 2^{i(q_i - n)} \\ &\geq c(v_{i-1} - v_i)^{q_i} 2^{i(q_i - n)}, \end{aligned}$$

where the constant c does not depend on q_i . It follows that

$$\varrho_{1,q(\cdot)}(v) \geq \sum_{i=2}^{\infty} \int_{A_i} |\nabla u|^{q_i} dx \geq c \sum_{i=2}^{\infty} (v_{i-1} - v_i)^{q_i} 2^{i(q_i - n)}.$$

Since the lower bound depends only on the v_i , we see that

$$\inf_{u \in L} \varrho_{1,q(\cdot)}(u) \geq c \inf_{\{v_i\}} \sum_{i=2}^{\infty} (v_{i-1} - v_i)^{q_i} 2^{i(q_i - n)},$$

where the second infimum is over sequences $\{v_i\}$ with $v_i \leq v_{i-1}$, $v_0 = 1$ and $\lim_{i \rightarrow \infty} v_i = 0$. Let us set $a_i = v_{i-1} - v_i$ so that $a_i \geq 0$ and $\sum a_i = 1$. Then we need to estimate

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i - n)},$$

with the infimum over partitions of unity $\{a_i\}$. Let N be such that

$$\frac{\varepsilon}{3} \geq q_i - n = (n - 1 + \varepsilon) \frac{\log_2(i + c)}{i + c} \geq (n - 1 + \varepsilon/2) \frac{\log_2(i)}{i}$$

for $i \geq N$. Note that such an N can be chosen independent of z . Since $a_i \leq 1$ we have $a_i^{q_i} \geq a_i^{n+\varepsilon/3}$ for such terms. Then we find that

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i - n)} \geq \inf_{\{a_i\}} \sum_{i=2}^{N-1} a_i^{q_i} 2^{i(q_i - n)} + \sum_{i=N}^{\infty} a_i^{n+\varepsilon/3} i^{n-1+\varepsilon/2}.$$

The first sum on the left-hand-side is finite, hence

$$\sum_{i=2}^{N-1} a_i^{q_i} 2^{i(q_i - n)} \geq \sum_{i=2}^{N-1} a_i^q \geq N^{1-q} \left(\sum_{i=2}^{N-1} a_i\right)^q,$$

where $q = \max_{2 \leq i \leq N-1} q_i$. It follows from Lemma 4.5 that

$$\sum_{i=N}^{\infty} a_i^{n+\varepsilon/3} i^{n-1+\varepsilon/2} \geq c \left(\sum_{i=N}^{\infty} a_i\right)^{n+\varepsilon/3}.$$

Combining these estimates we see that

$$\inf_{\{a_i\}} \sum_{i=2}^{\infty} a_i^{q_i} 2^{i(q_i-n)} \geq N^{1-q} \left(\sum_{i=2}^{N-1} a_i \right)^q + c \left(\sum_{i=N}^{\infty} a_i \right)^{n+\varepsilon/3}$$

is uniformly bounded from below by a positive constant, since the sum of the a_i 's is 1. We have thus shown that the condition of Theorem 4.3 holds, which concludes the proof. \square

References

- [1] S. M. Buckley and P. Koskela, *Sobolev-Poincaré implies John*, Math. Res. Lett. **2** (1995), 577–593.
- [2] ———, *Criteria for imbeddings of Sobolev-Poincaré type*, Internat. Math. Res. Notices (1996), 881–901.
- [3] B. Bojarski, *Remarks on Sobolev imbedding inequalities*, Complex Analysis, Joensuu 1987, Lecture Notes in Math., vol. 1351, Springer, Berlin, 1988, pp. 52–68.
- [4] D. Cruz-Uribe, A. Fiorenza, and C. J. Neugebauer, *The maximal function on variable L^p spaces*, Ann. Acad. Sci. Fenn. Math. **28** (2003), 223–238.
- [5] L. Diening, *Maximal function on generalized Lebesgue spaces $L^{p(\cdot)}$* , Math. Inequal. Appl. (to appear).
- [6] ———, *Riesz Potential and Sobolev Embeddings of generalized Lebesgue and Sobolev Spaces $L^{p(\cdot)}$ and $W^{k,p(\cdot)}$* , Math. Nachr. (to appear).
- [7] L. Diening and M. Růžička, *Calderón-Zygmund operators on generalized Lebesgue spaces $L^{p(\cdot)}$ and problems related to fluid dynamics*, J. Reine Angew. Math. **563** (2003), 197–220.
- [8] D. E. Edmunds and J. Rákosník, *Density of smooth functions in $W^{k,p(x)}(\Omega)$* , Proc. Roy. Soc. London Ser. A **437** (1992), 229–236.
- [9] ———, *Sobolev embeddings with variable exponent*, Studia Math. **143** (2000), 267–293.
- [10] ———, *Sobolev embeddings with variable exponent. II*, Math. Nachr. **246/247** (2002), 53–67.
- [11] X. Fan, J. Shen, and D. Zhao, *Sobolev embedding theorems for spaces $W^{k,p(x)}(\Omega)$* , J. Math. Anal. Appl. **262** (2001), 749–760.
- [12] P. Hajlasz and O. Martio, *Traces of Sobolev functions on fractal type sets and characterization of extension domains*, J. Funct. Anal. **143** (1997), 221–246.
- [13] P. Harjulehto, P. Hästö, M. Koskenoja, and S. Varonen, *Sobolev capacity on the space $W^{1,p(\cdot)}(\mathbb{R}^n)$* , J. Funct. Spaces Appl. **1** (2003), 17–33.
- [14] ———, *Dirichlet energy integral and Sobolev spaces with zero boundary values* (preprint. Available in <http://www.math.helsinki.fi/analysis/varsobgroup>).
- [15] J. Heinonen, T. Kilpeläinen, and O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1993, ISBN 0-19-853669-0.
- [16] V. Kokilasvili and S. Samko, *Maximal and fractional operators in weighted $L^{p(x)}$ spaces*, Rev. Mat. Iberoamericana (to appear).
- [17] O. Kováčik and J. Rákosník, *On spaces $L^{p(x)}$ and $W^{k,p(x)}$* , Czechoslovak Math. J. **41(116)** (1991), 592–618.

- [18] J. Malý and W. P. Ziemer, *Fine regularity of solutions of elliptic partial differential equations*, Mathematical Surveys and Monographs, vol. 51, American Mathematical Society, Providence, RI, 1997, ISBN 0-8218-0335-2.
- [19] O. Martio and J. Sarvas, *Injectivity theorems in plane and space*, Ann. Acad. Sci. Fenn. Ser. A I Math. **4** (1979), 383–401.
- [20] V. G. Maz'ja, *Sobolev spaces*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1985, ISBN 3-540-13589-8.
- [21] L. Pick and M. Růžička, *An example of a space $L^{p(x)}$ on which the Hardy-Littlewood maximal operator is not bounded*, Expo. Math. **19** (2001), 369–371.
- [22] Michael Růžička, *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics, vol. 1748, Springer-Verlag, Berlin, 2000, ISBN 3-540-41385-5.