

Anisotropic Besov Spaces and Approximation Numbers of Traces on Related Fractal Sets

Erika TAMÁSI

Mathematical Institute
Friedrich-Schiller-University Jena
D-07737 Jena — Germany
tamasi@minet.uni-jena.de

Received: December 25, 2005

Accepted: January 13, 2006

ABSTRACT

This paper deals with approximation numbers of the compact trace operator of an anisotropic Besov space into some L_p -space,

$$\mathrm{tr}_\Gamma : B_{pp}^{s,a}(\mathbb{R}^n) \hookrightarrow L_p(\Gamma), \quad s > 0, \quad 1 < p < \infty,$$

where Γ is an anisotropic d -set, $0 < d < n$. We also prove homogeneity estimates, a homogeneous equivalent norm and the localization property in $B_{pp}^{s,a}$.

Key words: anisotropic function spaces, fractals, approximation numbers, traces.

2000 Mathematics Subject Classification: 46E35, 42B35, 42C40.

1. Introduction

The theory of the anisotropic spaces has been developed from the very beginning parallel to the theory of isotropic function spaces. We refer in particular to the Russian school and works of S. M. Nikol'skiĭ, O. V. Besov, V. P. Il'in [1, 11].

Let $1 < p < \infty$ and (s_1, \dots, s_n) be an n -tuple of natural numbers. Then

$$W_p^{s,a}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \|f\|_{L_p(\mathbb{R}^n)} + \sum_{k=1}^n \left\| \frac{\partial^{s_k} f}{\partial x_k^{s_k}} \right\|_{L_p(\mathbb{R}^n)} < \infty \right\}$$

is the classical anisotropic Sobolev space on \mathbb{R}^n . It is obvious that unlike in case of the usual (isotropic) Sobolev space ($s_1 = \dots = s_n$) the smoothness properties of an

element from $W_p^{s,a}(\mathbb{R}^n)$ depend on the chosen direction in \mathbb{R}^n . The number s , defined by

$$\frac{1}{s} = \frac{1}{n} \left(\frac{1}{s_1} + \dots + \frac{1}{s_n} \right),$$

is usually called the *mean smoothness*, and $a = (a_1, \dots, a_n)$,

$$a_1 = \frac{s}{s_1}, \dots, a_n = \frac{s}{s_n}$$

characterizes the *anisotropy*. Similar to the isotropic situation the more general anisotropic Bessel potential spaces (fractional Sobolev spaces) $H_p^{s,a}(\mathbb{R}^n)$, where $1 < p < \infty$, $s \in \mathbb{R}$ and $a = (a_1, \dots, a_n)$ is a given anisotropy, fit in the scales of anisotropic Besov spaces $B_{pq}^{s,a}(\mathbb{R}^n)$, and anisotropic Triebel-Lizorkin spaces $F_{pq}^{s,a}(\mathbb{R}^n)$, respectively. It is well known that this theory has a more or less complete counterpart to the basic facts (definitions, description via differences and derivatives, elementary properties, embeddings for different metrics, interpolation) of isotropic spaces $B_{pq}^s(\mathbb{R}^n)$ and $F_{pq}^s(\mathbb{R}^n)$. We shall use the Fourier-analytical definition of $B_{pq}^{s,a}(\mathbb{R}^n)$, $F_{pq}^{s,a}(\mathbb{R}^n)$, where any function $f \in \mathcal{S}'(\mathbb{R}^n)$ is decomposed in a sum of entire analytic functions $(\varphi_j \hat{f})^\vee$ and this decomposition, measured in ℓ_q and $L_p(\mathbb{R}^n)$, respectively, is used to introduce the spaces. This concept goes back to [15, 16], see [13, chapter 4].

Our main aim in the present paper is to prove an anisotropic counterpart to the isotropic results, see [21].

As a first goal of this paper we define the anisotropic d -set as follows:

Let $0 < d < n$, a an anisotropy. Then $\Gamma \subset \mathbb{R}^n$ is called an anisotropic d -set if there exists a positive Radon measure μ with $\text{supp } \mu = \Gamma$ and

$$\mu(B^a(\gamma, r)) \sim r^d, \quad 0 < r < 1,$$

where $B^a(\gamma, r) = \{y \in \mathbb{R}^n : |y - \gamma|_a \leq r\}$ is an anisotropic ball and $\gamma \in \Gamma$.

We study the existence and properties of the trace operator tr_Γ ,

$$\text{tr}_\Gamma : B_{pp}^{s,a}(\mathbb{R}^n) \hookrightarrow L_p(\Gamma) \tag{1}$$

where Γ is an anisotropic d -set. It turns out that tr_Γ according to (1) exists if, and is compact if

$$\sum_{j \in \mathbb{N}_0} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} < \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1, \tag{2}$$

where $\mu_j = \sup_{m \in \mathbb{Z}^n} \mu(Q_{jm}^a)$, and Q_{jm}^a are rectangles centered at $2^{-j}m$ and with side length $2^{-ja_1}, \dots, 2^{-ja_n}$. If we can strengthen (2) by

$$\sum_{j \geq J} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} \sim 2^{-Jp'(s-\frac{n}{p})} \mu_J^{p'-1}, \quad J \in \mathbb{N}_0,$$

then one obtains for the approximation numbers a_k of the compact operator tr_Γ according to (1)

$$a_k(\text{tr}_\Gamma : B_{pp}^{s,a}(\mathbb{R}^n) \hookrightarrow L_p(\Gamma)) \sim k^{\frac{1}{d}(\frac{n}{p}-s)-\frac{1}{p}}, \quad \frac{n}{p} \geq s > \frac{n-d}{p},$$

as in the isotropic case, see [21]. In order to show the above result we prove, in addition, some important properties of spaces $B_{pq}^{s,a}(\mathbb{R}^n)$, with $0 < p \leq \infty, 0 < q \leq \infty$, and $s > n(\frac{1}{p} - 1)_+$, which might be of self-contained interest:

(i) We obtain the homogeneity estimate

$$\|f(R \cdot) | B_{pq}^{s,a}(\mathbb{R}^n)\| \leq cR^{s-\frac{n}{p}} \|f | B_{pq}^{s,a}(\mathbb{R}^n)\|$$

for all $f \in B_{pq}^{s,a}(\mathbb{R}^n)$ and $R \geq 1$.

(ii) We show that

$$\|(\varphi \hat{f})^\vee | L_p\| + \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|(\varphi_j^a \hat{f})^\vee | L_p\|^q \right)^{1/q}$$

and

$$\|f | L_p\| + \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|(\varphi_j^a \hat{f})^\vee | L_p\|^q \right)^{1/q}$$

are equivalent quasi-norms in $B_{pq}^{s,a}(\mathbb{R}^n)$.

(iii) Finally we prove the localization property of $B_{pp}^{s,a}(\mathbb{R}^n)$, that is

$$2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p \right)^{1/p} \|f | B_{pp}^{s,a}(\mathbb{R}^n)\| \sim \|f_j^a | B_{pp}^{s,a}(\mathbb{R}^n)\|,$$

where

$$f_j^a(x) = \sum_{k \in \mathbb{Z}^n} c_k f(2^{(j+1)a}(x - x_{j,k}^a)), \quad c_k \in \mathbb{C}, \quad j \in \mathbb{N},$$

and f is a product of one-dimensional functions

$$f(2^{(j+1)a}(x - x_{j,k}^a)) = \prod_{m=1}^n f_m(2^{(j+1)a_m}(x_m - 2^{-ja_m}k_m)).$$

The plan of the paper is the following. In the second section we give the definition and some important properties of anisotropic Besov spaces. In section 3 we introduce the anisotropic d -set and we formulate our main result. In the last section we collect the proofs and give a description of the wavelet frames according to [9].

2. Anisotropic Besov spaces

2.1. General notation

As usual, \mathbb{R}^n denotes the n -dimensional real Euclidean space, \mathbb{N} the collection of all natural numbers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, \mathbb{C} stands for the complex numbers, and \mathbb{Z}^n means the lattice of all points in \mathbb{R}^n with integer-valued components. We use the equivalence “ \sim ” in $\varphi(x) \sim \psi(x)$ always to mean that there are two positive numbers c_1 and c_2 such that

$$c_1 \varphi(x) \leq \psi(x) \leq c_2 \varphi(x)$$

for all admitted values of x , where φ, ψ are non-negative functions. If $a \in \mathbb{R}$ then $a_+ := \max(a, 0)$. Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ be a multi-index, then

$$|\alpha| = \alpha_1 + \dots + \alpha_n, \quad \alpha! = \alpha_1! \cdots \alpha_n!, \quad \alpha \in \mathbb{N}_0^n,$$

the derivatives D^α have the usual meaning, x^α means $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, and $\alpha\gamma = \alpha_1\gamma_1 + \dots + \alpha_n\gamma_n$, $\gamma \in \mathbb{R}^n$, stands for the scalar product in \mathbb{R}^n .

Given two quasi-Banach spaces X and Y , we write $X \hookrightarrow Y$ if $X \subset Y$ and the natural embedding of X in Y is continuous. All unimportant positive constants will be denoted by c , occasionally with additional subscripts within the same formula. We shall mainly deal with function spaces on \mathbb{R}^n ; so for convenience we shall usually omit the “ \mathbb{R}^n ” from their notation, if there is no danger of confusion.

2.2. Definitions

Let $a = (a_1, \dots, a_n)$ be a fixed n -tuple of positive numbers with $a_1 + \dots + a_n = n$, then we call a an *anisotropy*. We shall denote $a_{\min} = \min\{a_i : 1 \leq i \leq n\}$ and $a_{\max} = \max\{a_i : 1 \leq i \leq n\}$. If $a = (1, \dots, 1)$ we speak about the “isotropic case”.

The action of $t \in [0, \infty)$ on $x \in \mathbb{R}^n$ is defined by the formula

$$t^a x = (t^{a_1} x_1, \dots, t^{a_n} x_n). \quad (3)$$

For $t > 0$ and $s \in \mathbb{R}$ we put $t^{sa} x = (t^s)^a x$. In particular we write $t^{-a} x = (t^{-1})^a x$ and $2^{-ja} x = (2^{-j})^a x$.

Definition 2.1. An *anisotropic distance function* is a continuous function $u : \mathbb{R}^n \rightarrow \mathbb{R}$ with the properties $u(x) > 0$ if $x \neq 0$ and $u(t^a x) = tu(x)$ for all $t > 0$ and all $x \in \mathbb{R}^n$.

Remark 2.2. It is easy to see that $u_\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$u_\lambda(x) = \left(\sum_{i=1}^n |x_i|^{\frac{\lambda}{a_i}} \right)^{1/\lambda} \quad (4)$$

is an anisotropic distance function for every $0 < \lambda < \infty$, u_2 is usually called the anisotropic distance of x to the origin, see [13, 4.2.1]. It is well known, see [3, 1.2.3]

and [22, 1.4], that any two anisotropic distance functions u and u' are equivalent (in the sense that there exist constants $c, c' > 0$ such that $cu(x) \leq u'(x) \leq c'u(x)$ for all $x \in \mathbb{R}^n$) and that if u is an anisotropic distance function there exists a constant $c > 0$ such that $u(x+y) \leq c(u(x) + u(y))$ for all $x, y \in \mathbb{R}^n$. We are interested to use smooth anisotropic distance functions. Note that for appropriate values of λ one can obtain arbitrary (finite) smoothness of the function u_λ from (4), cf. [3, 1.2.4]. A standard method concerning the construction of anisotropic distance functions in $C^\infty(\mathbb{R}^n \setminus \{0\})$ was given in [14].

For $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, $x \neq 0$, let $|x|_a$ be the unique positive number t such that

$$\frac{x_1^2}{t^{2a_1}} + \dots + \frac{x_n^2}{t^{2a_n}} = 1$$

and let $|0|_a = 0$; then $|\cdot|_a$ is an anisotropic distance function in $C^\infty(\mathbb{R}^n \setminus \{0\})$, see [22, 1.4/3,8]. Plainly, $|x|_a$ is in the isotropic case the Euclidean distance of x to the origin.

Before introducing the function spaces under consideration we need to recall some notation. By \mathcal{S} we denote the Schwartz space of all complex-valued, infinitely differentiable and rapidly decreasing functions on \mathbb{R}^n and by \mathcal{S}' the dual space of all tempered distributions on \mathbb{R}^n . Furthermore, L_p with $0 < p \leq \infty$, stands for the usual quasi-Banach space with respect to the Lebesgue measure, quasi-normed by

$$\|f\|_{L_p} := \left(\int_{\mathbb{R}^n} |f(x)|^p dx \right)^{1/p},$$

with the obvious modification if $p = \infty$. If $\varphi \in \mathcal{S}$ then

$$\hat{\varphi}(\xi) \equiv (\mathcal{F}\varphi)(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-ix\xi} \varphi(x) dx, \quad \xi \in \mathbb{R}^n, \quad (5)$$

denotes the Fourier transform of φ . As usual, $\mathcal{F}^{-1}\varphi$ or $\check{\varphi}$ stands for the inverse Fourier transform, given by the right-hand side of (5) with i in place of $-i$. Here $x\xi$ denotes the scalar product in \mathbb{R}^n . Both \mathcal{F} and \mathcal{F}^{-1} are extended to \mathcal{S}' in the standard way. Let $\varphi \in \mathcal{S}$ be such that

$$\varphi(x) = 1 \quad \text{if} \quad |x|_a \leq 1 \quad \text{and} \quad \text{supp } \varphi \subset \{x \in \mathbb{R}^n : |x|_a \leq 2\}, \quad (6)$$

and for each $j \in \mathbb{N}$ let

$$\varphi_j^a(x) := \varphi(2^{-j}x) - \varphi(2^{-(j+1)}x), \quad x \in \mathbb{R}^n. \quad (7)$$

Then the sequence $(\varphi_j^a)_{j=0}^\infty$, with $\varphi_0 = \varphi$, forms a smooth anisotropic dyadic resolution of unity, cf. [13, 4.2]. Let $f \in \mathcal{S}'$, then the compact support of $\varphi_j^a \hat{f}$ implies by the Paley-Wiener-Schwartz theorem that $(\varphi_j^a \hat{f})^\vee$ is an entire analytic function on \mathbb{R}^n .

Definition 2.3. Assume $0 < p \leq \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, a an anisotropy, and $(\varphi_j^a)_{j=0}^\infty$ a smooth anisotropic dyadic resolution of unity. Then

$$B_{pq}^{s,a} = \left\{ f \in \mathcal{S}' : \|f\|_{B_{pq}^{s,a}} = \left(\sum_{j=0}^\infty 2^{jsq} \|(\varphi_j^a \hat{f})^\vee\|_{L_p}^q \right)^{1/q} < \infty \right\} \quad (8)$$

(with the usual modification if $q = \infty$).

Note that there is a parallel definition for spaces of type $F_{pq}^{s,a}$, $0 < p < \infty$, $0 < q \leq \infty$, $s \in \mathbb{R}$, a an anisotropy, when interchanging the order of ℓ_q - and L_p -quasi-norms in (8). It is obvious that the quasi-norm (8) depends on the chosen system $(\varphi_j^a)_{j \in \mathbb{N}_0}$, but not the space $B_{pq}^{s,a}$ (in the sense of equivalent quasi-norms); therefore we omit in our notation the subscript φ in the sequel. It is well-known that $B_{pq}^{s,a}$ are quasi-Banach spaces (Banach spaces if $p \geq 1$ and $q \geq 1$), and, as in the isotropic case, $\mathcal{S} \hookrightarrow B_{pq}^{s,a} \hookrightarrow \mathcal{S}'$ for all admissible values of p, q, s , see [17, 2.3.3]. If $s \in \mathbb{R}$ and $0 < p < \infty$, $0 < q < \infty$ then \mathcal{S} is dense in $B_{pq}^{s,a}$, see [3, 1.2.10; 22, 3.5]. Note that we indicated the only (formal) difference to the isotropic counterparts of (8) by the additional superscript at the smooth anisotropic dyadic resolution of unity $(\varphi_j^a)_{j=0}^\infty$.

Remark 2.4. A systematic treatment of the theory of (isotropic) B_{pq}^s (and F_{pq}^s) spaces may be found in the monographs [17–20]; see also [4, 12]. A survey on the basic results for the (anisotropic) spaces $B_{pq}^{s,a}$ (and $F_{pq}^{s,a}$) is given in [10, 2.1–2.2; 13, 4.2.1–4.2.4]. In addition to the literature mentioned in our introduction we essentially rely on [7, 8] in the sequel.

For convenience, in case of $p = q$ we shall stick to the notation

$$B_p^{s,a} = B_{pp}^{s,a} \quad \text{where } 0 < p \leq \infty, \quad s \in \mathbb{R},$$

in the sequel.

2.3. Properties

Our aim is to prove some new and important properties of anisotropic Besov spaces and thus to complement results in [2, 7, 13]. This also serves as preparation for our main results in section 3.

Let $\varphi \in \mathcal{S}$ as in section 2.2. In particular we have (6). We extend the definition of φ_j^a from (7) to all integers j . It should be noted that φ_0^a has now a different meaning as in 2.2, i.e., for $f \in \mathcal{S}'$ then we have that

$$f = (\varphi \hat{f})^\vee + \sum_{j=1}^\infty (\varphi_j^a \hat{f})^\vee \quad (\text{convergence in } \mathcal{S}'). \quad (9)$$

As usual, let $\sigma_p = n(\frac{1}{p} - 1)_+$, $0 < p \leq \infty$.

Theorem 2.5. *Let $0 < p \leq \infty$, $0 < q \leq \infty$, $s > \sigma_p$ and a an anisotropy, then*

$$\|(\varphi \hat{f})^\vee | L_p\| + \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|(\varphi_j^a \hat{f})^\vee | L_p\|^q \right)^{1/q} \tag{10}$$

and

$$\|f | L_p\| + \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|(\varphi_j^a f)^\vee | L_p\|^q \right)^{1/q} \tag{11}$$

(modification if $q = \infty$) are equivalent quasi-norms in $B_{pq}^{s,a}$.

Remark 2.6. The quasi-norms of type (10), (11) have a continuous counterpart. We introduce $\rho^a(t\xi) = \varphi(t^a\xi) - \varphi((2t)^a\xi)$ where $t > 0$. Then the counterpart of (10) reads as follows:

Let $0 < p \leq \infty$, $0 < q \leq \infty$, $s > \sigma_p$ and a an anisotropy, then

$$\|f | L_p\| + \left(\int_0^\infty t^{-sq} \|(\rho^a(t \cdot) \hat{f})^\vee | L_p\|^q \frac{dt}{t} \right)^{1/q} \tag{12}$$

(modification if $q = \infty$) is an equivalent quasi-norm in $B_{pq}^{s,a}$.

The proof is given in section 4.1. Now we can extend the well-known homogeneity estimate for B_{pq}^s (see [17, Prop. 3.4.1]) to anisotropic spaces.

Proposition 2.7. *Let $0 < p \leq \infty$, $0 < q \leq \infty$, $s > \sigma_p$ and a an anisotropy. There exists a constant $c > 0$ such that for all $R \geq 1$,*

$$\|f(R \cdot) | B_{pq}^{s,a}\| \leq cR^{s-\frac{n}{p}} \|f | B_{pq}^{s,a}\| \quad \text{for all } f \in B_{pq}^{s,a}. \tag{13}$$

The proof of this proposition is contained in section 4.2.

Our next aim is to extend the localization property, see [4, 2.3.2], to anisotropic spaces.

Let $x_{j,k}^a = 2^{-ja}k$ with $k \in \mathbb{Z}^n$ and $j \in \mathbb{N}$. Let $f \in \mathcal{S}'$ with

$$\text{supp } f \subset Q_b^a = \{x \in \mathbb{R}^n : x = (x_1, x_2, \dots, x_n), |x|_a < b\} \tag{14}$$

where $b > 0$ and $b \leq \frac{1}{4}(2^{a_{\min}} + 1)$. Let

$$f_j^a(x) = \sum_{k \in \mathbb{Z}^n} c_k f(2^{(j+1)a}(x - x_{j,k}^a)), \quad c_k \in \mathbb{C}, \quad j \in \mathbb{N} \tag{15}$$

where f is a product of one-dimensional functions,

$$f(2^{(j+1)a}(x - x_{j,k}^a)) = \prod_{m=1}^n f_m(2^{(j+1)a_m}(x_m - 2^{-ja_m}k_m)) \tag{16}$$

and $f_1(y) = \dots = f_n(y)$ where $y \in \mathbb{R}$.

Theorem 2.8. *Let $s > \sigma_p$, $0 < p \leq \infty$, a an anisotropy and $0 < b \leq \frac{1}{4}(2^{a_{\min}} + 1)$. There exist two constants $c' > 0$ and $c'' > 0$ such that for all $f \in B_p^{s,a}$ with $\text{supp } f \subset Q_b^a$ and all $j \in \mathbb{N}$ and all f_j^a given by (15)*

$$c' \|f_j^a | B_p^{s,a}\| \leq 2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p \right)^{1/p} \|f | B_p^{s,a}\| \leq c'' \|f_j^a | B_p^{s,a}\|. \tag{17}$$

We prove this result in section 4.3.

3. Traces and approximation numbers

3.1. General measures

Let μ be a positive Radon measure in \mathbb{R}^n with compact support

$$\Gamma = \text{supp } \mu, \quad 0 < \mu(\mathbb{R}^n) < \infty, \quad |\Gamma| = 0, \tag{18}$$

where $|\Gamma|$ is the Lebesgue measure of Γ . For $1 \leq p < \infty$ we denote by $L_p(\Gamma) = L_p(\Gamma, \mu)$ the usual complex Banach space, normed by

$$\|f | L_p(\Gamma, \mu)\| = \left(\int_{\mathbb{R}^n} |f(x)|^p \mu(dx) \right)^{1/p} = \left(\int_{\Gamma} |f(\gamma)|^p \mu(d\gamma) \right)^{1/p}.$$

Since μ is Radon, $S(\mathbb{R}^n) | \Gamma$ is dense in $L_p(\Gamma)$. If $\varphi \in \mathcal{S}$ then $\text{tr}_\Gamma \varphi = \varphi | \Gamma$ makes sense pointwise. If $1 < p < \infty$, $0 < q \leq \infty$ and $s > 0$ then the embedding $\text{tr}_\Gamma B_{pq}^{s,a} \hookrightarrow L_p(\Gamma)$ must be understood as follows: there exists a positive number $c > 0$ such that for any $\varphi \in \mathcal{S}$,

$$\|\text{tr}_\Gamma \varphi | L_p(\Gamma)\| \leq c \|\varphi | B_{pq}^{s,a}\|.$$

Since \mathcal{S} is dense in $B_{pq}^{s,a}$ for $0 < p, q < \infty$ this inequality can be extended by completion to any $f \in B_{pq}^{s,a}$ and the resulting function is denoted by $\text{tr}_\Gamma f$ and the independence of $\text{tr}_\Gamma f$ from the approximating sequence is shown in the standard way.

In the sequel, we only consider the case $p = q$. We proceed in a way similar to [21], dealing with the isotropic case. Let Q_{jm}^a be the rectangles in \mathbb{R}^n with side length $2^{-ja_1}, \dots, 2^{-ja_n}$ and centered at $2^{-ja}m$ where $m \in \mathbb{Z}^n$ and $j \in \mathbb{N}_0$. Let

$$\mu_j = \sup_{m \in \mathbb{Z}^n} \mu(Q_{jm}^a), \quad j \in \mathbb{N}_0. \tag{19}$$

Proposition 3.1. *Let*

$$1 < p < \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1, \quad s > 0.$$

Let μ be the Radon measure in \mathbb{R}^n with

$$\Gamma = \text{supp } \mu \text{ compact}, \quad 0 < \mu(\mathbb{R}^n) < \infty, \quad |\Gamma| = 0, \tag{20}$$

and

$$\sum_{j \in \mathbb{N}_0} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} < \infty \quad \text{where} \quad \mu_j = \sup_{m \in \mathbb{Z}^n} \mu(Q_{jm}^a). \tag{21}$$

Then

$$\text{tr}_\Gamma : B_p^{s,a}(\mathbb{R}^n) \hookrightarrow L_p(\Gamma) \tag{22}$$

exists and is compact. Furthermore there is a constant c (depending on p and s) such that for all measures μ with (20), (21),

$$\|\text{tr}_\Gamma\| \leq c \left(\sum_{j \in \mathbb{N}_0} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} \right)^{\frac{1}{p'}}. \tag{23}$$

The result above is the anisotropic version of [21, Proposition 3]. The proof can be found in section 4.5.

In the following we recall the concept of approximation numbers. Let A and B be two Banach spaces and let $T \in L(A, B)$. Then for any $k \in \mathbb{N}$ the k th approximation number $a_k(T)$ of T is given by

$$a_k(T) = \inf \{ \|T - L\| : L \in L(A, B), \text{rank } L < k \}, \tag{24}$$

where $\text{rank } L$ is the dimension of the range of L . These numbers have various properties given in the following lemma.

Lemma 3.2. *Let A and B be two Banach spaces and let $T, S \in L(A, B)$.*

- (i) $\|T\| = a_1(T) \geq a_2(T) \geq \dots \geq 0$.
- (ii) For all $n, m \in \mathbb{N}$,

$$a_{m+n-1}(S + T) \leq a_m(S) + a_n(T).$$

- (iii) For all $n, m \in \mathbb{N}$, and $R \in L(B, C)$

$$a_{m+n-1}(RT) \leq a_m(R)a_n(T).$$

- (iv) $a_n(T) = 0 \iff \text{rank } T < n$.

This is a well-known result and can be found for instance in [4, 1.3.1] and [5, II]. Let $T = \text{tr}_\Gamma$ according to Proposition 3.1. We strengthen (21) by

$$\sum_{j \geq J} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} \sim 2^{-Jp'(s-\frac{n}{p})} \mu_J^{p'-1}, \quad J \in \mathbb{N}_0, \tag{25}$$

where only the cases $s \leq \frac{n}{p}$ are of interest, otherwise (25) is always satisfied.

Proposition 3.3. *Let*

$$1 < p < \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1, \quad s > 0.$$

Let μ be a Radon measure in \mathbb{R}^n with (20) and (25). Let $a_k = a_k(\text{tr}_\Gamma)$ be the approximation numbers of the compact operator tr_Γ in (22). There are two positive numbers c and c' such that

$$a_{c2^{nJ}} \leq c' 2^{-J(s-\frac{n}{p})} \mu_J^{\frac{1}{p}}, \quad J \in \mathbb{N}_0, \quad (26)$$

where $c2^{nJ}$ is always assumed to be a natural number.

We prove the proposition in section 4.6.

3.2. Anisotropic d -sets in \mathbb{R}^n

We consider special measures μ and assume $\Gamma = \text{supp } \mu$ for some measure according to section 3.1, in particular with (18), now. Let again $a = (a_1, \dots, a_n)$ be a given anisotropy.

Definition 3.4. Let $0 < d < n$. Then $\Gamma \subset \mathbb{R}^n$ is called an anisotropic d -set if

$$\mu(B^a(\gamma, r)) \sim r^d, \quad 0 < r < 1, \quad (27)$$

where $B^a(\gamma, r) = \{y \in \mathbb{R}^n : |y - \gamma|_a \leq r\}$ and $\gamma \in \Gamma$.

In the following proposition we prove the existence of anisotropic d -sets.

Proposition 3.5. *For every $0 < d < n$ there exists an anisotropic d -set.*

Remark 3.6. One can show that our definition for the anisotropic d -set is a generalization of Farkas' definition in [8, 3.1].

3.3. Main assertion

We are now prepared to formulate our main result.

Theorem 3.7. *Let the anisotropic d -set Γ and μ be given according to (27), and*

$$0 < d < n, \quad 1 < p < \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1, \quad \frac{n}{p} \geq s > \frac{n-d}{p}.$$

Let $a_k = a_k(\text{tr}_\Gamma)$ be the approximation numbers of the compact operator tr_Γ according to (22). Then there exist numbers $c, c' > 0$ so that for all $k \in \mathbb{N}$

$$ck^{\frac{1}{d}(\frac{n}{p}-s)-\frac{1}{p}} \leq a_k(\text{tr}_\Gamma : B_p^{s,a}(\mathbb{R}^n) \hookrightarrow L_p(\Gamma)) \leq c' k^{\frac{1}{d}(\frac{n}{p}-s)-\frac{1}{p}}. \quad (28)$$

Remark 3.8. Let Γ be the anisotropic d -set considered in [8, 3.1], see Remark 3.6. Farkas proved in this situation that

$$e_k(\text{tr}_\Gamma : B_{p_1 q}^{\delta + \frac{n-d}{p_1}, a}(\mathbb{R}^n) \longrightarrow L_{p_2}(\Gamma)) \sim ck^{-\frac{\delta}{d}}$$

where $0 < p_1, p_2, q \leq \infty$, and $\delta > 0$. Let $p_1 = p_2 = q = p$ and $s = \delta + \frac{n-d}{p}$. Then

$$e_k(\text{tr}_\Gamma : B_p^{s, a}(\mathbb{R}^n) \longrightarrow L_p(\Gamma)) \sim ck^{\frac{1}{d}(\frac{n}{p} - s) - \frac{1}{p}}.$$

So we have the same results for the entropy and approximation numbers in the special case $p_1 = p_2$ which is not surprising, but cannot be expected for $p_1 \neq p_2$.

In view of the isotropic result [21, Theorem 2, Remark 9], if we restrict the outcome [21] to the classical example of a compact d -set with $0 < d < n$, then we have the same result like in the anisotropic setting.

4. Proofs

4.1. Proof of Theorem 2.5

Proof. We closely follow the proof in [18, 2.3.3] for the isotropic case.

Step 1. We prove that (10) is an equivalent quasi-norm in $B_{pq}^{s, a}$. It is sufficient to show that there exists a constant $c > 0$ such that

$$\|(\varphi_j^a \hat{f})^\vee | L_p\| \leq c2^{-j\sigma_p} \|(\varphi \hat{f})^\vee | L_p\|, \quad -j \in \mathbb{N}, \tag{29}$$

holds, because we need to prove that

$$\left(\sum_{j=-\infty}^{-1} 2^{jsq} \|(\varphi_j^a \hat{f})^\vee | L_p\|^q \right)^{1/q} \leq c \|(\varphi \hat{f})^\vee | L_p\|$$

and this is satisfied if (29) is true. For those j 's we have that $\varphi_j^a(x) = \varphi_j^a(x)\varphi(x)$ by the support condition (6) and (7) with $-j \in \mathbb{N}$, and hence

$$\begin{aligned} \|(\varphi_j^a \hat{f})^\vee | L_p\| &= \|(\varphi_j^a((\varphi \hat{f})^\vee)^\vee)^\vee | L_p\| \\ &\leq c \|\check{\varphi}_j^a | L_r\| \|(\varphi \hat{f})^\vee | L_p\|, \quad r = \min(1, p), \end{aligned} \tag{30}$$

where the inequality comes from the Fourier multiplier assertion for entire analytic functions, $\|F^{-1}MFf | L_p\| \leq \|F^{-1}M | L_{\tilde{p}}\| \|f | L_p\|$ where $\tilde{p} = \min(1, p)$, proved in [17, Proposition 1.5.1]. Elementary calculations show that $\check{\varphi}_j^a(x) = 2^{jn}\check{\varphi}_0(2^{ja}x)$ such that $\|\check{\varphi}_j^a | L_r\| = 2^{-j\frac{n}{r}} \|\check{\varphi}_0(2^{ja}\cdot) | L_r\| \leq c2^{-j\frac{n}{r} + ja}$ as $a_1 + \dots + a_n = n$. By (30) we thus have that

$$\|(\varphi_j^a \hat{f})^\vee | L_p\| \leq 2^{-jn(\frac{1}{r} - 1)} \|(\varphi \hat{f})^\vee | L_p\|$$

and we obtain (29) since $\sigma_p = n(\frac{1}{r} - 1)$.

Step 2. We prove that (11) is an equivalent quasi-norm in $B_{pq}^{s,a}$. By our assumption $s > \sigma_p$, we may assume that (9) converges not only in \mathcal{S}' , but also, say almost everywhere in \mathbb{R}^n . Then we have

$$\|f | L_p\| \leq c\|(\varphi\hat{f})^\vee | L_p\| + c\left(\sum_{j=1}^\infty \|(\varphi_j^a\hat{f})^\vee | L_p\|^p\right)^{1/p} \tag{31}$$

if $0 < p \leq 1$ and a corresponding estimate if $1 < p < \infty$. Now (10) and (31) prove that (11) can be estimated from above by $c\|f | B_{pq}^{s,a}\|$. We consider the converse inequality. Because f is a regular distribution we have a.e. that

$$(\varphi\hat{f})^\vee(x) = f(x) + ((1 - \varphi(\cdot))\hat{f})^\vee(x) = f(x) + \sum_{j=0}^\infty ((1 - \varphi(\cdot))\varphi_j^a(\cdot)\hat{f})^\vee(x).$$

By the above-mentioned Fourier multiplier assertion we have

$$\|(\varphi\hat{f})^\vee | L_p\| \leq c\|f | L_p\| + c\left(\sum_{j=0}^\infty \|(\varphi_j^a\hat{f})^\vee | L_p\|^p\right)^{1/p} \tag{32}$$

if $0 < p \leq 1$ and a corresponding estimate if $1 < p < \infty$. Now (10) and (32) prove that $\|f | B_{pq}^{s,a}\|$ can be estimated from above by the quasi-norm (11). \square

4.2. Proof of Proposition 2.7

Proof. We closely follow the proof in [4, Prop. 2.3.1] for the isotropic case. Let $\psi = \varphi_1$ be the same function as in (7). We have by (12)

$$\|f | L_p\| + \left(\int_0^\infty t^{-sq}\|(\psi(t\cdot)\hat{f})^\vee | L_p\|^q \frac{dt}{t}\right)^{1/q} \tag{33}$$

is an equivalent quasi-norm on $B_{pq}^{s,a}$. Elementary calculation shows that

$$\begin{aligned} (\psi(t\cdot)f(R\cdot)\widehat{(\cdot)})^\vee(x) &= (\psi(t\cdot)\hat{f}(R^{-1}\cdot))^\vee(x)R^{-n} \\ &= (\psi(t(R\cdot))\hat{f}(\cdot))^\vee(Rx). \end{aligned} \tag{34}$$

also in the anisotropic case, where $a_1 + \dots + a_n = n$. From (33), with $f(Rx)$ in place of $f(x)$, and (34) we obtain

$$\begin{aligned} \|f(R\cdot) | B_{pq}^{s,a}\| &\leq c_1\|f(R\cdot) | L_p\| + c_1\left(\int_0^\infty t^{-sq}\|\mathcal{F}^{-1}(\psi(t\cdot)\mathcal{F}[f(R\cdot)]) | L_p\|^q \frac{dt}{t}\right)^{1/q} \\ &\leq c_2R^{-\frac{n}{p}}\|f | L_p\| + c_3R^{s-\frac{n}{p}}\left(\int_0^\infty t^{-sq}\|\mathcal{F}^{-1}(\psi(t(R\cdot))\mathcal{F}f) | L_p\|^q \frac{dt}{t}\right)^{1/q} \end{aligned}$$

and from here follows (13) for $R \geq 1$, $c_1, c_2, c_3 > 0$ and $s > \sigma_p$. \square

4.3. Proof of Theorem 2.8

Proof. Step 1. At first we prove the left-hand side of (17). By (15) we have

$$f_j^a(2^{-(j+1)a}x) = \sum_{k \in \mathbb{Z}^n} c_k f(x - 2^a k), \quad c_k \in \mathbb{C}, \quad j \in \mathbb{N}, \tag{35}$$

where $f \in B_p^{s,a}$ and (14) is true. We would like to show that

$$\left\| \sum_{k \in \mathbb{Z}^n} c_k f(\cdot - 2^a k) \right\|_{B_p^{s,a}} \sim \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p \right)^{1/p} \|f\|_{B_p^{s,a}}. \tag{36}$$

We use the characterization of $B_p^{s,a}$ via local means; see [7, 4.4]. Recall notation (3). Let $k \in C^\infty$ so that $\text{supp } k \subset B^a = \{y \in \mathbb{R}^n : |y|_a \leq 1\}$ and

$$k(t, f)(x) = \int_{\mathbb{R}^n} k(y) f(x + t^a y) dy, \quad t > 0. \tag{37}$$

Let $k_0 \in C^\infty$ such that $\text{supp } k_0 \subset B^a$, and $s_1 > \max(s, \sigma_p) + \sigma_p$. Then

$$\|f\|_{B_p^{s,a}} \sim \|k_0(1, f)\|_{L_p} + \left(\sum_{j=1}^\infty 2^{j s p} \|k(2^{-j}, f)\|_{L_p}^p \right)^{1/p}$$

is an equivalent quasi-norm in $B_p^{s,a}$; see [7, 4.4]. We insert (35) in (37) and obtain

$$\begin{aligned} k\left(t, \sum_{m \in \mathbb{Z}^n} c_m f(\cdot - 2^a m)\right)(x) &= \int_{\mathbb{R}^n} k(y) \left(\sum_{m \in \mathbb{Z}^n} c_m f(x + t^a y - 2^a m) \right) dy \\ &= \sum_{m \in \mathbb{Z}^n} c_m \int_{\mathbb{R}^n} k(y) f(x - 2^a m + t^a y) m dy \\ &= \sum_{m \in \mathbb{Z}^n} c_m k(t, f)(x - 2^a m) \end{aligned}$$

and it follows

$$\begin{aligned} &\left\| \sum_{m \in \mathbb{Z}^n} c_m f(\cdot - 2^a m) \right\|_{B_p^{s,a}} \\ &\sim \left\| k_0\left(1, \sum_{m \in \mathbb{Z}^n} c_m f(\cdot - 2^a m)\right) \right\|_{L_p} \\ &\quad + \left(\sum_{j=1}^\infty 2^{j s p} \left\| k(2^{-j}, \sum_{m \in \mathbb{Z}^n} c_m f(\cdot - 2^a m)) \right\|_{L_p}^p \right)^{\frac{1}{p}} \\ &\sim \left(\sum_{m \in \mathbb{Z}^n} |c_m|^p \right)^{1/p} \left(\|k_0(1, f)\|_{L_p} + \left(\sum_{j=1}^\infty 2^{j s p} \|k(2^{-j}, f)\|_{L_p}^p \right)^{\frac{1}{p}} \right) \\ &\sim \left(\sum_{m \in \mathbb{Z}^n} |c_m|^p \right)^{1/p} \|f\|_{B_p^{s,a}}. \end{aligned}$$

Now the left-hand side of inequality of (17) is an easy consequence of Proposition 2.7, (35) and (36):

$$\begin{aligned} \|f_j^a | B_p^{s,a}\| &\leq c2^{j(s-n/p)} \left\| \sum_{m \in \mathbb{Z}^n} c_m f(\cdot - 2^a m) | B_p^{s,a} \right\| \\ &\leq c' 2^{j(s-n/p)} \left(\sum_{m \in \mathbb{Z}^n} |c_m|^p \right)^{1/p} \|f | B_p^{s,a}\|. \end{aligned}$$

Step 2. In this step we prove the right-hand side of (17). For this we would like to use the localization property given in [4, 2.3.2] if $n = 1$ and for the functions

$$f^{j\alpha}(x) = \sum_{m \in \mathbb{Z}} c_m f(2^{(j+1)\alpha}x - 2^\alpha m), \quad c_m \in \mathbb{C}, \quad j, \alpha \in \mathbb{N},$$

where $f \in \mathcal{S}'(\mathbb{R})$. By [4, 2.3.2/4] we know that there exist two constants $c' > 0$ and $c'' > 0$ such that for all $f \in B_{pp}^s$

$$c' \|f^{j\alpha} | B_{pp}^s\| \leq 2^{j\alpha(s-\frac{1}{p})} \left(\sum_{k \in \mathbb{Z}} |c_k|^p \right)^{1/p} \|f | B_{pp}^s\| \leq c'' \|f^{j\alpha} | B_{pp}^s\|, \quad (38)$$

as for $n = 1$ isotropic and anisotropic results coincide. For the functions f_j^a given in (15) we use the Fubini property of $B_p^{s,a}$; see [2, 6.], i.e.,

$$\begin{aligned} \|f_j^a | B_p^{s,a}(\mathbb{R}^n)\| &\sim \sum_{m=1}^n \left\| \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1}) \right\|_{x'}, \quad (39) \end{aligned}$$

where $x' = (x_1, \dots, x_{m-1}, x_{m+1}, \dots, x_n)$ and $s_m = \frac{s}{a_m}$. By (15) and (16)

$$\begin{aligned} &\|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} \\ &= \left\| \sum_{k_m=-\infty}^{\infty} f_m(2^{(j+1)a_m}x_m - 2^{a_m}k_m) \left[\sum_{\bar{k} \in \mathbb{Z}^{n-1}} c_{(k_1, \dots, k_n)} \bar{f} \right] | B_p^{s_m}(\mathbb{R}) \right\|_{x_m}, \end{aligned}$$

where $\bar{k} = (k_1, \dots, k_{m-1}, k_{m+1}, \dots, k_n)$ and

$$\begin{aligned} \bar{f} &= f_1(2^{(j+1)a_1}x_1 - 2^{a_1}k_1) \cdots f_{m-1}(2^{(j+1)a_{m-1}}x_{m-1} - 2^{a_{m-1}}k_{m-1}) \\ &\quad \times f_{m+1}(2^{(j+1)a_{m+1}}x_{m+1} - 2^{a_{m+1}}k_{m+1}) \cdots f_n(2^{(j+1)a_n}x_n - 2^{a_n}k_n). \end{aligned}$$

Let $d_{k_m} = \left(\sum_{l \in \mathbb{Z}^n, l_m=k_m} |c_l|^p \right)^{1/p}$ and without restriction of generality we may assume

that $d_{k_m} > 0$. We have that

$$\begin{aligned} & \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} \\ &= \left\| \sum_{k_m=-\infty}^{\infty} f_m(2^{(j+1)a_m} x_m - 2^{a_m} k_m) \left[\sum_{\bar{k} \in \mathbb{Z}^{n-1}} d_{k_m} \frac{c^{(k_1, \dots, k_n)}}{d_{k_m}} \bar{f} \right] | B_p^{s_m}(\mathbb{R}) \right\|_{x_m}. \end{aligned} \quad (40)$$

Let $\bar{c}_{\bar{k}} = \frac{c^{(k_1, \dots, k_n)}}{d_{k_m}}$ and by (40) we get that

$$\begin{aligned} & \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} \\ &= \left\| \sum_{k_m=-\infty}^{\infty} d_{k_m} f_m(2^{(j+1)a_m} x_m - 2^{a_m} k_m) \left[\sum_{\bar{k} \in \mathbb{Z}^{n-1}} \bar{c}_{\bar{k}} \bar{f} \right] | B_p^{s_m}(\mathbb{R}) \right\|_{x_m} \\ &= \left[\sum_{\bar{k} \in \mathbb{Z}^{n-1}} \bar{c}_{\bar{k}} \bar{f} \right] \left\| \sum_{k_m=-\infty}^{\infty} d_{k_m} f_m(2^{(j+1)a_m} x_m - 2^{a_m} k_m) | B_p^{s_m}(\mathbb{R}) \right\|_{x_m}. \end{aligned} \quad (41)$$

By (41),

$$\begin{aligned} & \left\| \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1}) \right\|_{x'} \\ &= \left\| \left[\sum_{\bar{k} \in \mathbb{Z}^{n-1}} \bar{c}_{\bar{k}} \bar{f} \right] \left\| \sum_{k_m=-\infty}^{\infty} d_{k_m} f_m(2^{(j+1)a_m} x_m - 2^{a_m} k_m) | B_p^{s_m}(\mathbb{R}) \right\|_{x_m} | L_p(\mathbb{R}^{n-1}) \right\|_{x'} \\ &= \left\| \sum_{k_m=-\infty}^{\infty} d_{k_m} f_m(2^{(j+1)a_m} x_m - 2^{a_m} k_m) | B_p^{s_m}(\mathbb{R}) \right\|_{x_m} \\ &\quad \times \left\| \sum_{\bar{k} \in \mathbb{Z}^{n-1}} \bar{c}_{\bar{k}} \bar{f} | L_p(\mathbb{R}^{n-1}) \right\|_{x'}. \end{aligned} \quad (42)$$

Note that

$$\begin{aligned} \left\| \sum_{\bar{k} \in \mathbb{Z}^{n-1}} \bar{c}_{\bar{k}} \bar{f} | L_p(\mathbb{R}^{n-1}) \right\|_{x'} &= \left(\sum_{\bar{k} \in \mathbb{Z}^{n-1}} |\bar{c}_{\bar{k}}|^p \right)^{1/p} 2^{(j+1)(a_m-n)/p} \\ &\quad \times \|f_1 \cdots f_{m-1} f_{m+1} \cdots f_n | L_p(\mathbb{R}^{n-1})\|_{x'}. \end{aligned} \quad (43)$$

(Recall that $-(a_1 + \dots + a_{m-1} + a_{m+1} + \dots + a_n) = a_m - n$.) Now we use (38) for the spaces $B_p^{s_m}(\mathbb{R})$ and by (42), (43)

$$\begin{aligned} & \left\| \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1}) \right\|_{x'} \\ &\geq c' 2^{ja_m(s_m - \frac{1}{p})} \left(\sum_{k_m=-\infty}^{\infty} |d_{k_m}|^p \right)^{1/p} \|f_m | B_p^{s_m}(\mathbb{R})\|_{x_m} \\ &\quad \times 2^{\frac{j}{p}(a_m-n)} \left(\sum_{\bar{k} \in \mathbb{Z}^{n-1}} |\bar{c}_{\bar{k}}|^p \right)^{\frac{1}{p}} \|f_1 \cdots f_{m-1} \cdot f_{m+1} \cdots f_n | L_p(\mathbb{R}^{n-1})\|_{x'}. \end{aligned} \quad (44)$$

On the other hand,

$$\left(\sum_{k_m=-\infty}^{\infty} |d_{k_m}|^p\right)^{1/p} = \left(\sum_{k_m=-\infty}^{\infty} \sum_{\substack{l \in \mathbb{Z}^n \\ l_m=k_m}} |c_l|^p\right)^{1/p} = \left(\sum_{l \in \mathbb{Z}^n} |c_l|^p\right)^{1/p} \tag{45}$$

and

$$\begin{aligned} \left(\sum_{\bar{k} \in \mathbb{Z}^{n-1}} |\bar{c}_{\bar{k}}|^p\right)^{1/p} &= \left(\sum_{\bar{k} \in \mathbb{Z}^{n-1}} \frac{|c_{(k_1, \dots, k_n)}|^p}{d_{k_m}^p}\right)^{1/p} \\ &= \frac{1}{d_{k_m}} \underbrace{\left(\sum_{\bar{k} \in \mathbb{Z}^{n-1}} |c_{(k_1, \dots, k_n)}|^p\right)^{1/p}}_{\geq d_{k_m}} \geq 1. \end{aligned} \tag{46}$$

By (44), (45) and (46) and $s_m \cdot a_m = s$, we conclude

$$\begin{aligned} &\| \|f_j^a(x_1, \dots, x_{m-1}, \cdot, x_{m+1}, \dots, x_n) | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1})\|_{x'} \\ &\geq c' 2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p\right)^{1/p} \|f_m | B_p^{s_m}(\mathbb{R})\|_{x_m} \\ &\quad \times \|f_1 \cdots f_{m-1} \cdot f_{m+1} \cdots f_n | L_p(\mathbb{R}^{n-1})\|_{x'} \\ &\geq c' 2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p\right)^{1/p} \| \|f_1 \cdots f_n | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1})\|_{x'}. \end{aligned} \tag{47}$$

By (47) and the Fubini property (39) we obtain the right-hand side of inequality of (17)

$$\begin{aligned} \|f_j^a | B_p^{s,a}\| &\geq c' 2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p\right)^{1/p} \sum_{m=1}^n \| \|f_1 \cdots f_n | B_p^{s_m}(\mathbb{R})\|_{x_m} | L_p(\mathbb{R}^{n-1})\|_{x'} \\ &\geq c' 2^{j(s-\frac{n}{p})} \left(\sum_{k \in \mathbb{Z}^n} |c_k|^p\right)^{1/p} \|f | B_p^{s,a}(\mathbb{R}^n)\|. \end{aligned} \quad \square$$

4.4. Wavelet frames

In the sequel we describe wavelet frames which are an effective instrument to estimate approximation numbers. This will be needed in our proofs below. Let k be a non-negative C^∞ function in \mathbb{R}^n with

$$\text{supp } k \subset \{y \in \mathbb{R}^n : |y|_a < 2^J, y_j > 0\}, \tag{48}$$

for some $J \in \mathbb{N}$, and

$$\sum_{m \in \mathbb{Z}^n} k(x - m) = 1, \quad x \in \mathbb{R}^n.$$

Recall that $x^\beta = x_1^{\beta_1} \cdots x_n^{\beta_n}$ where $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $\beta \in \mathbb{N}_0^n$, and put

$$k^\beta(x) = (2^{-J^a x})^\beta k(x) \geq 0, \quad x \in \mathbb{R}^n, \quad \beta \in \mathbb{N}_0^n.$$

Let

$$\omega \in \mathcal{S}, \quad \text{supp } \omega \subset (-\pi, \pi)^n, \quad \omega(x) = 1 \quad \text{if } |x|_a \leq 2,$$

and let

$$\omega^\beta(x) = \frac{i^{|\beta|} 2^{J^a \beta}}{(2\pi)^n \beta!} x^\beta \omega(x) \quad \text{for } x \in \mathbb{R}^n, \quad \beta \in \mathbb{N}_0^n,$$

and

$$\Omega^\beta(x) = \sum_{m \in \mathbb{Z}^n} (\omega^\beta)^\vee(m) e^{-imx}, \quad x \in \mathbb{R}^n, \quad \beta \in \mathbb{N}_0^n,$$

where $|\beta| = \beta_1 + \cdots + \beta_n$, and $\beta! = \beta_1! \cdots \beta_n!$ and $a\beta = a_1\beta_1 + \cdots + a_n\beta_n$. Let φ_0 be a C^∞ function in \mathbb{R}^n with

$$\varphi_0(x) = 1 \quad \text{if } |x|_a \leq 1 \quad \text{and} \quad \varphi_0(x) = 0 \quad \text{if } |x|_a \geq \frac{3}{2},$$

and let $\varphi(x) = \varphi_0(x) - \varphi_0(2^a x)$. Then

$$\Phi_{jm}^\beta(x) = \begin{cases} \Phi_F^\beta(x - m), & \text{if } j = 0, \\ \Phi_M^\beta(2^{ja}x - m), & \text{if } j \in \mathbb{N}, \end{cases}$$

are analytic wavelets where the *father wavelets* $\Phi_F^\beta(x)$ and the *mother wavelets* $\Phi_M^\beta(x)$ are given by their inverse Fourier transforms

$$\begin{aligned} (\Phi_F^\beta)^\vee(\xi) &= \varphi_0(\xi) \Omega^\beta(\xi), \quad \xi \in \mathbb{R}^n, \\ (\Phi_M^\beta)^\vee(\xi) &= \varphi(\xi) \Omega^\beta(\xi), \quad \xi \in \mathbb{R}^n. \end{aligned}$$

For the sequence

$$\lambda = \{ \lambda_{jm}^\beta \in \mathbb{C} : j \in \mathbb{N}_0, \quad m \in \mathbb{Z}^n, \quad \beta \in \mathbb{N}_0^n \},$$

$s \in \mathbb{R}$, $0 < p \leq \infty$, and $\varrho \geq 0$, we put

$$\|\lambda\|_{b_p^{s,\varrho}} = \left(\sum_{\beta \in \mathbb{N}_0^n} \sum_{j=0}^{\infty} \sum_{m \in \mathbb{Z}^n} 2^{\varrho a \beta p + j(s-n/p)p} |\lambda_{jm}^\beta|^p \right)^{1/p}. \tag{49}$$

For $f \in L_p(\mathbb{R}^n)$,

$$\lambda_{jm}^\beta(f) = 2^{jn} \int_{\mathbb{R}^n} f(x) \Phi_{jm}^\beta(x) dx, \quad j \in \mathbb{N}_0, \quad m \in \mathbb{Z}^n, \quad \beta \in \mathbb{N}_0^n.$$

We make use of the following wavelet characterization.

Theorem 4.1 ([9, Theorem 2.1]). *Let $0 < p < \infty$, $s > \sigma_p$ where $\sigma_p = n(\frac{1}{p} - 1)_+$, $\varrho \geq 0$, and a an anisotropy. Then $f \in \mathcal{S}'$ is an element of $B_p^{s,a}$ if, and only if, it can be represented as*

$$f = \sum_{\beta \in \mathbb{N}_0^n} \sum_{j=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta k^\beta(2^{ja}x - m), \quad x \in \mathbb{R}^n, \tag{50}$$

with $\|\lambda\|_{b_p^{s,\varrho}} < \infty$, absolute convergence being in $L_{\max(1,p)}$. Furthermore,

$$\|f\|_{B_p^{s,a}} \sim \inf \|\lambda\|_{b_p^{s,\varrho}},$$

where the infimum is taken over all admissible representations (50). In addition, any $f \in B_p^{s,a}$ can be optimally represented by

$$f = \sum_{\beta \in \mathbb{N}_0^n} \sum_{j=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k^\beta(2^{ja}x - m), \tag{51}$$

with

$$\|f\|_{B_p^{s,a}} \sim \|\lambda(f)\|_{b_p^{s,\varrho}}. \tag{52}$$

Remark 4.2. In the sequel we shall stick to the notation

$$k_{jm}^\beta(x) = k^\beta(2^{ja}x - m), \quad \beta \in \mathbb{N}_0^n, \quad j \in \mathbb{N}_0, \quad m \in \mathbb{Z}^n. \tag{53}$$

4.5. Proof of Proposition 3.1

We closely follow the ideas in [21].

Proof. Step 1. We prove the existence of (22) like in the isotropic case, see [20, Theorem 9.3, Corollary 9.8; 21, Prop. 2]. In our case we use the anisotropic local means and the equivalent norm in anisotropic function spaces, see [8, 2.2]. In comparison with [20, Theorem 9.3] we need only a special case where $u = v = p'$ and $\sigma = -s$. On the other hand, the existence of tr_Γ can also be shown by similar arguments as presented below.

Step 2. Let $f \in B_p^{s,a}$ be given by (51), (52). (We use the notation (53).) For any fixed $\beta \in \mathbb{N}_0^n$ we have

$$\begin{aligned} \left\| \sum_{j=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta \Big| L_p(\Gamma) \right\| &\leq \sum_{j=0}^{\infty} \left\| \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta \Big| L_p(\Gamma) \right\| \\ &\leq c \sum_{j=0}^{\infty} \left(\int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} |\lambda_{jm}^\beta(f)|^p |k_{jm}^\beta(x)|^p \mu(dx) \right)^{1/p} \end{aligned}$$

$$\begin{aligned} &\leq c \sum_{j=0}^{\infty} \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{jm}^\beta(f)|^p \int_{cQ_{jm}^a} |k_{jm}^\beta(x)|^p \mu(dx) \right)^{1/p} \\ &\leq c' \sum_{j=0}^{\infty} \mu_j^{\frac{1}{p}} \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{jm}^\beta(f)|^p \right)^{1/p} \end{aligned}$$

where we used the boundedness of k and (19). We apply the Hölder inequality, recall that $\frac{1}{p} + \frac{1}{p'} = 1$, and so we can continue

$$\begin{aligned} &\left\| \sum_{j=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta \mid L_p(\Gamma) \right\| \\ &\leq c' \left(\sum_{j=0}^{\infty} 2^{-jp'(s-\frac{n}{p})} \mu_j^{\frac{p'}{p}} \right)^{1/p'} \left(\sum_{j,m} 2^{j(s-\frac{n}{p})p} |\lambda_{jm}^\beta(f)|^p \right)^{1/p}. \end{aligned} \tag{54}$$

We choose $\varrho > 0$. Then it follows by (49) and (52) that

$$\| \text{tr}_\Gamma f \mid L_p(\Gamma) \| \leq c' \left(\sum_{j=0}^{\infty} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} \right)^{1/p'} \| f \mid B_p^{s,a} \| \tag{55}$$

where c' is independent of μ . This proves (23).

Step 3. We prove that tr_Γ is compact. Let $B \in \mathbb{N}$, $J \in \mathbb{N}$, $[a\beta] = \max\{r \in \mathbb{Z} : r \leq a\beta\}$, and let $\text{tr}_\Gamma^{B,J}$ be given by

$$\text{tr}_\Gamma^{B,J} f = \sum_{[a\beta] \leq B} \sum_{j \leq J} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta, \tag{56}$$

where again $f \in B_p^{s,a}$ is given by (51),(52) and where the sum $\sum_{m \in \mathbb{Z}^n}^\Gamma$ is restricted to those $m \in \mathbb{Z}^n$ such that the rectangles Q_{jm}^a have a non-empty intersection with Γ . For given $\delta > 0$ and suitably chosen $\varrho > 0$ it follows by the above arguments for $f \in B_p^{s,a}$ having norm of at most 1 that

$$\begin{aligned} &\| (\text{tr}_\Gamma - \text{tr}_\Gamma^{B,J}) f \mid L_p(\Gamma) \| \\ &\leq c \left(\sum_{[a\beta] \geq B} 2^{-\delta a\beta} \right) + c \left(\sum_{[a\beta] \leq B} 2^{-\delta a\beta} \right) \left(\sum_{j \geq J} 2^{-jp'(s-\frac{n}{p})} \mu_j^{p'-1} \right)^{1/p'}, \end{aligned} \tag{57}$$

see (49) and (54), (55). By (21) we find for any given $\varepsilon > 0$ sufficiently large numbers B and J such that

$$\| \text{tr}_\Gamma - \text{tr}_\Gamma^{B,J} \| \leq \varepsilon.$$

Then tr_Γ is compact, as $\text{tr}_\Gamma^{B,J}$ are finite rank operators. □

4.6. Proof of Proposition 3.3

Proof. Note that (25) implies (21), thus by Proposition 3.1 the operator tr_Γ is compact. We refine (56) by

$$\text{tr}_\Gamma^J f = \sum_{[a,\beta] \leq J} \sum_{j \leq J - [a,\beta]} \sum_{m \in \mathbb{Z}^n}^\Gamma \lambda_{jm}^\beta(f) k_{jm}^\beta, \quad J \in \mathbb{N}, \tag{58}$$

where again $f \in B_p^{s,a}$ is given by (51), (52) and the last sum has the same meaning as the last sum in (56). As μ is a measure in \mathbb{R}^n we have that

$$\mu_K \leq c2^{(J-K)n} \mu_J, \quad K \leq J, \tag{59}$$

also in the anisotropic case. (Recall that $a_1 + \dots + a_n = n$.) Let $\delta > 0$ be sufficiently large. By (59) we obtain for $f \in B_p^{s,a}$ having norm of at most 1 in analogy to (57) that

$$\begin{aligned} \|(\text{tr}_\Gamma - \text{tr}_\Gamma^J)f | L_p(\Gamma)\| &\leq c2^{-\delta J} + c \sum_{[a,\beta] \leq J} 2^{-\delta a\beta} \left(\sum_{j \geq J - [a,\beta]} 2^{-jp'(s - \frac{n}{p})} \mu_j^{\frac{p'}{p}} \right)^{1/p'} \\ &\leq c2^{-\delta J} + c \sum_{[a,\beta] \leq J} 2^{-\delta a\beta} 2^{-(J - [a,\beta])(s - \frac{n}{p})} \mu_{J - [a,\beta]}^{\frac{1}{p}} \\ &\leq c2^{-\delta J} + c\mu_J^{\frac{1}{p}} 2^{-J(s - \frac{n}{p})} \sum_{[a,\beta] \leq J} 2^{-\delta a\beta + a\beta(s - \frac{n}{p}) + a\beta \frac{n}{p}} \\ &\leq c'2^{-J(s - \frac{n}{p})} \mu_J^{\frac{1}{p}}. \end{aligned} \tag{60}$$

In the second estimate we used assumption (25) and in the next one (59). For the rank of tr_Γ^J we have the estimate

$$\text{rank}(\text{tr}_\Gamma^J) \leq c \sum_{[a,\beta] \leq J} 2^{n(J - [a,\beta])} \leq c'2^{nJ}.$$

This proves (26). □

4.7. Proof of Proposition 3.5

Proof. For simplicity we prove this proposition for the case $n = 2$. If $n > 2$ this can be done in a similar way.

We use the well-known mass distribution procedure to construct a measure μ with the desired properties. We refer to [19, ch. 4] for details. Let $Q = [0, 1]^2$ be the closed cube with side-length 1, we take the affine contractions $(A_m)_{m=1}^N$ on \mathbb{R}^2 which map the unit square to the rectangles $(A_m Q)_{m=1}^N$ with side-lengths r^{a_1} and r^{a_2} where $0 < a_1 < a_2$ and $a_1 + a_2 = 2$ as in figure 1, so that they are disjoint and




Figure 1

$\mu(A_m Q) = N^{-1}$. Furthermore we have $Nr^2 = N|A_m Q| < 1$. Let

$$AQ = (AQ)^1 = \bigcup_{m=1}^N A_m Q, \quad (AQ)^0 = Q,$$

$$(AQ)^k = A((AQ)^{k-1}).$$

The sequence of sets is monotonically decreasing and by [6, Theorem 8.3]

$$\Gamma = (AQ)^\infty = \bigcap_{k \in \mathbb{N}} (AQ)^k = \lim_{k \rightarrow \infty} (AQ)^k$$

is the uniquely determined fractal generated by the contractions $(A_m)_{m=1}^N$. But on the other hand we assume that $\mu(A_m Q) = r^d$, where $m = 1, \dots, N$, and from here we get that $d = \frac{\log N}{|\log r|}$. If $0 < d < 2$ then it follows by elementary geometrical reasoning that one can find (sufficiently small) numbers $r > 0$ and suitably chosen natural numbers $N \in \mathbb{N}$ with the desired properties. \square

4.8. Proof of Theorem 3.7

Proof. Step 1. First we prove the right-hand side of the estimate (28) in Theorem 3.7. Again we use the wavelet expansion (51), (52). For fixed $\beta \in \mathbb{N}_0^n$ we put

$$\text{tr}_\Gamma^\beta f = \sum_{j \in \mathbb{N}_0} \sum_{m \in \mathbb{Z}^n}^\Gamma \lambda_{jm}^\beta(f) k_{jm}^\beta$$

and

$$\text{tr}_\Gamma^{\beta,J} f = \sum_{j \leq J} \sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta, \tag{61}$$

where the second sum has the same meaning as the last sum in (56). By the same reasoning as in (58) and (60) but now for fixed β we have for $f \in B_p^{s,a}$ with norm of at most 1,

$$\|(\text{tr}_\Gamma^\beta - \text{tr}_\Gamma^{\beta,J})f | L_p(\Gamma)\| \leq c2^{-\delta a\beta} 2^{J(\frac{n}{p}-s)} \mu_J^{\frac{1}{p}}.$$

By Definition 3.4 there exists a constant $c > 0$ independent of $j \in \mathbb{N}_0$ with $\mu(Q_{jm}^a \cap \Gamma) \leq c2^{-jd}$ and we obtain that

$$\|(\text{tr}_\Gamma^\beta - \text{tr}_\Gamma^{\beta,J})f | L_p(\Gamma)\| \leq c2^{-\delta a\beta} 2^{J(\frac{n}{p}-s)} 2^{-J\frac{d}{p}}.$$

In definition (24) put $L = \text{tr}_\Gamma^{\beta,J}$, $T = \text{tr}_\Gamma^\beta$, and note that for $j \in \mathbb{N}_0$,

$$\text{rank}\left(\sum_{m \in \mathbb{Z}^n} \lambda_{jm}^\beta(f) k_{jm}^\beta\right) \leq c2^{jd}. \tag{62}$$

Thus we obtain by (61) that

$$\text{rank}(\text{tr}_\Gamma^{\beta,J}) \leq c \sum_{j \leq J} 2^{jd} \leq c' 2^{Jd}.$$

Then (62) implies that there are two positive numbers c and c' such that

$$a_{c2^{Jd}}(\text{tr}_\Gamma^\beta) \leq c' 2^{-\delta a\beta} 2^{J(\frac{n}{p}-s)} 2^{-J\frac{d}{p}}. \tag{63}$$

For $k \in \mathbb{N}$ there are numbers $J_k \in \mathbb{N}$ such that

$$2^{J_k d} \sim k \quad \text{with} \quad J_1 \leq J_2 \leq \dots \leq J_n \leq \dots; \tag{64}$$

inserted in (63) this leads to

$$a_{ck}(\text{tr}_\Gamma^\beta) \leq c2^{-\delta a\beta} 2^{J_k(\frac{n}{p}-s)} k^{-\frac{1}{p}}. \tag{65}$$

Let $\varepsilon > 0$, for given $k \in \mathbb{N}$ we apply (65) to $k_\beta \in \mathbb{N}$ with $k_\beta \sim 2^{-\varepsilon a\beta} k$. Then it follows by the additivity property of approximation numbers and from (65) that

$$\begin{aligned} a_{ck}(\text{tr}_\Gamma^\beta) &\leq \sum_{\beta \in \mathbb{N}_0^n} a_{k_\beta}(\text{tr}_\Gamma^\beta) \\ &\leq c' \sum_{\beta \in \mathbb{N}_0^n} 2^{-\delta a\beta} 2^{J_{k_\beta}(\frac{n}{p}-s)} (2^{-\varepsilon a\beta} k)^{-\frac{1}{p}} \\ &\leq c'' 2^{J_k(\frac{n}{p}-s)} k^{-\frac{1}{p}} \sum_{\beta \in \mathbb{N}_0^n} 2^{-a\beta(\delta - \frac{\varepsilon}{p})} \\ &\leq c''' 2^{J_k(\frac{n}{p}-s)} k^{-\frac{1}{p}} \end{aligned}$$

for $\varepsilon > 0$ small. We used $s \leq \frac{n}{p}$, such that $J_{k\beta}(\frac{n}{p} - s) \leq J_k(\frac{n}{p} - s)$. Finally (64) implies

$$a_{ck}(\text{tr}_\Gamma^\beta) \leq c''' k^{\frac{1}{d}(\frac{n}{p} - s) - \frac{1}{p}}$$

and so we finished the proof of the right-hand side of the estimate (28).

Step 2. To verify the left-hand side of the estimate (28) we closely follow the argument in [21, 4.4] for the isotropic case. Let $J \in \mathbb{N}$ and $c > 0$ be suitably chosen numbers such that there are lattice points

$$\gamma_{j,l} = 2^{(-j-J)a} m \quad \text{with} \quad m \in \mathbb{Z}^n, \quad l = 1, \dots, M_j \quad \text{where} \quad M_j \sim 2^{jd}$$

with

$$\text{dist}(\gamma_{j,l}, \Gamma) \leq c2^{-j} \quad \text{and disjoint anisotropic balls} \quad B^a(\gamma_{j,l}, c2^{-j+1}).$$

With k as in (48) we put for $j \in \mathbb{N}_0$,

$$f_j^a(x) = \sum_{l=1}^{M_j} c_{jl} 2^{-j(s-\frac{n}{p})} k(2^{ja}(x - \gamma_{j,l})), \quad c_{jl} \in \mathbb{C}, \quad x \in \mathbb{R}^n. \tag{66}$$

Then we obtain by Theorem 2.8

$$\|f_j^a|_{B_p^{s,a}}\| \sim 2^{j(s-\frac{n}{p})} \left(\sum_{l=1}^{M_j} 2^{-j(s-\frac{n}{p})} |c_{jl}|^p \right)^{\frac{1}{p}} = \left(\sum_{l=1}^{M_j} |c_{jl}|^p \right)^{\frac{1}{p}}$$

and

$$\begin{aligned} \|f_j^a|_{L_p(\Gamma)}\| &= \left(\int_\Gamma |f_j^a(x)|^p \mu(dx) \right)^{1/p} \\ &\sim 2^{-j(s-\frac{n}{p})} \left(\sum_{l=1}^{M_j} |c_{jl}|^p \int_\Gamma k^p(2^{ja}(x - \gamma_{j,l})) \mu(dx) \right)^{1/p} \\ &\sim 2^{-j(s-\frac{n}{p})} \left(\sum_{l=1}^{M_j} |c_{jl}|^p \int_{\Gamma \cap B^a(\gamma_{j,l}, c2^{-j})} k^p(2^{ja}(x - \gamma_{j,l})) \mu(dx) \right)^{1/p} \\ &\geq c2^{-j(s-\frac{n}{p})} 2^{-j\frac{d}{p}} \left(\sum_{l=1}^{M_j} |c_{jl}|^p \right)^{1/p} \end{aligned} \tag{67}$$

using our assumption (27) in the last estimate. Hence

$$\|f_j^a|_{L_p(\Gamma)}\| \geq c2^{-j(s-\frac{n}{p})} 2^{-j\frac{d}{p}} \quad \text{if} \quad \|f_j^a|_{B_p^{s,a}}\| \sim 1. \tag{68}$$

Now let T be an arbitrary linear operator,

$$T : B_p^{s,a} \hookrightarrow L_p(\Gamma) \quad \text{with} \quad \text{rank} T \leq M_j - 1. \tag{69}$$

Then we can find a function f_j^a according to (66) with norm 1 in $B_p^{s,a}$ and $Tf_j^a = 0$. Consequently, by (67) and (68),

$$\begin{aligned} \|\mathrm{tr}_\Gamma - T\| &= \sup\{ \|(\mathrm{tr}_\Gamma - T)f\|_{L_p(\Gamma)} : \|f\|_{B_p^{s,a}} \sim 1\} \\ &\geq \|(\mathrm{tr}_\Gamma - T)f_j^a\|_{L_p(\Gamma)} \\ &= \|f_j^a\|_{L_p(\Gamma)} \\ &\geq c2^{-j(s-\frac{n}{p})-j\frac{d}{p}}. \end{aligned}$$

As this is true for all T according to (69), we obtain

$$\begin{aligned} a_{M_j}(\mathrm{tr}_\Gamma) &= \inf\{ \|\mathrm{tr}_\Gamma - T\| : \mathrm{rank} T \leq M_{j-1} \} \\ &\geq c2^{-j(s-\frac{n}{p})-j\frac{d}{p}}. \end{aligned}$$

For $k \in \mathbb{N}$ there are numbers $j_k \in \mathbb{N}$ such that

$$2^{j_k d} \sim k \quad \text{with} \quad j_{k_1} \leq j_{k_2} \leq \dots \leq j_{k_n} \leq \dots,$$

inserted in (68) we obtain

$$a_k(\mathrm{tr}_\Gamma) \geq c2^{j_k(s-\frac{n}{p})} k^{-\frac{1}{p}} \geq c' k^{\frac{1}{d}(\frac{n}{p}-s)-\frac{1}{p}},$$

i.e., the left-hand side of the estimate (28). \square

Acknowledgements. I wish to thank Professor Hans Triebel and Dr. Dorothee D. Haroske for many inspiring discussions.

References

- [1] O. V. Besov, V. P. Il'in, and S. M. Nikol'skiĭ, *Integralnye predstavleniya funktsii i teoremy vlozheniya*, Izdat. "Nauka", Moscow, 1975.
- [2] S. Dachkovski, *Anisotropic function spaces and related semi-linear hypoelliptic equations*, Math. Nachr. **248/249** (2003), 40–61.
- [3] P. Dintelman, *On Fourier multipliers between anisotropic weighted function spaces*, Ph.D. thesis, TH Darmstadt, 1995.
- [4] D. E. Edmunds and H. Triebel, *Function spaces, entropy numbers, differential operators*, Cambridge Tracts in Mathematics, vol. 120, Cambridge University Press, Cambridge, 1996.
- [5] D. E. Edmunds and W. D. Evans, *Spectral theory and differential operators*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1987.
- [6] K. J. Falconer, *The geometry of fractal sets*, Cambridge Tracts in Mathematics, vol. 85, Cambridge University Press, Cambridge, 1986.
- [7] W. Farkas, *Atomic and subatomic decompositions in anisotropic function spaces*, Math. Nachr. **209** (2000), 83–113.

- [8] ———, *Eigenvalue distribution of some fractal semi-elliptic differential operators*, Math. Z. **236** (2001), no. 2, 291–320.
- [9] D. D. Haroske and E. Tamási, *Wavelet frames for distributions in anisotropic Besov spaces*, Georgian Math. J. **12** (2005), no. 4, 637–658.
- [10] J. Johnsen, *Pointwise multiplication of Besov and Triebel-Lizorkin spaces*, Math. Nachr. **175** (1995), 85–133.
- [11] S. M. Nikol'skiĭ, *Priblizhenie funktsii mnogikh peremennykh i teoremy vlozheniya*, “Nauka”, Moscow, 1977.
- [12] T. Runst and W. Sickel, *Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations*, de Gruyter Series in Nonlinear Analysis and Applications, vol. 3, Walter de Gruyter & Co., Berlin, 1996.
- [13] H.-J. Schmeisser and H. Triebel, *Topics in Fourier analysis and function spaces*, A Wiley-Interscience Publication, John Wiley & Sons Ltd., Chichester, 1987.
- [14] E. M. Stein and S. Wainger, *Problems in harmonic analysis related to curvature*, Bull. Amer. Math. Soc. **84** (1978), no. 6, 1239–1295.
- [15] B. Stöckert and H. Triebel, *Decomposition methods for function spaces of $B_{p,q}^s$ type and $F_{p,q}^s$ type*, Math. Nachr. **89** (1979), 247–267.
- [16] H. Triebel, *Fourier analysis and function spaces (selected topics)*, Teubner Verlagsgesellschaft, Leipzig, 1977.
- [17] ———, *Theory of function spaces*, Monographs in Mathematics, vol. 78, Birkhäuser Verlag, Basel, 1983.
- [18] ———, *Theory of function spaces. II*, Monographs in Mathematics, vol. 84, Birkhäuser Verlag, Basel, 1992.
- [19] ———, *Fractals and spectra*, Monographs in Mathematics, vol. 91, Birkhäuser Verlag, Basel, 1997.
- [20] ———, *The structure of functions*, Monographs in Mathematics, vol. 97, Birkhäuser Verlag, Basel, 2001.
- [21] ———, *Approximation numbers in function spaces and the distribution of eigenvalues of some fractal elliptic operators*, J. Approx. Theory **129** (2004), no. 1, 1–27.
- [22] M. Yamazaki, *A quasihomogeneous version of paradifferential operators. I. Boundedness on spaces of Besov type*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **33** (1986), no. 1, 131–174.