

About the Existence of Integrable Solutions of a Functional-Integral Equation ⁽¹⁾

G. EMMANUELE

ABSTRACT. We improve (in some sense) a recent theorem due to Banas and Knap ([2]) about the existence of integrable solutions of a functional-integral equation.

1. INTRODUCTION

Let $I = [0, 1]$ be. We consider the following functional-integral equation

$$x(t) = g(t) + f\left(t, \int_0^1 k(t, s)x(\varphi(s))ds\right) \quad t \in I \quad (1)$$

where $f: I \times \mathbb{R} \rightarrow \mathbb{R}^+ = [0, +\infty)$, $k: I \times I \rightarrow \mathbb{R}^+$, $g: I \rightarrow \mathbb{R}$ $\varphi: I \rightarrow I$ are functions verifying special hypotheses (see section 2) and we look-for solutions $x \in L^1(I)$. As remarked in the paper [2] this equation has been considered by a number of authors because of its importance in problems in physics, engineering and economics; further, problems in the theory of partial differential equations lead, sometimes, to the study of the equation (1). Recently, Banas and Knap ([2]) gave a result of existence of integrable solutions to (1). They were forced by the techniques used to consider certain monotonicity assumptions on g , f , k (see hypotheses i), ii) and iv) in [2]), that we are able to eliminate completely here. However, we must observe that Banas and Knap obtain a monotone solution, a fact that doesn't follow from our hypotheses. Prof Banas also observed that under our hypotheses we don't need to use the

⁽¹⁾ Work performed under the auspices of G.N.A.F.A. of C.N.R. and partially supported by M.U.R.S.T. of Italy (60%).

measure of weak noncompactness he considered in [2] because the operator we define following [2] actually has a relatively weakly compact range. So it is enough to apply Tychonoff fixed point Theorem ([5]). We take this opportunity to thank him very much for this remark that made our proof simpler.

2. PRELIMINARIES AND MAIN RESULT

As in the paper [2] we define the following four operators

$$(Kx)(t) = \int_0^1 k(t, s)x(s) ds$$

$$(Fx)(t) = f(t, x(t))$$

$$(Hx)(t) = f\left(t, \int_0^1 k(t, s)x(s) ds\right)$$

$$x = Ax = g + Hx(\varphi) = g + FKx(\varphi).$$

We consider the following hypotheses

- (i) $g \in L^1(I)$.
- (ii) $f: I \times \mathbb{R} \rightarrow \mathbb{R}^1$ satisfies Caratheodory hypotheses (i.e. f is measurable with respect to $t \in I$, for all $x \in \mathbb{R}$, and continuous in $x \in \mathbb{R}$, for a.a. $t \in I$) and there are $a \in L^1(I)$, $b \geq 0$ such that

$$f(t, x) \leq a(t) + b|x| \quad t \in I, \quad x \in \mathbb{R}$$

(this last inequality is a necessary and sufficient condition for F , and so H , to take values in $L^1(I)$ when acting on elements of $L^1(I)$; see Theorem 1 in [2])

- (iii) k verifies Caratheodory hypotheses and there is $\lambda \in L^1(I)$ such that

$$k(t, x) \leq \lambda(t) \quad t \text{ a.e. in } I, \quad x \in \mathbb{R}$$

(under (iii) the linear operator K maps $L^1(I)$ into $L^1(I)$ continuously; let us denote by $\|K\|$ its operator norm)

- (iv) $\varphi: I \rightarrow I$ is absolutely continuous and there exists $B > 0$ such that $\varphi'(t) \geq B$ for a.a. $t \in I$.

- (v) $b\|K\|/B < 1$.

The technique used in [2] is the following: under the above assumptions A is a weakly continuous operator from a suitable B_s into itself; furthermore there exists $L \in [0, 1]$ such that $\beta(A(Y)) \leq L\beta(Y)$, (β the measure of weak noncompactness introduced in [3]), for all nonempty subsets Y of B_s and hence results from [1] and [6] can be applied to get a fixed point of the operator $x \rightarrow g + FKx(\varphi)$. The difference between the result in [2] and our Theorem below resides in the technique we use to obtain the weak continuity of A ; indeed, Banas and Knap consider some monotonicity hypotheses on g, f, k we are able to dispense with. Further, we do not make use of the measure of weak noncompactness introduced in [3] as remarked in the Introduction.

Theorem. *Under the assumptions i)-v) above the equation [1] has at least a solution $x \in L^1(I)$.*

Proof. As in the paper [2] we can prove that $A: B_s \rightarrow B_s$, where $s = (||g|| + ||a||) / (1 - b||K||B^{-1})$. Furthermore, it is not difficult to see that the set $A(B_s)$ is relatively weakly compact ([5]), since it is bounded and uniformly integrable. Hence Tychonoff fixed point Theorem ([5]) will conclude the proof once we have the weak continuity of A . So, we need only to show that A is weakly continuous from B_s into B_s , i.e. A maps weakly convergent nets $(x_\alpha) \subset B_s$ into weakly convergent nets $(A(x_\alpha))$. It is clearly enough to show that H is weakly continuous. So let $(x_\alpha), x_0 \subset B_s$ be with $x_\alpha \rightharpoonup x_0$; if we prove that for any $\epsilon > 0$, any $y^* \in L^\infty(I), ||y^*|| \leq 1$ and any subnet (x_{α_β}) of (x_α) , there is another subnet $(x_{\alpha_{\beta_\gamma}})$ for which $|\langle H(x_{\alpha_{\beta_\gamma}}) - H(x_0), y^* \rangle| < \epsilon$ we are done (proceeding by contradiction, of course).

To reach our target, we start by noting that the operator $x \rightarrow x(\varphi)$ from $L^1(I)$ into itself is bounded and linear; hence it is weakly continuous and so $x_\alpha(\varphi) \rightharpoonup x_0(\varphi)$ in $L^1(I)$. Since B_s is bounded in $L^1(I)$, the set $\{x_\alpha(\varphi), x_0(\varphi)\}$ is even bounded in $L^1(I)$, by a number M . Now, given $\epsilon > 0$ choose

$\delta > 0$ such that $\text{meas}(\underline{D}) < \delta$, implies $\int_{\underline{D}} 2[a(t) + b\lambda(t)] dt < \frac{\epsilon}{2}$. Furthermore, choose a closed subset $I_1 \subset I$, $\text{meas}(I \setminus I_1) < \frac{\delta}{4}$, with $\lambda|_{I_1}$ continuous (use Lusin Theorem, [4]) $Q = \max_{I_1} \lambda$. Again consider a closed subset $I_2 \subset I$, $\text{meas}(I \setminus I_2) < \frac{\delta}{4}$, with $f|_{I_2 \times [-QM, QM]}$ continuous (and so uniformly continuous) and a closed subset $I_3 \subset I$, $\text{meas}(I \setminus I_3) < \frac{\delta}{4}$, with $k|_{I_3 \times I}$ continuous (and so uniformly continuous) (use Scorza-Dragnoni Theorem, [6]). Put $I_0 = \bigcap_{i=1}^3 I_i$. I_0 is a closed subset of I . Now, observe that, for $t', t'' \in I_0$, if $\psi_\alpha(t) = \int_0^1 k(t, s)x_\alpha(\varphi(s)) ds$, $\psi_0(t) = \int_0^1 k(t, s)x_0(\varphi(s)) ds$, one has

$$|\psi_\alpha(t') - \psi_\alpha(t'')| \leq \int_0^1 |k(t', s) - k(t'', s)| |x_\alpha(\varphi(s))| ds$$

(the same is true for ψ_0). Since $k|_{I_3 \times I_1}$ is uniformly continuous and $(x_\alpha) \subset B_s$, the set $\{\psi_\alpha, \psi_0\}$ is equicontinuous in $C^0(I_0)$. It is very easy to see that the same set is bounded by QM in the norm of $C^0(I_0)$, hence the Ascoli-Arzelà Theorem can be applied to get a relatively compact subset of $C^0(I_0)$. The net (ψ_{α_β}) admits a converging subnet $(\psi_{\alpha_{\beta_\gamma}})$. On the other hand, for $\bar{t} \in I_0$,

$$\psi_\alpha(\bar{t}) = \int_0^1 k(\bar{t}, s) x_\alpha(\varphi(s)) ds \rightarrow \psi_0(\bar{t}) = \int_0^1 k(\bar{t}, s) x_0(\varphi(s)) ds$$

since $x_\alpha(\varphi) \xrightarrow{w} x_0(\varphi)$ in $L^1(I)$ and $s \rightarrow k(\bar{t}, s)$ is in $L^\infty(I)$. Hence $\psi_{\alpha_{\beta_\gamma}} \rightarrow \psi_0$ in the C^0 -norm on I_0 . Now, recall that $f|_{I_0 \times [-QM, QM]}$ is uniformly continuous and so we have

$$\lim_{\gamma} f(t, \psi_{\alpha_{\beta_\gamma}}(t)) = f(t, \psi_0(t)) \quad \text{uniformly on } I_0 \quad (2)$$

Now, take $y^* \in L^\infty(I)$, with $\|y^*\|_\infty \leq 1$, calculate this y^* on $(f(\cdot, \psi_{\alpha_{\beta_\gamma}}(\cdot)) - f(\cdot, \psi_0(\cdot)))$

$$\begin{aligned} & \left| \int_0^1 y^*(t) [f(t, \psi_{\alpha_{\beta_\gamma}}(t)) - f(t, \psi_0(t))] dt \right| \leq \\ & \leq \int_{I_0} |y^*(t)| |f(t, \psi_{\alpha_{\beta_\gamma}}(t)) - f(t, \psi_0(t))| dt + \\ & + \int_{I \setminus I_0} |y^*(t)| |f(t, \psi_{\alpha_{\beta_\gamma}}(t)) - f(t, \psi_0(t))| dt \leq \\ & \leq \int_{I_0} |f(t, \psi_{\alpha_{\beta_\gamma}}(t)) - f(t, \psi_0(t))| dt + \int_{I \setminus I_0} 2[a(t) + b\lambda(t)] dt. \end{aligned}$$

Now, recall that (2) is true and observe that

$$\text{meas}(I \setminus I_0) \leq \sum_{i=1}^3 m(I \setminus I_i) \leq \frac{3}{4} \delta < \delta \text{ so that } \int_{I \setminus I_0} 2[a(t) + b\lambda(t)] dt < \frac{\epsilon}{2}.$$

Hence the last member of the chain of inequalities written above is smaller than ϵ for γ sufficiently large. This is what we need to show that H is weakly continuous on B_s . We are done.

References

- [1] J. APPELL, E. DE PASCALE, *Su alcuni parametri connessi con la misura di non compattezza di Hausdorff in spazi di funzioni misurabili*, Boll. Un. Mat. It. (6) 3-B (1984) 497-515.
- [2] J. BANAS; Z. KNAP, *Integrable solutions of a functional-integral equation*, Revista Matemática de la Univ. Complutense de Madrid, 2(1) (1989) 31-38.
- [3] F. S. DE BLASI, *On a property of the unit sphere in a Banach space*, Bull. Math. Soc. Sci. Math. R. S. Roumanie 21 (1977) 259-262.
- [4] N. DINULEANU, *Vector Measures*, Pergamon Press 1967.
- [5] N. DUNFORD, J. T. SCHWARTZ, *Linear operators*, part. I, Interscience 1958.
- [6] G. EMMANUELE, *Measure of weak noncompactness and fixed point theorems*, Bull. Math. Soc. Sci. Math. R. S. Roumanie 25 (1981) 253-258.
- [7] G. SCORZA-DRAGONI, *Un teorema sulle funzioni continue rispetto ad una e misurabili rispetto ad un'altra variabile*, Rend. Sem. Mat. Univ. Padova 17 (1948) 102-106.

Department of Mathematics,
University of Catania,
Viale A. Doria 6,
95125 Catania,
Italy

Recibido: 26 de abril de 1990
Revisado: 5 de julio de 1990