

Boundary Sentinels with Given Sensitivity

Gisèle MASSENGO MOPHOU and Jean-Pierre PUEL

Département de Mathématiques et Informatique
Université des Antilles et de La Guyane
Campus Fouillole
97159 Pointe-à-Pitre — Guadeloupe (FWI)
Gisele.Mophou@univ-ag.fr

Laboratoire de Mathématiques de Versailles
Université de Versailles St Quentin
45 avenue des Etats Unis
78035 Versailles — France
jppuel@math.uvsq.fr

Received: April 14, 2008

Accepted: July 30, 2008

ABSTRACT

The notion of sentinels with given sensitivity was introduced by J.-L. Lions in [10] in order to identify parameters in a problem of pollution ruled by a semilinear parabolic equation. He proves that the existence of such sentinels is reduced to the solution of exact controllability problem with constraints on the state. Reconsidering this notion of sentinels in a more general framework, we prove the existence of the new sentinels by solving a boundary null-controllability problem with constraint on the control. Our results use a Carleman inequality which is adapted to the constraint.

Key words: heat equation, optimal control, controllability, Carleman inequalities, sentinels.

2000 Mathematics Subject Classification: 35K05, 35K15, 35K20, 49J20, 93B05.

Introduction

In the models of boundary pollution problem, initial data are not completely known. Boundary conditions also are either unknown or known only on one part of the boundary, the other part being for example inaccessible to measurements. Let us formulate the problem more precisely. Let $N, M \in \mathbb{N} \setminus \{0\}$ and Ω be a bounded open subset of \mathbb{R}^N with boundary Γ of class C^2 . For a time $T > 0$, we set $\mathcal{Q} = \Omega \times (0, T)$, $\Sigma = \Gamma \times (0, T)$, and $\Sigma_0 = \Gamma_0 \times (0, T)$, where Γ_0 is a non empty open subset of Γ .

We consider the system modeling a problem of pollution [10]:

$$\begin{cases} \frac{\partial y}{\partial t} - \Delta y + f(y) = 0 & \text{in } \mathcal{Q}, \\ y = \begin{cases} \xi + \sum_{i=1}^M \lambda_i \hat{\xi}_i & \text{on } \Sigma_0, \\ 0 & \text{on } \Sigma \setminus \Sigma_0, \end{cases} \\ y(0) = y^0 + \tau \hat{y}^0 & \text{in } \Omega, \end{cases} \quad (1)$$

where:

- y represents the concentration of the pollutant.
- f is a given real function of class C^1 , globally Lipschitz.
- The boundary condition is unknown on a part Σ_0 of the boundary and represents here a pollution with a structure of the form $\xi + \sum_{i=1}^M \lambda_i \hat{\xi}_i$. In this structure, the functions ξ and $\{\hat{\xi}_i\}_{i=1}^M$ are known whereas the reals $\{\lambda_i\}_{i=1}^M$ are unknown.
- The initial condition is unknown and its structure is of the form $y^0 + \tau \hat{y}^0$ where the function y^0 is known and the term $\tau \hat{y}^0$ is unknown.

We set

$$\begin{aligned} H_0^{3/2}(\Gamma_0) &= \{ \rho \in H^{3/2}(\Gamma_0) \text{ such that } \rho = 0 \text{ on the boundary of } \Gamma_0 \}, \\ \mathcal{H}(\Sigma_0) &= \{ \rho \in H^1(0, T, H_0^{3/2}(\Gamma_0)) \text{ such that } \rho(x, 0) = 0 \text{ on } \Gamma_0 \} \end{aligned}$$

and we assume that:

- y^0 and \hat{y}^0 belong to $H_0^1(\Omega)$, ξ and $\hat{\xi}_i$ belong to $\mathcal{H}(\Sigma_0)$. Notice that if we write χ_{Σ_0} the characteristic function of the set Σ_0 , this implies that $\xi \cdot \chi_{\Sigma_0}$ and $\hat{\xi}_i \cdot \chi_{\Sigma_0}$ are elements of $\mathcal{H}(\Sigma)$ where

$$\mathcal{H}(\Sigma) = \{ \rho \in H^1(0, T, H^{3/2}(\Gamma)) \text{ such that } \rho(x, 0) = 0 \text{ on } \Gamma \}. \quad (2)$$

- The functions

$$\hat{\xi}_i \cdot \chi_{\Gamma_0}, \quad 1 \leq i \leq M, \quad \text{are linearly independent.} \quad (3)$$

- The real τ is small enough and $\|\hat{y}^0\|_{H_0^1(\Omega)} \leq 1$.
- The function f verifies (in order to simplify)

$$f(0) = 0. \quad (4)$$

For model (1), we are interested in identifying the parameters λ_i without any attempt at computing $\tau \hat{y}^0$. To identify these parameters, we use the theory of sentinel in a general framework. First of all, let us recall that the usual theory of sentinels lies on three considerations:

- A state equation represented here by (1) and where the solution $y = y(x, t, \lambda, \tau) = y(\lambda, \tau)$ depends on two families of parameters $\lambda = \{\lambda_1, \dots, \lambda_M\}$ and τ .
- An observation m_0 which is a measure of the flux of the concentration of the pollutant taken on a non-empty open subset O of $\Gamma \setminus \Gamma_0$ called observatory.
- A function S called “sentinel,” defined for $h_0 \in L^2(O \times (0, T))$ by

$$S(\lambda, \tau) = \int_0^T \int_O (h_0 + w) \frac{\partial y}{\partial \nu} dx dt \tag{5}$$

where ν is the unit exterior normal to Γ , $\frac{\partial y}{\partial \nu}$ is the derivative of y with respect to the normal ν and w , the control function. Moreover, the control w is to be found of minimal norm in $L^2(O \times (0, T))$ such that functions S satisfy

- S is stationary to the first order with respect to missing terms $\tau \hat{y}^0$:

$$\frac{\partial S}{\partial \tau}(0, 0) = 0 \quad \forall \hat{y}^0, \tag{6}$$

- S is sensitive to the first order with respect to pollution terms $\lambda_i \xi_i$:

$$\frac{\partial S}{\partial \lambda_i}(0, 0) = c_i, \quad 1 \leq i \leq M, \tag{7}$$

where $c_i, 1 \leq i \leq M$, are given constants not all identically zero.

Using the adjoint problem, J.-L. Lions in [10] shows that the existence of these sentinels is reduced to solution of exact controllability problem with constraints on the state, but he only proved results of approximate controllability. These types of approximate controllability problems with constraints on the state were also the subject of many numerical resolutions. It is in this context, for instance that J. P. Kernevez et al. use these sentinels in [1, 7] to identify parameters of pollution in a river. O. Bodart apply them in [2] to identify an unknown boundary.

Remark. To estimate the parameter λ_i , one proceeds as follows:

Assume that the solution of (1) when $\lambda = 0$ and $\tau = 0$ is known. Then, one has the following information:

$$S(\lambda, \tau) - S(0, 0) \approx \sum_{i=1}^M \lambda_i \frac{\partial S}{\partial \lambda_i}(0, 0).$$

Therefore, fixing $i \in \{1, \dots, M\}$ and choosing

$$\frac{\partial S}{\partial \lambda_j}(0, 0) = 0 \quad \text{for } j \neq i \quad \text{and} \quad \frac{\partial S}{\partial \lambda_i}(0, \dots, 0) = c_i,$$

one obtains the following estimate of the parameter λ_i :

$$\lambda_i \approx \frac{1}{c_i}(S(\lambda, \tau) - S(0, 0)).$$

Remark. J. L. Lions refers to the function S as a sentinel with given sensitivity c_i . In (7), the c_i are chosen according to the importance which is conferred to the component ξ_i of the pollution.

Remark. Notice that for the J. L. Lions’s sentinels defined by (5)–(7), the observatory $O \subset (\Gamma \setminus \Gamma_0)$ is also the support of the control function w .

In this paper, we consider the general framework where the support of the control function w may be different from the observatory O . More precisely, for any non-empty open subset γ of $\Gamma \setminus \Gamma_0$ where γ may be different from O , we look for a function $S = S(\lambda, \tau)$ solution of the following problem:

Given $h_0 \in L^2(O \times (0, T))$, find $w \in L^2(\gamma \times (0, T))$ such that

(i) the function S defined by

$$S(\lambda, \tau) = \int_0^T \int_O h_0 \frac{\partial y}{\partial \nu}(\lambda, \tau) d\Gamma dt + \int_0^T \int_\gamma w \frac{\partial y}{\partial \nu}(\lambda, \tau) d\Gamma dt. \quad (8)$$

satisfies the conditions (6) and (7);

(ii) The control w is of minimal norm in $L^2(\gamma \times (0, T))$ among “the admissible controls,” i.e.,

$$|w|_{L^2(\gamma \times (0, T))} = \min_{\bar{w} \in E} |\bar{w}|_{L^2(\gamma \times (0, T))} \quad (9)$$

where

$$E = \{ \bar{w} \in L^2(\gamma \times (0, T)), \text{ such that } (\bar{w}, S(\bar{w})) \text{ satisfies (6)–(8)} \}.$$

Under the above hypotheses on f and the data, one can prove as in [3] that there exists $\alpha > 0$ such that when

$$|\tau| + \sum_{i=1}^M |\lambda_i| \leq \alpha$$

problem (1) admits a unique solution $y = y(\lambda, \tau)$ in

$$H^{2,1}(\mathcal{Q}) = L^2(0, T; H^2(\Omega)) \cap H^1(0, T; L^2(\Omega)).$$

Moreover, if we denote by $I \subset \mathbb{R}$ a neighborhood of 0, the applications

$$\tau \longmapsto y(\lambda, \tau) \quad \text{and} \quad \lambda_i \longmapsto y(\lambda, \tau) \quad (1 \leq i \leq M)$$

are in $C^1(I; L^2(0, T; H^2(\Omega)))$. In fact, if we denote by $y_0 = y(0, 0) \in H^{2,1}(Q)$ the solution of (1) when $\lambda = 0$ and $\tau = 0$, and if we write the derivatives of y at $(0, 0)$ with respect to τ and λ_i as

$$y_\tau = \frac{\partial}{\partial \tau} y(\lambda, \tau)|_{\tau=0, \lambda_i=0}$$

and

$$y_{\lambda_i} = \frac{\partial}{\partial \lambda_i} y(\lambda, \tau)|_{\tau=0, \lambda_i=0},$$

the functions y_τ and y_{λ_i} are respectively solutions of

$$\begin{cases} \frac{\partial y_\tau}{\partial t} - \Delta y_\tau + f'(y_0)y_\tau = 0 & \text{in } \mathcal{Q}, \\ y_\tau = 0 & \text{on } \Sigma, \\ y_\tau(0) = \hat{y}^0 & \text{in } \Omega \end{cases} \tag{10}$$

and

$$\begin{cases} \frac{\partial y_{\lambda_i}}{\partial t} - \Delta y_{\lambda_i} + f'(y_0)y_{\lambda_i} = 0 & \text{in } \mathcal{Q}, \\ y_{\lambda_i} = \hat{\xi}_i \chi_{\Sigma_0} & \text{on } \Sigma, \\ y_{\lambda_i}(0) = 0 & \text{in } \Omega, \end{cases} \tag{11}$$

where $f'(y_0)$ denotes the derivative of f at point y_0 . Since f is a real C^1 and globally Lipschitz function, we have

$$a_0 = f'(y_0) \in L^\infty(\mathcal{Q}). \tag{12}$$

Consequently, \hat{y}^0 and $\hat{\xi}_i$ being respectively in $H_0^1(\Omega)$ and $\mathcal{H}(\Sigma_0)$, we deduce that problems (10) and (11) have respectively a unique solution y_τ and y_{λ_i} in $H^{2,1}(\mathcal{Q})$.

Remark. Since $y \in L^2(0, T; H^2(\Omega))$, we have $\frac{\partial y}{\partial \nu} \in L^2(0, T; H^{1/2}(\Gamma)) \subset L^2(\Sigma)$. Thus, h_0 and w being respectively in $L^2(O \times (0, T))$ and in $L^2(\gamma \times (0, T))$, relation (8) is well defined. Moreover, y_τ and y_{λ_i} belong to $L^2(0, T; H^2(\Omega))$. This means that $\frac{\partial y_\tau}{\partial \nu}$ and $\frac{\partial y_{\lambda_i}}{\partial \nu}$ belong to $L^2(0, T; H^{1/2}(\Gamma)) \subset L^2(\Sigma)$. Thus, conditions (6) and (7) are well defined.

Let χ_γ be the characteristic function of the set γ . We set

$$Y = \text{Span} \left(\frac{\partial y_{\lambda_1}}{\partial \nu} \chi_\gamma, \dots, \frac{\partial y_{\lambda_M}}{\partial \nu} \chi_\gamma \right) \tag{13}$$

the vector subspace of $L^2(\gamma \times (0, T))$ generated by the M functions $\frac{\partial y_{\lambda_i}}{\partial \nu} \chi_\gamma$, $1 \leq i \leq M$, which will be proved to be independent (see Lemma 1.2 in subsection 1.1) and we denote by Y^\perp the orthogonal of Y in $L^2(\gamma \times (0, T))$. Let also

$$Y_\theta = \frac{1}{\theta} Y$$

be the vector subspace of $L^2(\gamma \times (0, T))$ generated by the M functions $\frac{1}{\theta} \frac{\partial y_{\lambda_i}}{\partial \nu} \chi_\gamma$, $1 \leq i \leq M$, where θ is the positive function precisely defined later on by (26). Clearly, these functions will also be independent.

We consider the following controllability problem:

Given $h_0 \in L^2(O \times (0, T))$, $w_0 \in Y_\theta$, and $a_0 \in L^\infty(Q)$, find $v \in L^2(\gamma \times (0, T))$ such that

$$v \in Y^\perp, \tag{14}$$

and if $q = q(x, t, v) \in L^2(Q)$ is solution of

$$\begin{cases} -\frac{\partial q}{\partial t} - \Delta q + a_0 q = 0 & \text{in } Q, \\ q = h_0 \chi_O + (w_0 - v) \chi_\gamma & \text{on } \Sigma, \\ q(T) = 0 & \text{in } \Omega, \end{cases} \tag{15}$$

q satisfies

$$q(x, 0, v) = 0 \quad \text{in } \Omega. \tag{16}$$

Remark. Let us notice that if v exists, the set

$$\mathcal{E} = \{ \bar{v} \in Y^\perp \quad \text{such that} \quad (\bar{v}, \bar{q} = q(x, t, \bar{v})) \text{ satisfies (15), (16)} \}$$

is a non-empty, closed, and convex set in $L^2(\gamma \times (0, T))$. Therefore there exists $v \in \mathcal{E}$ of minimal norm.

The problem (14)–(16) is a null boundary controllability problem with constraint on the control. When $Y^\perp = L^2(\gamma \times (0, T))$, this problem becomes a null boundary controllability problem without constraint on the control. This problem has been studied by many authors.

In the linear case, D. Russell in [15] has proved that exact controllability for the wave equation implies exact controllability for the heat equation. Inspired by this work of D. Russell, G. Lebeau and L. Robbiano in [8] solved the problem of null boundary controllability in the case $a_0 = 0$ using observability inequalities deriving from Carleman inequalities. At the same time, O. Y. Imanuvilov and A. Fursikov in [5] obtained the same result for more general operators including variable coefficients and nonzero potentials using more directly global Carleman inequalities for the

evolution operator. They could extend their method to the case of some nonlinear heat equations where they prove that the problem of null boundary controllability holds for sufficiently small initial data. We can also mention results in this direction by Y. J. Lin Guo and W. Littman in [9].

O. Nakoulima gives in [14] a result of null controllability for the heat equation with constraint on a distributed control. His results is based on an observability inequality adapted to the constraint.

In this paper we solve the null boundary controllability problem with constraint on the control (14)–(16). This allows us to prove the existence of the sentinel for given sensitivity (6)–(9). More precisely, we have the following results:

Theorem A. *Let Ω be a bounded open subset of \mathbb{R}^N with boundary Γ of class C^2 , Γ_0 a non-empty open subset of Γ , O and γ two non-empty open subsets of $\Gamma \setminus \Gamma_0$. Let also f be real function of class C^1 , globally Lipschitz verifying (4). Then the existence of the sentinel (6)–(9) holds if and only if the null boundary controllability problem with constraint on the control (14)–(16) has a solution.*

The proof of the boundary null-controllability problem with constraint on the control (14)–(16) lies on the existence of a function θ and a Carleman inequality adapted to the constraint (see subsection 2.1) for which we have the following result:

Theorem B. *Assume that the hypotheses of Theorem A are satisfied. Then there exists a positive real weight function θ (a precise definition of θ will be given later on) such that, for any function $h_0 \in L^2(O \times (0, T))$ with $\theta h_0 \in L^2(O \times (0, T))$, there exists a unique control $\hat{v} \in L^2(\gamma \times (0, T))$ such that (\hat{v}, \hat{q}) with $\hat{q} = q(\hat{v})$ is solution of the null boundary controllability problem with constraint on the control (14)–(16) and provides a control $\hat{w} = w_0 \chi_\gamma - \hat{v}$ of the sentinel problem satisfying (9).*

Moreover, the control \hat{w} is given by

$$\hat{w} = P(w_0) + (I - P) \left(\frac{\partial \hat{\rho}}{\partial \nu} \chi_\gamma \right),$$

where P is the orthogonal projection operator from $L^2(\gamma \times (0, T))$ into Y , $w_0 \in Y_\theta$ depends on h_0 and c_i , $i \in \{1, \dots, M\}$, and will be precisely determined in (22), and $\hat{\rho}$ satisfies

$$\begin{cases} \frac{\partial \hat{\rho}}{\partial t} - \Delta \hat{\rho} + a_0 \hat{\rho} = 0 & \text{in } \mathcal{Q}, \\ \hat{\rho} = 0 & \text{on } \Sigma. \end{cases}$$

The paper is organized as follows: Section 1 is devoted to the equivalence between the sentinel problem and the controllability problem with constraint on the control. In this section we give the proof of Theorem A, the formulation of the sentinel and the estimate on the parameter λ_i . In section 2, we study the null-controllability problem with constraint on the control (14)–(16) and prove Theorem B.

1. Equivalence between the sentinel problem and the controllability problem with constraint on the control

1.1. Proof of Theorem A

To obtain the boundary null-controllability problem (14)–(16), we interpret (6) and (7).

In view of (8), the stationary condition (6) and respectively the sensitivity conditions (7) hold if and only if

$$\int_0^T \int_O h_0 \frac{\partial y_\tau}{\partial \nu} d\Gamma dt + \int_0^T \int_\gamma w \frac{\partial y_\tau}{\partial \nu} d\Gamma dt = 0 \quad \forall \hat{y}^0 \in H_0^1(\Omega) \quad (17)$$

and

$$\int_0^T \int_O h_0 \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt + \int_0^T \int_\gamma w \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt = c_i, \quad 1 \leq i \leq M. \quad (18)$$

Therefore, in order to transform equation (17), we introduce the classical adjoint state. More precisely, we consider the following linear problem

$$\begin{cases} -\frac{\partial q}{\partial t} - \Delta q + a_0 q = 0 & \text{in } \mathcal{Q}, \\ q = h_0 \chi_O + w \chi_\gamma & \text{on } \Sigma, \\ q(T) = 0 & \text{in } \Omega. \end{cases} \quad (19)$$

Since $h_0 \chi_O + w \chi_\gamma \in L^2(\Sigma)$, one can prove using transposition method that problem (19) admits a unique solution q in $L^2(\mathcal{Q}) \cap \mathcal{C}([0, T], H^{-1}(\Omega))$. Moreover we have $\frac{\partial q}{\partial \nu} \in H^{-1}(0, T; H^{-3/2}(\Gamma))$ (see [12]).

Remark 1.1. In fact, we have $\frac{\partial q}{\partial \nu} \in \mathcal{H}'(\Sigma)$ where $\mathcal{H}(\Sigma)$ is defined in (2). In order to show this, we can take $T' > T$, extend h_0 and w by zero on (T, T') and take $q(T') = 0$ so that $q = 0$ on (T, T') . Now if $\xi \in \mathcal{H}(\Sigma)$, we can extend it smoothly on (T, T') so that $\xi(T') = 0$. Therefore $\xi \in H_0^1(0, T'; H^{3/2}(\Gamma))$ and the duality $\langle \frac{\partial q}{\partial \nu}, \xi \rangle_{H^{-1}(0, T'; H^{-3/2}(\Gamma)), H_0^1(0, T'; H^{3/2}(\Gamma))}$ makes perfect sense with $\frac{\partial q}{\partial \nu} = 0$ on (T, T') .

Now, multiplying both sides of the differential equation in (19) by y_τ solution of (10) and integrating by parts in \mathcal{Q} , we get for all $\hat{y}^0 \in H_0^1(\Omega)$ that

$$\int_0^T \int_O h_0 \frac{\partial y_\tau}{\partial \nu} d\Gamma dt + \int_0^T \int_\gamma w \frac{\partial y_\tau}{\partial \nu} d\Gamma dt = -\langle q(0), \hat{y}^0 \rangle_{H^{-1}(\Omega), H_0^1(\Omega)}$$

where $\langle \cdot, \cdot \rangle_{X, X'}$ represents now and in the sequel, the duality bracket between the space X and X' .

Thus, condition (6) (or (17)) holds if and only if

$$q(0) = 0 \quad \text{in } \Omega. \quad (20)$$

Then, multiplying both sides of the differential equation in (19) by y_{λ_i} solution of (11) and integrating by parts in \mathcal{Q} , we have

$$\left\langle \frac{\partial q}{\partial \nu}, \hat{\xi}_i \chi_{\Gamma_0} \right\rangle_{\mathcal{H}'(\Sigma), \mathcal{H}(\Sigma)} = \int_0^T \int_O h_0 \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt + \int_0^T \int_{\gamma} w \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt, \quad 1 \leq i \leq M.$$

Thus, condition (7) (or (18)) is equivalent to

$$\left\langle \frac{\partial q}{\partial \nu}, \hat{\xi}_i \chi_{\Gamma_0} \right\rangle_{\mathcal{H}'(\Sigma), \mathcal{H}(\Sigma)} = c_i, \quad 1 \leq i \leq M. \tag{21}$$

The above considerations show that the existence of the sentinel of given sensitivity (6)–(9) is reduced to the solution of the following null controllability problem:

Given $h_0 \in L^2(O \times (0, T))$ and $a_0 \in L^\infty(\mathcal{Q})$, find the control w of minimal norm in $L^2(\gamma \times (0, T))$ such that the pair (w, q) is solution of (19)–(21).

This new problem is a null-controllability problem with constraints (21) on the state q . Actually, let us show that the insensitivity conditions (7) (or the constraints (21)) on the state q are equivalent to constraint on the control.

Lemma 1.2. *Assume that (3) holds. Then the functions $\frac{\partial y_{\lambda_i}}{\partial \nu} \chi_{\gamma}$, $1 \leq i \leq M$, are linearly independent. Moreover the functions $\frac{1}{\theta} \frac{\partial y_{\lambda_i}}{\partial \nu} \chi_{\gamma}$, $1 \leq i \leq M$, are also linearly independent.*

Proof. Let $\alpha_i \in \mathbb{R}$, $1 \leq i \leq M$ be such that $\sum_{i=1}^M \alpha_i \frac{\partial y_{\lambda_i}}{\partial \nu} \chi_{\gamma} = 0$. Then, in view of (11), $k = \sum_{i=1}^M \alpha_i y_{\lambda_i}$ is such that

$$\left\{ \begin{array}{ll} \frac{\partial k}{\partial t} - \Delta k + f'(y_0)k = 0 & \text{in } \mathcal{Q}, \\ k = \sum_{i=1}^M \alpha_i \hat{\xi}_i \chi_{\Sigma_0} & \text{on } \Sigma, \\ \frac{\partial k}{\partial \nu} = 0 & \text{on } \gamma \times (0, T), \\ k(0) = 0 & \text{in } \Omega. \end{array} \right.$$

As $\gamma \subset \Gamma \setminus \Gamma_0$, we have $k = 0$ and $\frac{\partial k}{\partial \nu} = 0$ on $\gamma \times (0, T)$. Thanks to the unique continuation property ([13, 16]), this implies $k = 0$ in \mathcal{Q} . Therefore, we deduce that $\sum_{i=1}^M \alpha_i \hat{\xi}_i = 0$ on $\Gamma_0 \times (0, T)$. Hence, assumption (3) allows to conclude that $\alpha_i = 0$ for $1 \leq i \leq M$.

The second assertion of the lemma follows immediately. □

Now the matrix

$$\left(\int_0^T \int_{\gamma} \frac{1}{\theta} \frac{\partial y_{\lambda_j}}{\partial \nu} \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt \right)_{i,j}$$

is symmetric positive definite and, therefore, there exists a unique $w_0 \in Y_\theta$ such that

$$c_i - \int_0^T \int_O h_0 \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt = \int_0^T \int_\gamma w_0 \frac{\partial y_{\lambda_i}}{\partial \nu} d\Gamma dt, \quad 1 \leq i \leq M. \tag{22}$$

Consequently, combining (18) with (22), we observe that condition (7) (or the constraints (21)) holds if and only if

$$w - w_0 = -v \in Y^\perp,$$

where Y is given by (13). Replacing w by $-v + w_0$ in the second expression of (19), we obtain the second one in (15). Therefore, the sentinel problem (6)–(9) holds if and only if null boundary controllability problem with constraint on the control (14)–(16) has a solution. \square

Remark 1.3. The function w_0 is such that $\theta w_0 \in L^2(\gamma \times (0, T))$. The choice of w_0 in Y_θ will be necessary for the construction of the optimal control for the null boundary controllability problem (14)–(16) in section 2.

Remark 1.4. If \mathcal{E} is the set of admissible controls $v \in L^2(\gamma \times (0, T))$ such that (14)–(16) is satisfied, then \mathcal{E} is a closed convex subset of $L^2(\gamma \times (0, T))$. Since $w_0 - \mathcal{E}$ is also a closed convex subset of $L^2(\gamma \times (0, T))$, we can obtain w to be of minimal norm in $L^2(\gamma \times (0, T))$ by minimizing the norm of $w_0 - v$ when $v \in \mathcal{E}$. Then the pair $(v, q(v))$ satisfying (14)–(16) necessarily provides a control w satisfying (9).

1.2. Formulation of the sentinel with given sensitivity and identification of parameter λ_i

Assume that Theorem B holds. Then if we replace in (8) w with

$$\hat{w} = P(w_0) + (I - P) \frac{\partial \hat{\rho}}{\partial \nu} \chi_\gamma,$$

the function S defined by

$$S(\lambda, \tau) = \int_0^T \int_O h_0 \frac{\partial y}{\partial \nu}(\lambda, \tau) d\Gamma dt + \int_0^T \int_\gamma \left(P(w_0) + (I - P) \frac{\partial \hat{\rho}_\theta}{\partial \nu} \chi_\gamma \right) \frac{\partial y}{\partial \nu}(\lambda, \tau) d\Gamma dt$$

is such that (\hat{w}, S) verifies (6)–(9). Therefore proceeding as in the remark in page 167, we deduce that

$$\lambda_i \approx \frac{1}{c_i} \left\{ \int_0^T \int_O h_0 \left(m_0 - \frac{\partial y_0}{\partial \nu} \right) d\Gamma dt \right\} + \frac{1}{c_i} \left\{ \int_0^T \int_\gamma \left(P(w_0) + (I - P) \frac{\partial \hat{\rho}_\theta}{\partial \nu} \chi_\gamma \right) \left(m_0 - \frac{\partial y_0}{\partial \nu} \right) d\Gamma dt \right\},$$

where m_0 is a measure of the flux of the concentration of the pollutant taken on the observatory $O \cup \gamma$ and y_0 is solution of (1) when $\lambda = 0$ and $\tau = 0$.

2. Study of the boundary null-controllability problem with constraint on the control

In this section, we prove existence of the solution of the boundary null controllability problem (14)–(16) and of course uniqueness if we want the control to be of minimal norm among admissible controls. The main tool we use is an observability inequality adapted to the constraint (14) which itself will be a consequence of an adapted Carleman inequality.

2.1. An adapted Carleman inequality

Let us consider an auxiliary function $\psi \in C^2(\bar{\Omega})$ which satisfies the following conditions:

$$\begin{aligned} \Psi(x) &> 0 \quad \forall x \in \Omega, \\ |\nabla \Psi| &> \alpha > 0 \quad \forall x \in \bar{\Omega}, \\ \Psi(x) &= 0 \quad \text{on } \Gamma \setminus \gamma, \\ \frac{\partial \Psi}{\partial \nu}(x) &< 0 \quad \forall x \in \Gamma \setminus \gamma. \end{aligned} \tag{23}$$

Such a function ψ exists according to A. Fursikov and O. Yu. Imanuvilov [5].

For $(x, t) \in \mathcal{Q}$, we define for any positive parameter value λ the following weight functions:

$$\varphi(x, t) = \frac{e^{\lambda(m|\Psi|_\infty + \Psi(x))}}{t(T-t)}, \tag{24}$$

$$\eta(x, t) = \frac{e^{2\lambda m|\Psi|_\infty} - e^{\lambda(m|\Psi|_\infty + \Psi(x))}}{t(T-t)} \tag{25}$$

with $m > 1$. Weight functions of this kind were first introduced by O. Yu. Imanuvilov. Since φ does not vanish in \mathcal{Q} , for all $s > 0$ and $\lambda > 0$, we set

$$\frac{1}{\theta^2} = \min \left[e^{-2s\eta} \left(\varphi^{-1}, \varphi, \varphi^3, \varphi \left| \frac{\partial \Psi}{\partial \nu} \right| \right) \right] \tag{26}$$

and we adopt the following notations:

$$\begin{aligned} L &= \frac{\partial}{\partial t} - \Delta + a_0 I, \\ L^* &= -\frac{\partial}{\partial t} - \Delta + a_0 I, \\ \mathcal{V} &= \{ \rho \in C^\infty(\bar{\mathcal{Q}}), \quad \rho = 0 \text{ on } \Sigma \}, \end{aligned} \tag{27}$$

where $a_0 \in L^\infty(Q)$ is defined by (12). Then, using the notations given by (27), we have the following inequality:

Proposition 2.1 (Adapted Carleman inequality). *Assume that (3) holds. Let Y be the real vector subspace of $L^2(\gamma \times (0, T))$ of finite dimension defined in (13) and P be the orthogonal projection operator from $L^2(\gamma \times (0, T))$ into Y . Let also θ be the function defined by (26). Then, there exist numbers $\lambda_0 = \lambda_0(\Omega, \gamma, a_0) > 1$, $s_0 = s_0(\Omega, \gamma, a_0, T) > 1$, $C = C(\Omega, \gamma, a_0) > 0$ such that, for fixed $\lambda \geq \lambda_0$ and $s \geq s_0$ and for any $\rho \in \mathcal{V}$,*

$$\int_0^T \int_\Gamma \frac{1}{\theta^2} \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt \leq C \left[\int_0^T \int_\Omega |L\rho|^2 dx dt + \int_0^T \int_\gamma \left| P \frac{\partial \rho}{\partial \nu} - \frac{\partial \rho}{\partial \nu} \chi_\gamma \right|^2 d\Gamma dt \right]. \tag{28}$$

To prove Proposition 2.1, we need some preliminary results.

Proposition 2.2 (Boundary Carleman inequality). *Let Ψ , φ and η be the functions defined respectively by (23)–(25). Then, there exist numbers $\lambda_0 = \lambda_0(\Omega, \gamma, a_0) > 1$, $s_0 = s_0(\Omega, \gamma, a_0, T) > 1$, $C_0 = C_0(\Omega, \gamma, a_0) > 0$, and $C_1 = C_1(\Omega, \gamma, a_0) > 0$ such that, for any $\lambda \geq \lambda_0$, for any $s \geq s_0$, and for any $\rho \in \mathcal{V}$, the following estimate holds:*

$$\begin{aligned} & \int_0^T \int_\Omega e^{-2s\eta} \left((s\varphi)^{-1} \left(\left| \frac{\partial \rho}{\partial t} \right|^2 + |\Delta\rho|^2 \right) + s\lambda^2\varphi |\nabla\rho|^2 + s^3\lambda^4\varphi^3 |\rho|^2 \right) dx dt \\ & \quad + C_0 \int_0^T \int_{\Gamma \setminus \gamma} s e^{-2s\eta} \varphi \left(-\frac{\partial \Psi}{\partial \nu} \right) \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt \\ & \leq C_1 \left[\int_0^T \int_\Omega e^{-2s\eta} |L\rho|^2 dx dt + \int_0^T \int_\gamma s e^{-2s\eta} \varphi \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt \right]. \tag{29} \end{aligned}$$

Proof. See [4, 6]. □

As a consequence of Proposition 2.2, we have this other inequality:

Proposition 2.3. *Let θ be defined as in (26). Then, there exist numbers $\lambda_0 = \lambda_0(\Omega, \gamma, a_0) > 1$, $s_0 = s_0(\Omega, \gamma, a_0, T) > 1$, $C_0 = C_0(\Omega, \gamma, a_0) > 0$, and $C_1 = C_1(\Omega, \gamma, a_0) > 0$ such that, for fixed $\lambda \geq \lambda_0$ and $s \geq s_0$ and for any $\rho \in \mathcal{V}$,*

$$\begin{aligned} & \int_0^T \int_\Omega \frac{1}{\theta^2} \left(\left| \frac{\partial \rho}{\partial t} \right|^2 + |\Delta\rho|^2 + |\nabla\rho|^2 + |\rho|^2 \right) dx dt + C_0 \int_0^T \int_\Gamma \frac{1}{\theta^2} \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt \\ & \leq C_1 \left[\int_0^T \int_\Omega |L\rho|^2 dx dt + \int_0^T \int_\gamma \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt \right]. \tag{30} \end{aligned}$$

Proof. As Ψ belongs to $\mathcal{C}^2(\bar{\Omega})$ and $\varphi e^{-2s\eta}$ is bounded, it is immediate that $\frac{1}{\theta}$ is also bounded in Q . Consequently, adding the term $\int_0^T \int_\gamma s\varphi e^{-2s\eta} \left| \frac{\partial \Psi}{\partial \nu} \right| \left| \frac{\partial \rho}{\partial \nu} \right|^2 d\Gamma dt$ to each side of (29), then using the property of Ψ given by the fourth condition in (23) and the fact that $s > 1$ and $\lambda > 1$, we deduce (30). □

Lemma 2.4. *Assume that (3) holds. Let Y be defined by (13). Then any function ρ such that*

$$\begin{cases} \frac{\partial \rho}{\partial t} - \Delta \rho + a_0 \rho = 0 & \text{in } \mathcal{Q}, \\ \rho = 0 & \text{on } \Sigma, \\ \frac{\partial \rho}{\partial \nu} \Big|_{\gamma} \in Y \end{cases} \quad (31)$$

is identically zero.

Proof. For any ρ verifying (31), there exists $\alpha_i \in \mathbb{R}$, $1 \leq i \leq M$, such that $\frac{\partial \rho}{\partial \nu} = \sum_{i=1}^M \alpha_i \frac{\partial y_{\lambda_i}}{\partial \nu}$ on γ . We set $z = \rho - \sum_{i=1}^M \alpha_i y_{\lambda_i}$. Then in view of (11), we have

$$\begin{cases} \frac{\partial z}{\partial t} - \Delta z + a_0 z = 0 & \text{in } \mathcal{Q}, \\ z = \sum_{i=1}^M \alpha_i \hat{\xi}_i \chi_{\Sigma_0} & \text{on } \Sigma, \\ \frac{\partial z}{\partial \nu} = 0 & \text{on } \gamma \times (0, T). \end{cases}$$

As $\gamma \subset \Gamma \setminus \Gamma_0$, we have

$$z = 0 \quad \text{and} \quad \frac{\partial z}{\partial \nu} = 0 \quad \text{on } \gamma \times (0, T).$$

Therefore, thanks to the unique continuation property (see [13, 16]), we have $z = 0$ in \mathcal{Q} . Consequently, we deduce on the one hand that $\rho = \sum_{i=1}^M \alpha_i y_{\lambda_i}$ and on the other hand that $\sum_{i=1}^M \alpha_i \hat{\xi}_i = 0$ on Σ_0 . Hence, it follows from assumption (3) that $\alpha_i = 0$ for $1 \leq i \leq M$. Thus, $\rho = 0$ in \mathcal{Q} . \square

Proof of Proposition 2.1. The proof uses a well known compactness-uniqueness argument and inequality (30). We argue by contradiction. We suppose that for any $n \in \mathbb{N}$ there exists $\rho_n \in \mathcal{V}$ such that

$$\int_0^T \int_{\Omega} |L\rho_n|^2 dx dt \leq \frac{1}{n}, \quad (32)$$

$$\int_0^T \int_{\gamma} \left| P \frac{\partial \rho_n}{\partial \nu} - \frac{\partial \rho_n}{\partial \nu} \chi_{\gamma} \right|^2 d\Gamma dt \leq \frac{1}{n} \quad (33)$$

and

$$\int_0^T \int_{\Gamma} \frac{1}{\theta^2} \left| \frac{\partial \rho_n}{\partial \nu} \right|^2 d\Gamma dt = 1. \quad (34)$$

Now let us prove that (32)–(34) yield a contradiction. We do it in three steps.

• *Step 1.* We have

$$\int_0^T \int_\gamma \frac{1}{\theta^2} \left| P \frac{\partial \rho_n}{\partial \nu} \right|^2 d\Gamma dt \leq 2 \int_0^T \int_\gamma \frac{1}{\theta^2} \left| P \frac{\partial \rho_n}{\partial \nu} - \frac{\partial \rho_n}{\partial \nu} \right|^2 d\Gamma dt + 2 \int_0^T \int_\gamma \frac{1}{\theta^2} \left| \frac{\partial \rho_n}{\partial \nu} \right|^2 d\Gamma dt.$$

Since the function $\frac{1}{\theta^2}$ is bounded in \mathcal{Q} , using (33) and (34), we conclude that there exists a constant C independent of n such that

$$\int_0^T \int_\gamma \frac{1}{\theta^2} \left| P \frac{\partial \rho_n}{\partial \nu} \right|^2 d\Gamma dt \leq C.$$

Therefore, Y being a finite dimensional vector subspace of $L^2(\gamma \times (0, T))$, we deduce that

$$\left| P \frac{\partial \rho_n}{\partial \nu} \right|_{L^2(\gamma \times (0, T))} \leq C. \tag{35}$$

Consequently, using (33) and (35), we get

$$\left| \frac{\partial \rho_n}{\partial \nu} \right|_{L^2(\gamma \times (0, T))} \leq C. \tag{36}$$

• *Step 2.* Let us define $L^2(\frac{1}{\theta}, X) = \{ \varphi \in L^2(X), \int_X \frac{1}{\theta^2} |\varphi|^2 dX < \infty \}$. Then $L^2(X) \subset L^2(\frac{1}{\theta}, X)$ and the canonical injection is continuous. Thus, according to (32) and (36), we deduce from (30) that $\rho_n, \nabla \rho_n, \Delta \rho_n$ are bounded in $L^2(\frac{1}{\theta}, \mathcal{Q})$ and $\frac{\partial \rho_n}{\partial \nu}$ is bounded in $L^2(\frac{1}{\theta}, \Sigma)$. Let us then take a subsequence still denoted by (ρ_n) such that

$$(\rho_n) \rightharpoonup \rho \text{ weakly in } L^2(\frac{1}{\theta}, \mathcal{Q}).$$

If we refer to (24) and (25) and the definition of $\frac{1}{\theta}$ given by (26), we can see that (ρ_n) is bounded in $L^2(] \beta, T - \beta[; H^2(\Omega)), \forall \beta > 0$. Then, we have in particular, for every $\beta > 0$,

$$\begin{aligned} \rho_n &\rightharpoonup \rho \text{ weakly in } L^2(] \beta, T - \beta[\times \Omega), \\ \frac{\partial \rho_n}{\partial \nu} &\rightharpoonup \frac{\partial \rho}{\partial \nu} \text{ weakly in } L^2(] \beta, T - \beta[\times \Gamma). \end{aligned}$$

This implies that

$$\begin{aligned} \rho_n &\rightharpoonup \rho \text{ weakly in } D'(\mathcal{Q}), \\ \frac{\partial \rho_n}{\partial \nu} &\rightharpoonup \frac{\partial \rho}{\partial \nu} \text{ weakly in } D'(\Sigma). \end{aligned}$$

Therefore, we get from (32) and (36) that

$$L\rho_n \longrightarrow L\rho = 0 \quad \text{strongly in } L^2(\mathcal{Q}), \tag{37}$$

$$\frac{\partial\rho_n}{\partial\nu} \rightharpoonup \frac{\partial\rho}{\partial\nu} \quad \text{weakly in } L^2(\gamma \times (0, T)). \tag{38}$$

And, since P is a compact operator, we deduce from (38) that

$$P\frac{\partial\rho_n}{\partial\nu} \longrightarrow P\frac{\partial\rho}{\partial\nu} \quad \text{strongly in } L^2(\gamma \times (0, T)). \tag{39}$$

From (33), we also have

$$P\frac{\partial\rho_n}{\partial\nu} - \frac{\partial\rho_n}{\partial\nu}\chi_\gamma \longrightarrow 0 \quad \text{strongly in } L^2(\gamma \times (0, T)). \tag{40}$$

Thus, combining (39) with (40), we obtain

$$\frac{\partial\rho_n}{\partial\nu}\chi_\gamma \longrightarrow P\frac{\partial\rho}{\partial\nu} \quad \text{strongly in } L^2(\gamma \times (0, T)). \tag{41}$$

Thanks to the uniqueness of the limit in $L^2(\gamma \times (0, T))$, the convergence relations (38) and (41) imply $P\frac{\partial\rho}{\partial\nu} = \frac{\partial\rho}{\partial\nu}\chi_\gamma$. This means that $\frac{\partial\rho}{\partial\nu}\chi_\gamma \in Y$. In short, we have proved that ρ verifies

$$\begin{cases} L\rho = 0 & \text{in } \mathcal{Q}, \\ \rho = 0 & \text{on } \Sigma, \\ \frac{\partial\rho}{\partial\nu}\chi_\gamma \in Y. \end{cases}$$

Thanks to Lemma 2.4, ρ is identically zero. Thus, $\frac{\partial\rho}{\partial\nu} = 0$ on $\gamma \times (0, T)$ and (41) becomes

$$\frac{\partial\rho_n}{\partial\nu} \longrightarrow 0 \quad \text{strongly in } L^2(\gamma \times (0, T)). \tag{42}$$

- *Step 3.* Since $(\rho_n) \in \mathcal{V}$, applying Carleman inequality (30) to ρ_n , then, passing to the limit on n , it results from (37) and (42) that $\frac{1}{\theta}\frac{\partial\rho_n}{\partial\nu} \rightarrow 0$ strongly in $L^2(\Sigma)$. Therefore, the contradiction occurs with (34). \square

2.2. Solution of the boundary null-controllability problem with constraints on the control

The proof of Theorem B can be obtained in different ways, all of them using the adapted Carleman estimate which has been proved in the previous section. Here we choose to present a proof which uses a penalization argument and which will be divided in three steps.

• *Step 1.* For every $\varepsilon > 0$ let us consider the following optimal control problem. Let w_0 be defined by (22). If $v \in Y^\perp$ let q be solution of (15). We know that $q \in C([0, T]; H^{-1}(\Omega))$ and we can define the functional

$$J_\varepsilon(v) = \frac{1}{2} \|w_0 - v\|_{L^2(\gamma \times (0, T))}^2 + \frac{1}{2\varepsilon} \|q(0)\|_{H^{-1}(\Omega)}^2$$

with $\|\varphi\|_{H^{-1}(\Omega)}^2 = \langle -\Delta)^{-1}, \varphi \rangle_{H_0^1(\Omega), H^{-1}(\Omega)}$ where $(-\Delta)^{-1}\varphi = \psi$ satisfies $-\Delta\psi = \varphi$ in Ω , $\psi = 0$ on Γ .

The optimal control problem is then to find $v_\varepsilon \in Y^\perp$ such that

$$J_\varepsilon(v_\varepsilon) = \min_{v \in Y^\perp} J_\varepsilon(v). \tag{43}$$

As Y^\perp is closed and convex (it is a linear subspace), it is classical to show that there exists a unique solution v_ε to (43) (see for example [11]). If we write q_ε the solution to (15) corresponding to v_ε , using an adjoint state ρ_ε , the triple $(q_\varepsilon, \rho_\varepsilon, v_\varepsilon)$ is solution of the following first order optimality system

$$\begin{cases} L^* q_\varepsilon = 0 & \text{in } \mathcal{Q}, \\ q_\varepsilon = h_0 \chi_O + w_0 \chi_\gamma - v_\varepsilon \chi_\gamma & \text{on } \Sigma, \\ q_\varepsilon(T) = 0 & \text{in } \Omega, \end{cases} \tag{44}$$

$$\begin{cases} L \rho_\varepsilon = 0 & \text{in } \mathcal{Q}, \\ \rho_\varepsilon = 0 & \text{on } \Sigma, \\ \rho_\varepsilon(0) = (-\Delta)^{-1} \left(\frac{1}{\varepsilon} q_\varepsilon(0) \right) & \text{in } \Omega, \end{cases} \tag{45}$$

$$v_\varepsilon = \left(w_0 \chi_\gamma - \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) - P \left(w_0 \chi_\gamma - \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \in Y^\perp. \tag{46}$$

• *Step 2.* We will here obtain estimates independent of ε on the control v_ε and on the state and adjoint state q_ε and ρ_ε .

In view of the value of v_ε given in (46), in order to obtain bounds for v_ε it is enough to obtain bounds for $\left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma - P \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right)$.

Let us multiply the state equation (44) by ρ_ε . We obtain

$$\langle \rho_\varepsilon(0), q_\varepsilon(0) \rangle_{H_0^1(\Omega), H^{-1}(\Omega)} + \int_0^T \int_\Gamma q_\varepsilon \frac{\partial \rho_\varepsilon}{\partial \nu} d\Gamma dt = 0$$

and therefore

$$\frac{1}{\varepsilon} \|q_\varepsilon(0)\|_{H^{-1}(\Omega)}^2 + \int_0^T \int_\gamma (w_0 - v_\varepsilon) \frac{\partial \rho_\varepsilon}{\partial \nu} d\Gamma dt + \int_0^T \int_O h_0 \frac{\partial \rho_\varepsilon}{\partial \nu} d\Gamma dt = 0,$$

so that

$$\begin{aligned} \frac{1}{\varepsilon} |q_\varepsilon(0)|_{H^{-1}(\Omega)}^2 + \int_0^T \int_\gamma |w_0 - v_\varepsilon|^2 d\Gamma dt + \int_0^T \int_O h_0 \frac{\partial \rho_\varepsilon}{\partial \nu} d\Gamma dt \\ - \int_0^T \int_\gamma (w_0 - v_\varepsilon) P \left(w_0 \chi_\gamma - \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) d\Gamma dt = 0. \end{aligned}$$

As $v_\varepsilon \in Y^\perp$, this gives

$$\begin{aligned} 2J_\varepsilon(v_\varepsilon) = \int_0^T \int_\gamma |P w_0 \chi_\gamma|^2 d\Gamma dt + \int_0^T \int_\gamma w_0 (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) d\Gamma dt \\ - \int_0^T \int_\gamma w_0 \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma d\Gamma dt - \int_0^T \int_O h_0 \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma d\Gamma dt. \end{aligned}$$

As we have on $\gamma \times (0, T)$

$$w_0 - v_\varepsilon = P w_0 \chi_\gamma + (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right)$$

we have

$$\int_0^T \int_\gamma |w_0 - v_\varepsilon|^2 d\Gamma dt = \int_0^T \int_\gamma |P w_0 \chi_\gamma|^2 d\Gamma dt + \int_0^T \int_\gamma \left| (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \right|^2 d\Gamma dt$$

and we obtain

$$\begin{aligned} \frac{1}{\varepsilon} |q_\varepsilon(0)|_{H^{-1}(\Omega)}^2 + \int_0^T \int_\gamma \left| (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \right|^2 d\Gamma dt \\ = \int_0^T \int_\gamma w_0 (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) d\Gamma dt \\ - \int_0^T \int_\gamma w_0 \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma d\Gamma dt - \int_0^T \int_O h_0 \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma d\Gamma dt. \end{aligned}$$

This implies

$$\begin{aligned} \frac{1}{\varepsilon} |q_\varepsilon(0)|_{H^{-1}(\Omega)}^2 + \frac{1}{2} \int_0^T \int_\gamma \left| (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \right|^2 d\Gamma dt \\ \leq \frac{1}{2} \int_0^T \int_\gamma |w_0|^2 d\Gamma dt + \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2} \\ \times \left(\int_0^T \int_\Gamma \frac{1}{\theta^2} \left| \frac{\partial \rho_\varepsilon}{\partial \nu} \right|^2 d\Gamma dt \right)^{1/2}. \quad (47) \end{aligned}$$

Now, if we apply the adapted Carleman inequality (28) to ρ_ε , we obtain

$$\int_0^T \int_\Gamma \frac{1}{\theta^2} \left| \frac{\partial \rho_\varepsilon}{\partial \nu} \right|^2 d\Gamma dt \leq C \int_0^T \int_\gamma \left| (I - P) \frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right|^2 d\Gamma dt. \quad (48)$$

From (47), the hypothesis on h_0 and the choice of $w_0 \in Y_\theta$, we see that

$$\begin{aligned} & \frac{1}{\varepsilon} |q_\varepsilon(0)|_{H^{-1}(\Omega)}^2 + \int_0^T \int_\gamma \left| (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \right|^2 d\Gamma dt \\ & \leq C \left(\int_0^T \int_\gamma |w_0|^2 d\Gamma dt + \int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right). \end{aligned}$$

This implies

$$\begin{aligned} & \left| (I - P) \left(\frac{\partial \rho_\varepsilon}{\partial \nu} \chi_\gamma \right) \right|_{L^2(\gamma \times (0, T))} \\ & \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}, \quad (49a) \end{aligned}$$

$$|v_\varepsilon|_{L^2(\gamma \times (0, T))} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}, \quad (49b)$$

$$|q_\varepsilon(0)|_{H^{-1}(\Omega)} \leq C \sqrt{\varepsilon} \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}, \quad (49c)$$

where $C > 0$ is independent of ε . Then, the properties of the heat equation (44) allow us to conclude that

$$|q_\varepsilon|_{L^2(\mathcal{Q})} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}. \quad (50)$$

From (49a) and (48), we see that

$$\left| \frac{1}{\theta} \frac{\partial \rho_\varepsilon}{\partial \nu} \right|_{L^2(\Sigma)} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}$$

and using again (49a) and the fact that $\frac{1}{\theta}$ is bounded, we obtain

$$\left| \frac{1}{\theta} P \frac{\partial \rho_\varepsilon}{\partial \nu} \right|_{L^2(\gamma \times (0, T))} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}$$

Therefore, Y being a finite dimensional vector subspace of $L^2(\gamma \times (0, T))$, we deduce that

$$\left| P \frac{\partial \rho_\varepsilon}{\partial \nu} \right|_{L^2(\gamma \times (0, T))} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}.$$

Therefore, from (49a) we obtain

$$\left| \frac{\partial \rho_\varepsilon}{\partial \nu} \right|_{L^2(\gamma \times (0, T))} \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right)^{1/2}. \tag{51}$$

Using Proposition 2.3 we then have

$$\begin{aligned} \int_0^T \int_\Omega \frac{1}{\theta^2} \left(\left| \frac{\partial \rho_\varepsilon}{\partial t} \right|^2 + |\Delta \rho_\varepsilon|^2 + |\nabla \rho_\varepsilon|^2 + |\rho_\varepsilon|^2 \right) dx dt \\ \leq C \left(\int_0^T \int_\gamma \theta^2 |w_0|^2 d\Gamma dt + \int_0^T \int_O \theta^2 |h_0|^2 d\Gamma dt \right) \end{aligned} \tag{52}$$

• *Step 3.* In view of (49), (50), and (52), we can extract subsequences of (v_ε) , (q_ε) , and ρ_ε (still called (v_ε) , (q_ε) , and ρ_ε) such that

$$\begin{aligned} v_\varepsilon &\rightharpoonup \tilde{v} && \text{weakly in } L^2(\gamma \times (0, T)), \\ q_\varepsilon &\rightharpoonup \tilde{q} && \text{weakly in } L^2(\mathcal{Q}), \\ \rho_\varepsilon &\rightharpoonup \tilde{\rho} && \text{weakly in } L^2\left(\frac{1}{\theta}, \mathcal{Q}\right). \end{aligned} \tag{53}$$

Therefore, $\tilde{v} \in Y^\perp$, $L^*q_\varepsilon \rightarrow L^*\tilde{q}$ weakly in $D'(\mathcal{Q})$ and, in view of the first equality in (44), we have

$$L^*\tilde{q} = 0 \quad \text{in } \mathcal{Q}.$$

Thus, $\tilde{q} \in L^2(\mathcal{Q})$ and $L^*\tilde{q} \in L^2(\mathcal{Q})$. Consequently, we can give a sense to the traces \tilde{q} on Σ and $\frac{\partial \tilde{q}}{\partial \nu}$ on Σ respectively in $H^{-1}(0, T, H^{-1/2}(\Gamma))$ and in $H^{-1}(0, T, H^{-3/2}(\Gamma))$ on the one hand, and to $\tilde{q}(T)$ and $\tilde{q}(0)$ in $H^{-1}(\Omega)$ on the other hand. Now, it can be shown that

$$q_\varepsilon(0) \longrightarrow \tilde{q}(0) \quad \text{in } \mathcal{D}'(\Omega)$$

but from (49c) we know that

$$q_\varepsilon(0) \longrightarrow 0 \quad \text{in } H^{-1}(\Omega) \text{ strongly,}$$

so that

$$\tilde{q}(0) = 0.$$

Therefore it is clear that (\tilde{v}, \tilde{q}) verifies (14)–(16) and there exists a solution to the boundary null-controllability problem. Moreover, it is clear from (45) that $\tilde{\rho}$ satisfies

$$\begin{cases} L\tilde{\rho} = 0 & \text{in } \mathcal{Q}, \\ \tilde{\rho} = 0 & \text{on } \Sigma. \end{cases}$$

Notice that $\tilde{\rho}$ may not have an initial value at time 0. Nevertheless because of (51) we see that

$$\frac{\partial \rho_\varepsilon}{\partial \nu} \rightharpoonup \frac{\partial \tilde{\rho}}{\partial \nu} \quad \text{weakly in } L^2(\gamma \times (0, T))$$

and therefore

$$\tilde{v} = (I - P) \left(w_0 \chi_\gamma - \frac{\partial \tilde{\rho}}{\partial \nu} \chi_\gamma \right).$$

As there exists a solution to the boundary null-controllability problem, it has already been noticed that we can find a unique $\hat{v} \in \mathcal{E}$ (admissible control) such that $(w_0 - \hat{v})$ is of minimal norm in $L^2(\gamma \times (0, T))$. If we denote by \hat{q} the corresponding solution of (15), we have $\hat{q}(0) = 0$ and, as $\tilde{v} \in \mathcal{E}$,

$$\frac{1}{2} |w_0 - v_\varepsilon|_{L^2(\gamma \times (0, T))}^2 \leq J_\varepsilon(v_\varepsilon) \leq J_\varepsilon(\hat{v}) = \frac{1}{2} |w_0 - \hat{v}|_{L^2(\gamma \times (0, T))}^2$$

and

$$\frac{1}{2} |w_0 - \hat{v}|_{L^2(\gamma \times (0, T))}^2 \leq \frac{1}{2} |w_0 - \tilde{v}|_{L^2(\gamma \times (0, T))}^2.$$

But because of (53),

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{2} |w_0 - v_\varepsilon|_{L^2(\gamma \times (0, T))}^2 \geq \frac{1}{2} |w_0 - \tilde{v}|_{L^2(\gamma \times (0, T))}^2.$$

Therefore we have

$$\tilde{v} = \hat{v}$$

and

$$v_\varepsilon \longrightarrow \hat{v} \quad \text{strongly in } L^2(\gamma \times (0, T)).$$

Writing $\hat{\rho} = \tilde{\rho}$ we have

$$\hat{v} = (I - P) \left(w_0 \chi_\gamma - \frac{\partial \hat{\rho}}{\partial \nu} \chi_\gamma \right).$$

This finishes the proof of Theorem B. □

References

- [1] B.-E. Aïnseba, J.-P. Kernévez, and R. Luce, *Identification de paramètres dans les problèmes non linéaires à données incomplètes*, RAIRO Modél. Math. Anal. Numér. **28** (1994), no. 3, 313–328 (French, with English and French summaries).
- [2] O. Bodart and P. Demeestere, *Sentinels for the identification of an unknown boundary*, Math. Models Methods Appl. Sci. **7** (1997), no. 6, 871–885.
- [3] T. Cazenave and A. Haraux, *Introduction aux problèmes d'évolution semi-linéaires*, Mathématiques & Applications, vol. 1, Ellipses, Paris, 1990.
- [4] E. Fernández-Cara and S. Guerrero, *Global Carleman inequalities for parabolic systems and applications to controllability*, SIAM J. Control Optim. **45** (2006), no. 4, 1399–1446.

- [5] A. V. Fursikov and O. Yu. Imanuilov, *Controllability of evolution equations*, Lecture Notes, vol. 34, Research Institute of Mathematics, Seoul National University, Seoul, 1996.
- [6] O. Yu. Imanuilov, *Controllability of parabolic equations*, Mat. Sb. **186** (1995), no. 6, 109–132 (Russian); English transl., Sb. Math. **186** (1995), no. 6, 879–900.
- [7] J.-P. Kernévez, *The sentinel method and its application to environmental pollution problems*, CRC Mathematical Modelling Series, CRC Press, Boca Raton, FL, 1997.
- [8] G. Lebeau and L. Robbiano, *Contrôle exact de l'équation de la chaleur*, Comm. Partial Differential Equations **20** (1995), no. 1-2, 335–356.
- [9] Y.-J. Lin Guo and W. Littman, *Null boundary controllability for semilinear heat equations*, Appl. Math. Optim. **32** (1995), no. 3, 281–316.
- [10] J.-L. Lions, *Sentinelles pour les systèmes distribués à données incomplètes*, Recherches en Mathématiques Appliquées, vol. 21, Masson, Paris, 1992.
- [11] ———, *Optimal control of systems governed by partial differential equations*, Grundlehren der mathematischen Wissenschaften, vol. 170, Springer-Verlag, New York, 1971.
- [12] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications*. 1, Travaux et Recherches Mathématiques **17**; 2, Travaux et Recherches Mathématiques **18**, Dunod, Paris, 1968.
- [13] S. Mizohata, *Unicité du prolongement des solutions pour quelques opérateurs différentiels paraboliques*, Mem. Coll. Sci. Univ. Kyoto. Ser. A. Math. **31** (1958), 219–239.
- [14] O. Nakoulima, *Contrôlabilité à zéro avec contraintes sur le contrôle*, C. R. Math. Acad. Sci. Paris **339** (2004), no. 6, 405–410 (French, with English and French summaries).
- [15] D. L. Russell, *A unified boundary controllability theory for hyperbolic and parabolic partial differential equations*, Studies in Appl. Math. **52** (1973), 189–211.
- [16] J.-C. Saut and B. Scheurer, *Unique continuation for some evolution equations*, J. Differential Equations **66** (1987), no. 1, 118–139.