

# The $SL(2, \mathbb{C})$ -Character Varieties of Torus Knots

Vicente Muñoz

Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM  
Consejo Superior de Investigaciones Científicas  
Serrano 113 bis  
28006 Madrid — Spain  
vicente.munoz@imaff.cfmac.csic.es

Facultad de Matemáticas  
Universidad Complutense de Madrid  
Plaza Ciencias 3  
28040 Madrid — Spain

Received: September 29, 2008

Accepted: January 13, 2009

## ABSTRACT

Let  $G$  be the fundamental group of the complement of the torus knot of type  $(m, n)$ . This has a presentation  $G = \langle x, y \mid x^m = y^n \rangle$ . We find the geometric description of the character variety  $X(G)$  of characters of representations of  $G$  into  $SL(2, \mathbb{C})$ .

*Key words:* Torus knot, characters, representations.

*2000 Mathematics Subject Classification:* Primary: 14D20. Secondary: 57M25, 57M27.

## Introduction

Since the foundational work of Culler and Shalen [1], the varieties of  $SL(2, \mathbb{C})$ -characters have been extensively studied. Given a manifold  $M$ , the variety of representations of  $\pi_1(M)$  into  $SL(2, \mathbb{C})$  and the variety of characters of such representations both contain information of the topology of  $M$ . This is specially interesting for 3-dimensional manifolds, where the fundamental group and the geometrical properties of the manifold are strongly related.

This can be used to study knots  $K \subset S^3$ , by analysing the  $SL(2, \mathbb{C})$ -character variety of the fundamental group of the knot complement  $S^3 - K$ . In this paper, we study the case of the torus knots  $K_{m,n}$  of any type  $(m, n)$ . The case  $(m, n) = (m, 2)$  was analysed in [3] and the general case was recently determined in [2] by a method different from ours.

---

Partially supported through grant MEC (Spain) MTM2007-63582

## 1. Character varieties

A *representation* of a group  $G$  in  $\mathrm{SL}(2, \mathbb{C})$  is a homomorphism  $\rho : G \rightarrow \mathrm{SL}(2, \mathbb{C})$ . Consider a finitely presented group  $G = \langle x_1, \dots, x_k \mid r_1, \dots, r_s \rangle$ , and let  $\rho : G \rightarrow \mathrm{SL}(2, \mathbb{C})$  be a representation. Then  $\rho$  is completely determined by the  $k$ -tuple  $(A_1, \dots, A_k) = (\rho(x_1), \dots, \rho(x_k))$  subject to the relations  $r_j(A_1, \dots, A_k) = 0$ ,  $1 \leq j \leq s$ . Using the natural embedding  $\mathrm{SL}(2, \mathbb{C}) \subset \mathbb{C}^4$ , we can identify the space of representations as

$$\begin{aligned} R(G) &= \mathrm{Hom}(G, \mathrm{SL}(2, \mathbb{C})) \\ &= \{(A_1, \dots, A_k) \in \mathrm{SL}(2, \mathbb{C})^k \mid r_j(A_1, \dots, A_k) = 0, 1 \leq j \leq s\} \subset \mathbb{C}^{4k}. \end{aligned}$$

Therefore  $R(G)$  is an affine algebraic set.

We say that two representations  $\rho$  and  $\rho'$  are equivalent if there exists  $P \in \mathrm{SL}(2, \mathbb{C})$  such that  $\rho'(g) = P^{-1}\rho(g)P$ , for every  $g \in G$ . This produces an action of  $\mathrm{SL}(2, \mathbb{C})$  in  $R(G)$ . The moduli space of representations is the GIT quotient

$$M(G) = \mathrm{Hom}(G, \mathrm{SL}(2, \mathbb{C})) // \mathrm{SL}(2, \mathbb{C}).$$

A representation  $\rho$  is *reducible* if the elements of  $\rho(G)$  all share a common eigenvector, otherwise  $\rho$  is *irreducible*.

Given a representation  $\rho : G \rightarrow \mathrm{SL}(2, \mathbb{C})$ , we define its *character* as the map  $\chi_\rho : G \rightarrow \mathbb{C}$ ,  $\chi_\rho(g) = \mathrm{tr} \rho(g)$ . Note that two equivalent representations  $\rho$  and  $\rho'$  have the same character, and the converse is also true if  $\rho$  or  $\rho'$  is irreducible [1, Proposition 1.5.2].

There is a character map  $\chi : R(G) \rightarrow \mathbb{C}^G$ ,  $\rho \mapsto \chi_\rho$ , whose image

$$X(G) = \chi(R(G))$$

is called the *character variety of  $G$* . Let us give  $X(G)$  the structure of an algebraic variety. By the results of [1], there exists a collection  $g_1, \dots, g_a$  of elements of  $G$  such that  $\chi_\rho$  is determined by  $\chi_\rho(g_1), \dots, \chi_\rho(g_a)$ , for any  $\rho$ . Such collection gives a map

$$\Psi : R(G) \rightarrow \mathbb{C}^a, \quad \Psi(\rho) = (\chi_\rho(g_1), \dots, \chi_\rho(g_a)).$$

We have a bijection  $X(G) \cong \Psi(R(G))$ . This endows  $X(G)$  with the structure of an algebraic variety. Moreover, this is independent of the chosen collection as proved in [1].

**Lemma 1.1.** *The natural algebraic map  $M(G) \rightarrow X(G)$  is a bijection.*

*Proof.* The map  $R(G) \rightarrow X(G)$  is algebraic and  $\mathrm{SL}(2, \mathbb{C})$ -invariant, hence it descends to an algebraic map  $\varphi : M(G) \rightarrow X(G)$ . Let us see that  $\varphi$  is a bijection.

For  $\rho$  an irreducible representation, if  $\varphi(\rho) = \varphi(\rho')$  then  $\rho$  and  $\rho'$  are equivalent representations; so they represent the same point in  $M(G)$ .

Now suppose that  $\rho$  is reducible. Consider  $e_1 \in \mathbb{C}^2$  the common eigenvector of all  $\rho(g)$ . This gives a sub-representation  $\rho' : G \rightarrow \mathbb{C}^*$  of  $G$ . We have a quotient

representation  $\rho'' = \rho/\rho' : G \rightarrow \mathbb{C}^*$ , defined as the representation induced by  $\rho$  in the quotient space  $\mathbb{C}^2/\langle e_1 \rangle$ . As characters,  $\rho'' = \rho'^{-1}$ . The representation  $\rho' \oplus \rho''$  is the *semisimplification* of  $\rho$ . It is in the closure of the  $SL(2, \mathbb{C})$ -orbit through  $\rho$ . Clearly,  $\chi_\rho(g) = \rho'(g) + \rho'(g)^{-1}$ . Now if  $\rho$  and  $\tilde{\rho}$  are two reducible representations and  $\varphi(\rho) = \varphi(\tilde{\rho})$ , then their semisimplifications have the same character, that is

$$\chi_\rho(g) = \chi_{\tilde{\rho}}(g) \Rightarrow \rho'(g) + \rho'(g)^{-1} = \tilde{\rho}'(g) + \tilde{\rho}'(g)^{-1}.$$

Therefore  $\rho' = \tilde{\rho}'$  or  $\rho' = \tilde{\rho}'^{-1}$ . In either case  $\rho$  and  $\tilde{\rho}$  represent the same point in  $M(G)$ , which is actually the point represented by  $\rho' \oplus \rho'^{-1}$ .  $\square$

## 2. Character varieties of torus knots

Let  $T^2 = S^1 \times S^1$  be the 2-torus and consider the standard embedding  $T^2 \subset S^3$ . Let  $m, n$  be a pair of coprime positive integers. Identifying  $T^2$  with the quotient  $\mathbb{R}^2/\mathbb{Z}^2$ , the image of the straight line  $y = \frac{m}{n}x$  in  $T^2$  defines the *torus knot* of type  $(m, n)$ , which we shall denote as  $K_{m,n} \subset S^3$  (see [4, Chapter 3]).

For any knot  $K \subset S^3$ , we denote by  $G(K)$  the fundamental group of the exterior  $S^3 - K$  of the knot. It is known that

$$G_{m,n} = G(K_{m,n}) \cong \langle x, y \mid x^m = y^n \rangle.$$

The purpose of this paper is to describe the character variety  $X(G_{m,n})$ .

In [3], the character variety  $X(G_{m,2})$  is computed. We want to extend the result to arbitrary  $m, n$ , and give a simpler argument than that of [3].

After the completion of this work, we became aware of the paper [2] where the character varieties of  $X(G_{m,n})$  are determined (even without the assumption of  $m, n$  being coprime). However, our method is more direct than the one presented in [2].

To start with, note that

$$R(G_{m,n}) = \{(A, B) \in SL(2, \mathbb{C}) \mid A^m = B^n\}.$$

Therefore we shall identify a representation  $\rho$  with a pair of matrices  $(A, B)$  satisfying the required relation  $A^m = B^n$ .

We decompose the character variety

$$X(G_{m,n}) = X_{red} \cup X_{irr},$$

where  $X_{red}$  is the subset consisting of the characters of reducible representations (which is a closed subset by [1]), and  $X_{irr}$  is the closure of the subset consisting of the characters of irreducible representations.

**Proposition 2.1.** *There is an isomorphism  $X_{red} \cong \mathbb{C}$ . The correspondence is defined by*

$$\rho = \left( A = \begin{pmatrix} t^n & 0 \\ 0 & t^{-n} \end{pmatrix}, B = \begin{pmatrix} t^m & 0 \\ 0 & t^{-m} \end{pmatrix} \right) \mapsto s = t + t^{-1} \in \mathbb{C}.$$

*Proof.* By the discussion in Lemma 1.1, an element in  $X_{red}$  is described as the character of a split representations  $\rho = \rho' \oplus \rho'^{-1}$ . This means that in a suitable basis,

$$A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix}.$$

The equality  $A^m = B^n$  implies  $\lambda^m = \mu^n$ . Therefore there is a unique  $t \in \mathbb{C}$  with  $t \neq 0$  such that

$$\begin{cases} \lambda = t^n, \\ \mu = t^m. \end{cases}$$

(Here we use the coprimality of  $(m, n)$ ). Note that the pair  $(A, B)$  is well-defined up to permuting the two vectors in the basis. This corresponds to the change  $(\lambda, \mu) \mapsto (\lambda^{-1}, \mu^{-1})$ , which in turn corresponds to  $t \mapsto t^{-1}$ . So  $(A, B)$  is parametrized by  $s = t + t^{-1} \in \mathbb{C}$ .  $\square$

**Lemma 2.2.** *Suppose that  $\rho = (A, B) \in R(G_{m,n})$ . In any of the following cases:*

- (a)  $A^m = B^n \neq \pm \text{Id}$ ,
- (b)  $A = \pm \text{Id}$  or  $B = \pm \text{Id}$ ,
- (c)  $A$  or  $B$  is non-diagonalizable,

*the representation  $\rho$  is reducible.*

*Proof.* First suppose that  $A$  is diagonalizable with eigenvalues  $\lambda, \lambda^{-1}$ , and suppose that  $\lambda^m \neq \pm 1$ . Then there is a basis  $e_1, e_2$  in which  $A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$ , which is well-determined up to multiplication of the basis vectors by non-zero scalars. Then

$$B^n = A^m = \begin{pmatrix} \lambda^m & 0 \\ 0 & \lambda^{-m} \end{pmatrix}$$

is a diagonal matrix, different from  $\pm \text{Id}$ . Therefore  $B$  must be diagonal in the same basis,  $B = \begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix}$ , with  $\lambda^m = \mu^n$ . This proves the reducibility in case (a).

Now suppose that  $A = \lambda \text{Id}$ ,  $\lambda = \pm 1$ . Then  $B^n = \lambda^m \text{Id}$ , so it must be that  $B$  is diagonalizable. Using a basis in which  $B$  is diagonal, we get the reducibility in case (b).

Finally, suppose that  $A$  is not diagonalizable. Then there is a suitable basis on which  $A$  takes the form  $A = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$ , with  $\lambda = \pm 1$ . Clearly

$$B^n = A^m = \lambda^m \begin{pmatrix} 1 & m\lambda \\ 0 & 1 \end{pmatrix}$$

and so

$$B = \begin{pmatrix} \mu & x \\ 0 & \mu \end{pmatrix},$$

with  $\mu = \pm 1$ ,  $\mu^n = \lambda^m$  and  $\mu nx = \lambda m$ . In this basis, the vector  $e_1$  is an eigenvector for both  $A$  and  $B$ . Hence the representation  $(A, B)$  is reducible, completing the case (c).  $\square$

**Proposition 2.3.** *Let  $X_{irr}^o$  be the set of irreducible characters, and  $X_{irr}$  its closure. Then*

$$\begin{aligned} X_{irr}^o &\cong \{(\lambda, \mu, r) \mid \lambda^m = \mu^n = \pm 1, \lambda \neq \pm 1, \mu \neq \pm 1, r \in \mathbb{C} - \{0, 1\}\} / \mathbb{Z}_2 \times \mathbb{Z}_2, \\ X_{irr} &\cong \{(\lambda, \mu, r) \mid \lambda^m = \mu^n = \pm 1, \lambda \neq \pm 1, \mu \neq \pm 1, r \in \mathbb{C}\} / \mathbb{Z}_2 \times \mathbb{Z}_2. \end{aligned}$$

where  $\mathbb{Z}_2 \times \mathbb{Z}_2$  acts as  $(\lambda, \mu, r) \sim (\lambda^{-1}, \mu, 1 - r) \sim (\lambda, \mu^{-1}, 1 - r) \sim (\lambda^{-1}, \mu^{-1}, r)$ .

*Proof.* Let  $\rho = (A, B)$  be an element of  $R(G_{m,n})$  which is an irreducible representation. By Lemma 2.2,  $A$  is diagonalizable but not equal to  $\pm \text{Id}$ , and  $A^m = \pm \text{Id}$ . So the eigenvalues  $\lambda, \lambda^{-1}$  of  $A$  satisfy  $\lambda^m = \pm 1$  and  $\lambda \neq \pm 1$ . Analogously,  $B$  is diagonalizable but not equal to  $\pm \text{Id}$ , with eigenvalues  $\mu, \mu^{-1}$ , with  $\mu^n = \pm 1$ ,  $\mu \neq \pm 1$ . Moreover,

$$\lambda^m = \mu^n.$$

We may choose a basis  $\{e_1, e_2\}$  under which  $A$  diagonalizes. This is well-defined up to multiplication of  $e_1$  and  $e_2$  by two non-zero scalars. Let  $\{f_1, f_2\}$  be a basis under which  $B$  diagonalizes, which is well-defined up to multiplication of  $f_1, f_2$  by non-zero scalars. Then  $\{[e_1], [e_2], [f_1], [f_2]\}$  are four points of the projective line  $\mathbb{P}^1 = \mathbb{P}(\mathbb{C}^2)$ . Note that the pair  $(A, B)$  is irreducible if and only if the four points are different.

The only invariant of four points in  $\mathbb{P}^1$  is the double ratio

$$r = ([e_1] : [e_2] : [f_1] : [f_2]) \in \mathbb{P}^1 - \{0, 1, \infty\} = \mathbb{C} - \{0, 1\}.$$

So  $(A, B)$  is parametrized, up to the action of  $SL(2, \mathbb{C})$ , by  $(\lambda, \mu, r)$ . Permuting the two basis vectors  $e_1, e_2$  corresponds to  $(\lambda, \mu, r) \mapsto (\lambda^{-1}, \mu, 1 - r)$ , since

$$([e_2] : [e_1] : [f_1] : [f_2]) = 1 - ([e_1] : [e_2] : [f_1] : [f_2]).$$

Analogously, permuting the two basis vectors  $f_1, f_2$  corresponds to

$$(\lambda, \mu, r) \mapsto (\lambda, \mu^{-1}, 1 - r).$$

Note that this gives an action of  $\mathbb{Z}_2 \times \mathbb{Z}_2$  and  $X_{irr}^o$  is the quotient of the set of  $(\lambda, \mu, r)$  as above by this action.

To describe the closure of  $X_{irr}^o$ , we have to allow  $f_1$  to coincide with  $e_1$ . This corresponds to  $r = 1$  (the same happens if  $f_2$  coincides with  $e_2$ ). In this case,  $e_1$  is

an eigenvector of both  $A$  and  $B$ , so the representation  $(A, B)$  has the same character as its semisimplification  $(A', B')$  given by

$$A' = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \quad B' = \begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix}.$$

This means that the point  $(\lambda, \mu, 1)$  corresponds under the identification  $X_{red} \cong \mathbb{C}$  given by Proposition 2.1 to  $s_1 = t_1 + t_1^{-1}$ , where  $t_1 \in \mathbb{C}$  satisfies

$$\begin{cases} \lambda = t_1^n, \\ \mu = t_1^m. \end{cases} \quad (1)$$

Also, we have to allow  $f_1$  to coincide with  $e_2$  (or  $f_2$  to coincide with  $e_1$ ). This corresponds to  $r = 0$ . The representation  $(A, B)$  has semisimplification  $(A', B')$  where

$$A' = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \quad B' = \begin{pmatrix} \mu^{-1} & 0 \\ 0 & \mu \end{pmatrix}.$$

So the point  $(\lambda, \mu, 1)$  corresponds to  $s_0 = t_0 + t_0^{-1} \in X_{red} \cong \mathbb{C}$ , where  $t_0 \in \mathbb{C}$  satisfies

$$\begin{cases} \lambda = t_0^n, \\ \mu^{-1} = t_0^m. \end{cases} \quad (2)$$

□

Proposition 2.3 says that  $X_{irr}$  is a collection of  $\frac{(m-1)(n-1)}{2}$  lines. A pair  $(\lambda, \mu)$  with  $\lambda^m = \pm 1$  and  $\mu^n = \pm 1$  is given as

$$\lambda = e^{\pi i k/m}, \quad \mu = e^{\pi i k'/n},$$

where  $0 \leq k < 2m$ ,  $0 \leq k' < 2n$ . The condition  $\lambda \neq \pm 1$ ,  $\mu \neq \pm 1$  gives  $k \neq 0, m$ ,  $k' \neq 0, n$ . Finally, the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -action allows us to restrict to  $0 < k < m$ ,  $0 < k' < n$ . The condition  $\lambda^m = \mu^n$  means that

$$k \equiv k' \pmod{2}.$$

Denote by  $X_{irr}^{k,k'}$  the line of  $X_{irr}$  corresponding to the values of  $k, k'$ . Then

$$X_{irr} = \bigsqcup_{\substack{0 < k < m, 0 < k' < n \\ k \equiv k' \pmod{2}}} X_{irr}^{k,k'}.$$

The line  $X_{irr}^{k,k'}$  intersects  $X_{red}$  in two points. This gives a collection of  $(m-1)(n-1)$  points in  $X_{red}$ , which are defined as follows: under the identification  $X_{red} \cong \mathbb{C}$ , these are the points  $s_l = t_l + t_l^{-1}$ , where

$$t_l = e^{\pi i l/nm},$$




Figure 1 – Picture of  $X(G_{m,n})$ .

and  $0 < l < mn$ ,  $m \nmid l$ ,  $n \nmid l$ . Assume that  $n$  is odd (note that either  $m$  or  $n$  should be odd). Then from (1) and (2), the line  $X_{irr}^{k,k'}$  intersects at the points  $s_{l_0}, s_{l_1} \in X_{red}$  where

$$\begin{aligned} nl_0 &\equiv k \pmod{m}, & ml_0 &\equiv n - k' \pmod{n}, \\ nl_1 &\equiv k \pmod{m}, & ml_1 &\equiv k' \pmod{n}. \end{aligned}$$

These two points are different since  $k' \not\equiv n - k' \pmod{n}$ , as  $n$  is odd.

In the case  $(m, n) = (2, n)$ , this result coincides with [3, Corollary 4.2].

### 3. The algebraic structure of $X(G_{m,n})$

We want to give a geometric realization of  $X(G_{m,n})$  which shows that the algebraic structure of this variety is that of a collection of rational lines as in Figure 1 intersecting with nodal curve singularities.

The map  $R(G_{m,n}) \rightarrow \mathbb{C}^3$ ,  $\rho = (A, B) \mapsto (\text{tr}(A), \text{tr}(B), \text{tr}(AB))$ , defines a map

$$\Psi : X(G_{m,n}) \rightarrow \mathbb{C}^3.$$

**Theorem 3.1.** *The map  $\Psi$  is an isomorphism with its image  $C = \Psi(X(G_{m,n}))$ .  $C$  is a curve consisting of  $\frac{(n-1)(m-1)}{2} + 1$  irreducible components, all of them smooth and isomorphic to  $\mathbb{C}$ . They intersect with nodal normal crossing singularities following the pattern in Figure 1.*

*Proof.* Let us look first at  $\Psi_0 = \Psi|_{X_{red}} : X_{red} \rightarrow \mathbb{C}^3$ . For a given  $\rho = (A, B) \in X_{red}$ , with the shape given in Proposition 2.1, we have that

$$\Psi_0 : s = t + t^{-1} \mapsto (t^n + t^{-n}, t^m + t^{-m}, t^{n+m} + t^{-(n+m)}).$$

This map is clearly injective: the image recovers

$$\{t^n, t^{-n}\}, \{t^m, t^{-m}\}, \{t^{n+m}, t^{-(n+m)}\}.$$

From this, we recover  $\{(t^n, t^m), (t^{-n}, t^{-m})\}$  and hence the pair  $t, t^{-1}$  (since  $n, m$  are coprime).

Let us see that  $\Psi_0$  is an immersion. The differential is

$$\frac{d\Psi_0}{dt} = (nt^{-n-1}(t^{2n} - 1), mt^{-m-1}(t^{2m} - 1), (n+m)t^{-n-m-1}(t^{2n+2m} - 1)). \quad (3)$$

This is non-zero at all  $t \neq \pm 1$ . As  $\frac{ds}{dt} \neq 0$ , we have  $\frac{d\Psi_0}{ds} \neq (0, 0, 0)$ . For  $t = \pm 1$ , we note that  $\frac{ds}{dt} = t^{-2}(t^2 - 1)$ , so

$$\frac{d\Psi_0}{ds} = \left( nt^{-n+1} \frac{t^{2n} - 1}{t^2 - 1}, mt^{-m+1} \frac{t^{2m} - 1}{t^2 - 1}, (n+m)t^{-n-m+1} \frac{t^{2n+2m} - 1}{t^2 - 1} \right),$$

which is non-zero again.

Now, consider a component of  $X_{irr}$  corresponding to a pair  $(\lambda, \mu)$ . Take  $r \in \mathbb{C}$ . Fix the basis  $\{e_1, e_2\}$  of  $\mathbb{C}^2$  which is given as the eigenbasis of  $A$ . Let  $\{f_1, f_2\}$  be the eigenbasis of  $B$ . As the double ratio  $(0 : \infty : 1 : r/(r-1)) = r$ , we can take  $f_1 = (1, 1)$  and  $f_2 = (r-1, r)$ . This corresponds to the matrices:

$$\begin{aligned} A &= \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \\ B &= \begin{pmatrix} 1 & r-1 \\ 1 & r \end{pmatrix} \begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix} \begin{pmatrix} 1 & r-1 \\ 1 & r \end{pmatrix}^{-1} \\ &= \begin{pmatrix} r(\mu - \mu^{-1}) + \mu^{-1} & (1-r)(\mu - \mu^{-1}) \\ r(\mu - \mu^{-1}) & \mu - r(\mu - \mu^{-1}) \end{pmatrix}. \end{aligned}$$

Therefore:

$$\begin{aligned} \Psi(A, B) &= (\text{tr}(A), \text{tr}(B), \text{tr}(AB)) \\ &= (\lambda + \lambda^{-1}, \mu^{-1} + \mu, (\lambda\mu^{-1} + \lambda^{-1}\mu) + r(\lambda - \lambda^{-1})(\mu - \mu^{-1})). \end{aligned}$$

The image of this component is a line in  $\mathbb{C}^3$ . Its direction vector is  $(0, 0, 1)$ . At an intersection point with  $\Psi_0(X_{red})$ , the tangent vector to  $\Psi_0(X_{red})$ , given in (3), has non-zero first and second component, since  $\lambda = t^n, \mu = t^m$  and  $t \neq 0, \lambda^2 \neq 1, \mu^2 \neq 1$ . So the intersection of these components is a transverse nodal singularity.

Finally, note that the map  $\Psi : X(G_{m,n}) \rightarrow C$  is an algebraic map, it is a bijection, and  $C$  is a nodal curve (the mildest possible type of singularities). Therefore  $\Psi$  must be an isomorphism.  $\square$



**Corollary 3.2.**  $M(G) \cong X(G)$ , for  $G = G_{m,n}$ .

*Proof.* By Lemma 1.1,  $\varphi : M(G) \rightarrow X(G)$  is an algebraic map which is a bijection. As the singularities of  $X(G)$  are just transverse nodes,  $\varphi$  must be an isomorphism.  $\square$

**Acknowledgement.** The author wishes to thank the referee for useful comments, specially for pointing out the reference [2].

## References

- [1] M. Culler and P. B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. (2) **117** (1983), no. 1, 109–146.
- [2] J. Martín-Morales and A-M. Oller-Marcén, *On the varieties of representations and characters of a family of one-relator subgroups*, arXiv:0805.4716.
- [3] A-M. Oller-Marcén, *The  $SL(2, \mathbb{C})$  character variety of a class of torus knots*, Extracta Math. **23** (2008), no. 2, 163–172.
- [4] D. Rolfsen, *Knots and links*, Mathematics Lecture Series, vol. 7, Publish or Perish Inc., Houston, TX, 1990. Corrected reprint of the 1976 original.