

On Singular Cut-and-Pastes in the 3-Space with Applications to Link Theory

Fujitsugu HOSOKAWA and Shin'ichi SUZUKI

ABSTRACT. In the study of surfaces in 3-manifolds, the so-called “cut-and-paste” of surfaces is frequently used. In this paper, we generalize this method, in a sense, to singular-surfaces, and as an application, we prove that two collections of singular-disks in the 3-space R^3 which span the same trivial link are link-homotopic in the upper-half 4-space $R^3[0, \infty)$ keeping the link fixed.

Throughout the paper, we work in the piecewise linear category, consisting of simplicial complexes and piecewise linear maps.

1. SINGULAR LOOPS IN A 2-CELL

We denote by ∂X and ${}^{\circ}X$, respectively, the boundary and the interior of a manifold X . For a subcomplex P in a complex M , by $N(P; M)$ we denote a regular neighborhood of P in M , that is, we construct the second derived of M and take the closed star of P , see [H], [RS].

We shall say that a submanifold X of a manifold Y is *proper* iff $X \cap \partial Y = \partial X$.

By R^n , D^n and S^{n-1} we shall denote the Euclidean n -space, the standard n -cell and the standard $(n-1)$ -sphere ∂D^n , respectively.

1.1. Definition. (1) Let $f : D^1 \rightarrow M$ and $g : S^1 \rightarrow M$ be non-degenerate continuous maps into a manifold M . Then, the images $f(D^1) = A$ and $g(S^1) = J$ will be called a *singular-arc* (or simply an arc) and a *singular-loop* (or simply a loop), respectively. In particular, A and J will be called a *simple arc* and a *simple loop*, respectively, if f and g are embeddings. The boundary of an arc $f(D^1) = A$ is the image $f(\partial D^1)$ of the boundary ∂D^1 , and we denote it by $\partial^* A$.

(2) An arc A in a manifold M is said to be *proper* iff $A \cap \partial F = \partial^* A$. A loop J in a manifold M is said to be *proper* iff $J \subset {}^o F$.

(3) Let $B = B_1 \cup \dots \cup B_n$ be a finite union of proper arcs and proper loops in a 2-manifold F^2 . A point p in B is said to be a *singular-point* of multiplicity k , iff the number of the preimages of p is k with $k \geq 2$.

We shall say that B is *normal*, iff

(i) B has only a finite number of singular-points of multiplicity 2, and

(ii) at every singular point of B , B crosses transversally.

1.2. Lemma. Let $\mathcal{J}_1 = J_{11} \cup \dots \cup J_{1m(1)}$ and $\mathcal{J}_2 = J_{21} \cup \dots \cup J_{2m(2)}$ be finite unions of proper loops in a simply connected 2-manifold F^2 such that $\mathcal{J}_1 \cap \mathcal{J}_2 = \emptyset$. Then, there exists $j \in \{1, \dots, m(1)\}$ or $k \in \{1, \dots, m(2)\}$ so that J_{1j} is contractible in $F^2 - \mathcal{J}_2$ or J_{2k} is contractible in $F^2 - \mathcal{J}_1$.

Proof. We may assume that $\mathcal{J}_1 \cup \mathcal{J}_2$ is polygonal and normal.

Let $\mathcal{R} = \{R_1, \dots, R_r\}$ be the set of regions of $F^2 - {}^o N(\mathcal{J}_1; F^2)$. It will be noticed that $R_1 \cup \dots \cup R_r \supset \mathcal{J}_2$.

If there exist a loop, say J_{2k} , of \mathcal{J}_2 , and a simply connected region, say R_h , of \mathcal{R} with $J_{2k} \subset R_h$, then J_{2k} is contractible in $R_h \subset F^2 - \mathcal{J}_1$, and so the proof is complete.

So, we may assume that there exist some non-simply connected regions, say Q_1, \dots, Q_q , of \mathcal{R} , so that $Q_1 \cup \dots \cup Q_q \supset \mathcal{J}_2$. Let $C_1 \cup \dots \cup C_s = \partial Q_1 \cup \dots \cup \partial Q_q$ be the disjoint union of simple loops on F^2 , and let Δ_h be the 2-cell on F^2 with $\partial \Delta_h = C_h (h = 1, \dots, s)$. We choose an innermost 2-cell, say Δ_1 , in $\{\Delta_1, \dots, \Delta_n\}$, i.e. there is no other Δ_h in Δ_1 . Since Δ_1 is not belong to \mathcal{R} and $C_1 = \partial \Delta_1$ is the one of the boundary curves $\partial Q_1 \cup \dots \cup \partial Q_q$, it holds that $\Delta_1 \cap \mathcal{J}_1 \neq \emptyset$, and since Δ_1 does not contain any Q_1, \dots, Q_q and $\mathcal{J}_2 \subset Q_1 \cup \dots \cup Q_q$, it holds that $\Delta_1 \cap \mathcal{J}_2 = \emptyset$. Now, any J_{1j} of \mathcal{J}_1 with $J_{1j} \cap \Delta_1 \neq \emptyset$ is contractible in $\Delta_1 \subset F^2 - \mathcal{J}_2$, and so the proof is complete. ■

In the same way as that Lemma 1.2. we have the following:

1.3. Theorem. *Let $\mathcal{J}_i = J_1 \cup \dots \cup J_{im(i)}$ be a finite union of proper loops in a simply connected 2-manifold F^2 for $i = 1, \dots, \mu$, such that $\mathcal{J}_i \cap \mathcal{J}_h = \emptyset$ for $i \neq h$. Then, there exist $j \in \{1, \dots, \mu\}$ and $k \in \{1, \dots, m(j)\}$ so that J_{jk} is contractible in $F^2 - \bigcup_{i \neq j} \mathcal{J}_i$.*

Proof. We prove this by induction on the number μ of the classes \mathcal{J}_i . The case of $\mu = 1$ is trivial, and the case $\mu = 2$ is Lemma 1.2. So, we assume that $\mu \geq 3$ and Theorem is true for $\mu - 1$. We may assume that every \mathcal{J}_i is polygonal and normal.

Let $\mathcal{R} = \{R_1, \dots, R_r\}$ be the set of regions of $F^2 - \mathcal{N}(\mathcal{J}_1; F^2)$. It will be noted that $R_1 \cup \dots \cup R_r \supset \mathcal{J}_2 \cup \dots \cup \mathcal{J}_\mu$.

If there exist a loop, say J_{jk} , of \mathcal{J}_j and a simply connected region, say R_h , of \mathcal{R} with $J_{jk} \subset R_h$, then $\mathcal{J}'_i = \mathcal{J}_i \cap R_h (i = 2, \dots, \mu)$ is a finite union of loops in the simply connected region R_h satisfying the conditions of Theorem. By induction hypothesis, we have a loop, say J_{jk} , of $\mathcal{J}'_j \subset \mathcal{J}_j$ so that J_{jk} is contractible in $R_h - \bigcup_{i \neq 1, j} \mathcal{J}'_i \subset F^2 - \bigcup_{i \neq j} \mathcal{J}_i$, and so the proof is complete.

So, we may assume that there exist some non-simply connected regions, say Q_1, \dots, Q_q of \mathcal{R} , so that $Q_1 \cup \dots \cup Q_q \supset \mathcal{J}_2 \cup \dots \cup \mathcal{J}_\mu$. Now, the proof of this case, which is omitted here, is the same as that of Lemma 1.2. ■

In general, we have the following:

1.4. Theorem. Let $\mathcal{A}_i = A_{i1} \cup \cdots \cup A_{im(i)}$ be a finite union of proper arcs in a simply connected 2-manifold F^2 for $i = 1, \dots, \mu$, and let $\mathcal{J}_i = J_{i1} \cup \cdots \cup J_{im(i)}$ be a finite union of proper loops in F^2 , such that $(\mathcal{A}_i \cup \mathcal{J}_i) \cap (\mathcal{A}_h \cup \mathcal{J}_h) = \emptyset$ for $i \neq h$. Then, there exist $j \in \{1, \dots, \mu\}$ and $k \in \{1, \dots, m(j)\}$ so that J_{jk} is contractible in $F^2 - \bigcup_{i \neq j} (\mathcal{A}_i \cup \mathcal{J}_i)$.

Proof. We may assume that every $\mathcal{A}_i \cup \mathcal{J}_i$ is polygonal and normal.

Since every region of $F^2 - {}^{\circ}N(\mathcal{A}_i; F^2)$ is simply connected, the proof of Theorem is similar to that of Theorem 1.3, and so it is omitted here. ■

2. SINGULAR SPHERES IN A 3-CELL

In this section, we will discuss singular 2-spheres in a 3-cell and prove similar theorems to those in the previous section.

First let us explain several well-known facts to be used in the sequel.

If a compact 3-manifold M is embeddable in the 3-sphere S^3 , then there is a 1-complex G in S^3 such that the exterior $S^3 - {}^{\circ}N(G; S^3)$ is homeomorphic to M by Fox [F].

A 1-complex G in S^3 is said to be split, iff there exists a 2-sphere $S \subset S^3 - G$, such that both components of $S^3 - S$ contain points of G . If a 1-complex $G \subset S^3$ is not split, then the exterior $S^3 - {}^{\circ}N(G; S^3)$ is aspherical, i.e. the second homotopy group $\pi_2(S^3 - {}^{\circ}N(G; S^3)) = \{0\}$, by Papakyriakopoulos [P]. In particular, if $G \subset S^3$ is a connected 1-complex, then $S^3 - {}^{\circ}N(G; S^3)$ is aspherical.

We will call a compact 3-manifold M an *aspherical region*, iff M is embeddable in S^3 and aspherical.

It holds the following:

2.1. Proposition. (i) If a compact 3-manifold M is embeddable in S^3 and ∂M is connected, then M is an aspherical region.

(2) Let M be an aspherical region with connected boundary ∂M and let $F \subset {}^{\circ}M$ be a closed connected 2-manifold. Then, there exists an aspherical region R in M with $\partial R = F$. ■

The following corresponds to Definition 1.1.

2.2. Definition. (1) Let $f : F^2 \rightarrow M$ be a non-degenerate continuous map of a compact 2-manifold F^2 into a manifold M . Then, the image $f(F^2) = F$ will be called a singular-surface. In particular, singular-surfaces $f(D^2) = D$ and $g(S^2) = S$ will be called a singular-disk and singular-sphere, respectively.

The boundary of a singular-surface $f(F^2) = F$ is the image $f(\partial F^2)$, and we denote it by $\partial^* F$.

(2) A singular-surface F in a manifold M is said to be proper iff $F \cap \partial M = \partial^* F$.

(3) Let F be a proper singular-surface in a 3-manifold M . A point p in F is a singular-point of multiplicity k , iff the number of the preimages of p is k with $k \geq 2$.

We shall say that F is normal, iff

(i) F has only singular-points of multiplicity 2 and 3,

(ii) the set of singular-points of multiplicity 2 is a finite number of polygonal curves, that is, singular-arcs and singular-loops, which will be called double-lines,

(iii) the set of singular-points of multiplicity 3 consists of a finite number of points which are intersection points of the double-lines, which will be called triple-points, and

(iv) at every singular-point of multiplicity 2, F crosses transversally.

In fact, every singular-point p of F has one of the neighborhood described in Figure 1, and it is well known that every singular-surface may be ε -approximated by such a normal one.




Figure 1

2.3. Lemma. *Let $\mathcal{S}_1 = S_{11} \cup \cdots \cup S_{1m(1)}$ and $\mathcal{S}_2 = S_{21} \cup \cdots \cup S_{2m(2)}$ be finite unions of proper singular-spheres in an aspherical region M with connected boundary ∂M such that $\mathcal{S}_1 \cap \mathcal{S}_2 = \emptyset$. Then, there exists $j \in \{1, \dots, m(1)\}$ or $k \in \{1, \dots, m(2)\}$ so that S_{1j} is contractible in $M - \mathcal{S}_2$ or S_{2k} is contractible in $M - \mathcal{S}_1$.*

Proof. We may assume that $\mathcal{S}_1 \cup \mathcal{S}_2$ is normal. The proof of this Lemma is similar to that of Lemma 1.2.

Let $\mathcal{R} = \{R_1, \dots, R_r\}$ be the set of regions of $M - \circ N(\mathcal{S}_1; M)$. It will be noted that $R_1 \cup \cdots \cup R_r \supset \mathcal{S}_2$.

If there exist a singular-sphere, say S_{2k} , of \mathcal{S}_2 and an aspherical region, say R_h , of \mathcal{R} with $S_{2k} \subset R_h$, then S_{2k} is contractible in $R_h \subset M - \mathcal{S}_1$, and we are finished.

So, we may assume that there exist some spherical regions, say Q_1, \dots, Q_q , in \mathcal{R} , so that $Q_1 \cup \cdots \cup Q_q \supset \mathcal{S}_2$. Let $F_1 \cup \cdots \cup F_s = \partial Q_1 \cup \cdots \cup \partial Q_q$ be the disjoint union of closed connected 2-manifolds, and let M_h be the aspherical region in M with $\partial M_h = F_h$ for $h = 1, \dots, s$, see Proposition 2.1 (2). We choose an innermost region, say M_1 , in these aspherical regions, that is, there are no other M_h in M_1 . Then, by the same way as the proof of Lemma 1.2, it is easily checked that $M_1 \cap \mathcal{S}_1 \neq \emptyset$ and $M_1 \cap \mathcal{S}_2 = \emptyset$. Now, any S_{1j} of \mathcal{S}_1 with $S_{1j} \cap M_1 \neq \emptyset$ is contractible in $M_1 \subset M - \mathcal{S}_2$, and completing the proof. ■

The following theorems correspond to Theorems 1.3 and 1.4, respectively.

2.4. Theorem. *Let $S_i = S_{i1} \cup \dots \cup S_{im(i)}$ be a finite union of proper singular-spheres in an aspherical region M with connected boundary ∂M for $i = 1, \dots, \mu$, such that $S_i \cap S_h = \emptyset$ for $i \neq h$. Then, there exist $j \in \{1, \dots, \mu\}$ and $k \in \{1, \dots, m(j)\}$ so that S_{jk} is contractible in $M - \bigcup_{i \neq j} S_i$.*

Proof. The proof is similar to that of Lemma 2.3, and is word for word that of Theorem 1.3. ■

2.5. Theorem. *Let M be an aspherical region with connected boundary ∂M . Let $\mathcal{D}_i = D_{i1} \cup \dots \cup D_{in(i)}$ and $S_i = S_{i1} \cup \dots \cup S_{im(i)}$ be finite unions of proper singular-disks and proper singular-spheres in M , respectively, for $i = 1, \dots, \mu$, such that $(\mathcal{D}_i \cup S_i) \cap (\mathcal{D}_h \cup S_h) = \emptyset$ for $i \neq h$. Then, there exist $j \in \{1, \dots, \mu\}$ and $k \in \{1, \dots, m(j)\}$, so that S_{jk} is contractible in $M - \bigcup_{i \neq j} (\mathcal{D}_i \cup S_i)$.*

Proof. We may assume that every $\mathcal{D}_i \cup S_i$ is normal. Since every region R of $M - \circ N(\mathcal{D}_i; M)$ is an aspherical region, the proof of this Theorem is similar to that of Theorem 2.4, and is word for word that of Theorem 1.4. ■

3. SINGULAR CUT-AND-PASTES

3.1. Definition. *Let M^3 be a 3-manifold, and let E^2 be a compact 2-manifold in $\circ M^3$. Let $f : F^2 \rightarrow M^3$ be a non-degenerate continuous map of a compact 2-manifold F^2 into M^3 such that*

- (i) $f(F^2) = F$ is a normal singular-surface,
- (ii) F intersects with E^2 transversally, and
- (iii) no triple-point and no branch point of F lie on E^2 .

Then, the intersection $F \cap E^2$ consists of a finite number of arcs and loops. Let J be a loop in $F \cap E^2$, and let J^ be the preimage of J in F^2 : J^* is a simple loop. We suppose that J^* is 2-sided on F^2 , and let F'^2 be the 2-manifold obtained from F^2 by attaching a 2-handle along J^* . In fact, we define F'^2 as follows: We take a homeomorphism*

$h^2 : \partial D^2 \times D^1 \rightarrow N(J^*; F^2)$ with $h^2(\partial D^2 \times 0) = J^*$, and let $F'^2 = (F^2 - {}^oN(J^*; F^2)) \cup h^2(D^2 \times \partial D^1)$.

Now, we suppose that J is contractible on E^2 . Then, we have a non-degenerate continuous map, say g , of D^2 into $E^2 \subset M^3$ such that $g(\partial D^2) = J$. Using the product structure $N(E^2; M^3) \cong E^2 \times D^1$, we define a non-degenerate continuous map $f' : F'^2 \rightarrow M^3$ as follows:

$$f'|_{F'^2 - h^2(D^2 \times \partial D^1)} = f|_{F^2 - h^2(\partial D^2 \times D^1)},$$

$$f'(h^2(D^2 \times \partial D^1)) = g(D^2) \times \partial D^1 \subset E^2 \times D^1.$$

We say that $F' = f'(F'^2)$ is obtained from $F = f(F^2)$ by a cut-and-paste along $J \subset E^2$, and we denote simply by $F \rightarrow F'$.

It will be noticed that $F' \cap E^2 = F \cap E^2 - J$ and that $F'^2 = D^2 \amalg S^2$ (a disjoint union) provided that $F^2 = D^2$ and $F'^2 = S^2 \amalg S^2$ provided that $F^2 = S^2$.

3.2. Theorem. Let $\mathcal{O}_i = O_{i1} \cup \dots \cup O_{in(i)}$ be a trivial link in the 3-sphere S^3 (or the 3-space R^3) for $i = 1, \dots, \mu$, such that $\mathcal{O}_1 \cup \dots \cup \mathcal{O}_\mu$ is also a trivial link. Let $\mathcal{D}_i = D_{i1} \cup \dots \cup D_{in(i)}$ be a finite union of normal singular-disks in S^3 for $i = 1, \dots, \mu$, such that $\partial^* D_{ij} = O_{ij}$ for $i = 1, \dots, \mu$ and $j = 1, \dots, n(i)$, and $\mathcal{D}_i \cap \mathcal{D}_h = \emptyset$ for $i \neq h$.

Let $\mathcal{D}_i^* = D_{i1}^* \cup \dots \cup D_{in(i)}^*$ be mutually disjoint 2-cells in S^3 (or R^3) for $i = 1, \dots, \mu$, such that $\partial D_{ij}^* = O_{ij}$ for $i = 1, \dots, \mu$ and $j = 1, \dots, n(i)$, and $\mathcal{D}_i^* \cap \mathcal{D}_h^* = \emptyset$ for $i \neq h$.

We suppose that $\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu$ intersects with $\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*$ transversally, and any triple-point and any branch-point of $\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu$ do not lie on $\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*$.

Then, there exists a finite sequence of cut-and-pastes

$$\begin{aligned} \mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu &= \mathcal{D}_1^{(0)} \cup \dots \cup \mathcal{D}_\mu^{(0)} \rightarrow \mathcal{D}_1^{(1)} \cup \dots \cup \mathcal{D}_\mu^{(1)} \rightarrow \dots \\ &\rightarrow \mathcal{D}_1^{(u)} \cup \dots \cup \mathcal{D}_\mu^{(u)} \rightarrow \dots \rightarrow \mathcal{D}_1^{(w)} \cup \dots \cup \mathcal{D}_\mu^{(w)} \end{aligned}$$

along $(\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu) \cap (\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*) \subset \mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*$ such that

(1) $\mathcal{D}_i^{(u)} = D_{i1}^{(u)} \cup \dots \cup D_{in(i)}^{(u)} \cup S_{i1}^{(u)} \cup \dots \cup S_{im(i)}^{(u)}$, where $D_{ij}^{(u)}$ is a singular-disk with $\partial^* D_{ij}^{(u)} = O_{ij}$ and $S_{is}^{(u)}$ is a singular-sphere, for $i = 1, \dots, \mu$; $j = 1, \dots, n(i)$; $u = 1, \dots, w$; $s = 1, \dots, m(i)$,

(2) $\mathcal{D}_i^{(u)} \cap \mathcal{D}_h^{(u)} = \emptyset$ for $i \neq h$, $u = 1, \dots, w$, and

(3) $\mathcal{D}_i^{(w)} \cap \mathcal{D}_h^* = \emptyset$ for $i \neq h$, and $\mathcal{D}_i^{(w)} \cap \mathcal{D}_i^* = (D_{i1}^{(w)} \cup \dots \cup D_{in(i)}^{(w)}) \cap \mathcal{D}_i^*$ consists of a finite number of proper arcs in \mathcal{D}_i^* .

Proof. From our hypothesis, $D_{ij} \cap D_{hk}^*$ consists of proper loops in D_{hk}^* provided that $i \neq h$, and $D_{ij} \cap D_{ik}^*$ consists of proper loops and proper arcs in D_{ik}^* for every i, j, k . Therefore, by the induction on the number $n = n(1) + \dots + n(\mu)$ of 2-cells in $\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*$, it suffices to show that there exists a finite sequence of cut-and-pastes of $\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu$ along proper loops $(\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu) \cap D_{11}^* \subset D_{11}^*$ so that $\mathcal{D}_1^{(u)} \cup \dots \cup \mathcal{D}_\mu^{(u)}$ satisfies the conditions (1), (2) and

(3) $\mathcal{D}_i^{(w)} \cap D_{11}^* = \emptyset$ and $\mathcal{D}_i^{(w)} \cap D_{1j}^* = \mathcal{D}_i \cap D_{1j}^*$ for $i = 2, \dots, t$ and $j = 2, \dots, n(1)$, and $\mathcal{D}_1^{(w)} \cap D_{11}^*$ consists of a finite number of proper arcs in D_{11}^* and $\mathcal{D}_1^{(w)} \cap D_{1j}^* = \mathcal{D}_1 \cap D_{1j}^*$ for $j = 2, \dots, n(1)$.

We consider $\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu$ and D_{11}^* . Let $\mathcal{A}_1 = A_{11} \cup \dots \cup A_{1a(1)}$ be the collection of proper arcs in $\mathcal{D}_1 \cap D_{11}^*$ on D_{11}^* , and let $\mathcal{A}_i = \emptyset$ be the collection of proper arcs in $\mathcal{D}_i \cap D_{11}^*$ for $i = 2, \dots, \mu$. Let $\mathcal{J}_i = J_{i1} \cup \dots \cup J_{ib(i)}$ be a collection of proper loops in $\mathcal{D}_i \cap D_{11}^*$ on D_{11}^* for $i = 1, \dots, \mu$. Then, $\mathcal{A}_i \cup \mathcal{J}_i$ satisfies the assumptions in Theorem 1.4, and so there exists a loop J_{jk} of some \mathcal{J}_j such that J_{jk} is contractible in $D_{11}^* - \bigcup_{i \neq j} (\mathcal{A}_i \cup \mathcal{J}_i)$. We have a non-degenerate continuous map $g : D^2 \rightarrow$

D_{11}^* such that $g(D^2) \cap (\mathcal{A}_i \cup \mathcal{J}_i) = \emptyset$ for $i \neq j$. Using this g , we perform the first cut-and-paste for $\mathcal{D}_j \subset \mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu = \mathcal{D}_1^{(0)} \cup \dots \cup \mathcal{D}_\mu^{(0)}$ and obtain $\mathcal{D}_1^{(1)} \cup \dots \cup \mathcal{D}_\mu^{(1)}$. Let w be the number of loops in $(\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu) \cap D_{11}^*$. By the repetition of the procedure w times, we can get rid of all loops in $(\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu) \cap D_{11}^*$, and it is easily checked that $\mathcal{D}_1^{(u)} \cup \dots \cup \mathcal{D}_\mu^{(u)}$ satisfies the required conditions for $u = 1, \dots, w$, and we complete the proof of Theorem. ■

3.3. Remarks. (1) From the proof of Theorem 3.2, we know that w is the number of loops in $(\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu) \cap (\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*)$

and $w = m(1) + \dots + m(\mu)$, which is the number of singular-spheres in $\mathcal{D}_1^{(w)} \cup \dots \cup \mathcal{D}_\mu^{(w)}$.

(2) Let D and D^* be a normal singular-disk and a 2-cell, respectively, in S^3 (or R^3) such that $\partial^*D = \partial D^* = O$ (a trivial knot). Let A be a proper arc of $D \cap D^*$ in D^* and let α be a simple arc in O with $\partial\alpha = \partial^*A$. Since $A \cup \alpha$ is contractible in D^* , we can formulate a cut-and-paste of D along $A \cup \alpha \subset D^*$ as the same way as Definition 3.1 except for obvious modifications, so that $D \rightarrow D' = D'_1 \cup S'_1$, where S'_1 is a singular-sphere and D'_1 is a singular-disk with $\partial^*D'_1 = O$.

Now, in the notation and assumptions of Theorem 3.2, we suppose that $D_{ij} \cap D_{ik}^*$ does not contain proper arcs on D_{ik}^* for $i = 1, \dots, \mu$ and $j \neq k$. Then, we can remove proper arcs of $\mathcal{D}_i^{(w)} \cap \mathcal{D}_i^*$ by a finite sequence of the modified cut-and-pastes.

4. APPLICATIONS TO LINK THEORY

A continuous image of the 3-cell D^3 will be called a *singular-ball*. The *boundary* of a singular-ball B is the image of ∂D^3 , and we denote it by ∂^*B .

We use here the same notation as that of Section 0 in [KSS].

The following is a generalization of Horibe-Yanagawa's Lemma [KSS, Lemma 1.6] in a sense.

4.1. Theorem. *In the notation and assumptions of Theorem 3.2, let $\Sigma_i = \Sigma_{i1} \cup \dots \cup \Sigma_{in(i)}$ be a finite union of singular-spheres in $R^3[0, 1]$ defined by*

$$\Sigma_{ij} = D_{ij}[0] \cup O_{ij} \times [0, 1] \cup D_{ij}^*[1]$$

for $i = 1, \dots, \mu$ and $j = 1, \dots, n(i)$. Then, we can find a finite union of singular-balls $\mathcal{B}_i = \mathcal{B}_{i1} \cup \dots \cup \mathcal{B}_{in(i)}$ in $R^3[0, \infty)$ for $i = 1, \dots, \mu$, such that $\partial^\mathcal{B}_{ij} = \Sigma_{ij}$ for every i and j , and $\mathcal{B}_i \cap \mathcal{B}_h = \emptyset$ for $i \neq h$.*

Proof. The proof is similar to that of [KSS, Lemma 1.6]. We shall construct the required singular-balls $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu$ by specifying the cross-sections $\mathcal{B}_{ij} \cap R^3[t]$ for all i and j .

Under the notation of Theorem 3.2, we also use Theorem 3.2. Let $g_u : D^2 \rightarrow \mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*$ ($u = 1, \dots, w$) be a non-degenerate continuous map such that we perform the u -th cut-and-paste

$$\mathcal{D}_1^{(u-1)} \cup \dots \cup \mathcal{D}_\mu^{(u-1)} \rightarrow \mathcal{D}_1^{(u)} \cup \dots \cup \mathcal{D}_\mu^{(u)}$$

in Theorem 3.2 along the loop $g_u(\partial D^2)$ under g_u . We extend g_u to a continuous map

$$g_u^\# : h^2(D^2 \times D^1) \rightarrow N(\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*; R^3) \cong (\mathcal{D}_1^* \cup \dots \cup \mathcal{D}_\mu^*) \times D^1$$

of the 3-cell $h^2(D^2 \times D^1)$ naturally, and we denote the singular-ball $g_u^\#(h^2(D^2 \times D^1))$ by H_u . We divide the interval $[0,1]$ into the subintervals $[0, t_1], [t_1, t_2], \dots, [t_{w-1}, t_w], [t_w, 1]$, where $t_u = u/(w + 1)$, $u = 1, \dots, w$. Let

$$\begin{aligned} (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu)[t] \text{ for } 0 \leq t < t_1 \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t_1] &= (\mathcal{D}_1 \cup \dots \cup \mathcal{D}_\mu \cup H_1)[t_1], \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{D}_1^{(1)} \cup \dots \cup \mathcal{D}_\mu^{(1)})[t] \text{ for } t_1 < t < t_2, \end{aligned}$$

.....

$$\begin{aligned} (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{D}_1^{(u-1)} \cup \dots \cup \mathcal{D}_\mu^{(u-1)})[t] \text{ for } t_{u-1} < t < t_u, \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t_u] &= (\mathcal{D}_1^{(u-1)} \cup \dots \cup \mathcal{D}_\mu^{(u-1)} \cup H_u)[t_u], \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{D}_1^{(u)} \cup \dots \cup \mathcal{D}_\mu^{(u)})[t] \text{ for } t_u < t < t_{u+1}, \end{aligned}$$

.....

$$\begin{aligned} (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t_w] &= (\mathcal{D}_1^{(w-1)} \cup \dots \cup \mathcal{D}_\mu^{(w-1)} \cup H_w)[t_w], \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{D}_1^{(w)} \cup \dots \cup \mathcal{D}_\mu^{(w)})[t] \text{ for } t_w < t \leq 1. \end{aligned}$$

Thus, we constructed $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[0,1]$ which consists of $n = n(1) + \dots + n(\mu)$ singular-balls with $w = m(1) + \dots + m(\mu)$ singular-balls removed.

Let $S_{ij}^{(w)} = D_{ij}^{(w)} \cup D_{ij}^*$ be the singular-sphere for $i = 1, \dots, \mu$ and $j = m(i)+1, \dots, m(i)+n(i)$, and let $\mathcal{S}_i = \mathcal{D}_i^{(w)} \cup \mathcal{D}_i^* = S_{i1}^{(w)} \cup \dots \cup S_{i, m(i)+n(i)}^{(w)}$, which consists of $m(i) + n(i)$ singular-spheres in R^3 . From Theorem 3.2(2) and (3), it is easy to see that $\mathcal{S}_i \cap \mathcal{S}_h = \emptyset$ for $i \neq h$, which is the assumption of Theorem 2.4.

We divide the interval $[1, 2]$ into the $n + w + 1$ subintervals $[1, s_1], [s_1, s_2], \dots, [s_{n+w-1}, s_{n+w}], [s_{n+w}, 2]$, where $s_v = 1 + v/(n + w + 1)$, $v = 1, \dots, n + w$. From now on, we construct $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[1, 2]$ so that $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[0, 2]$ forms the required singular-balls. By Theorem 2.4, there exist $j \in \{1, \dots, \mu\}$ and $k \in \{1, \dots, m(j) + n(j)\}$ so that $S_{jk}^{(w)}$ is contractible in $R^3 - \bigcup_{i \neq j} \mathcal{S}_i$. Let $g_1 : D^3 \rightarrow R^3 - \bigcup_{i \neq j} \mathcal{S}_i$ be a continuous map such that $g_1(\partial D^3) = S_{jk}^{(w)}$, and we denote $g_1(D^3)$ by E_1 . We set $\mathcal{S}_j^{(1)} = \mathcal{S}_j - S_{jk}^{(w)}$, and $\mathcal{S}_i^{(1)} = \mathcal{S}_i$ for $i \neq j$. Then, we define $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[1, s_2]$ as follows:

$$\begin{aligned} (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{S}_1 \cup \dots \cup \mathcal{S}_\mu)[t] \text{ for } 1 \leq t < s_1, \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[s_1] &= (\mathcal{S}_1 \cup \dots \cup \mathcal{S}_\mu \cup E_1)[s_1], \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{S}_1^{(1)} \cup \dots \cup \mathcal{S}_\mu^{(1)})[t] \text{ for } s_1 < t < s_2. \end{aligned}$$

By Theorem 2.4, there exist $j' \in \{1, \dots, \mu\}$ and $k' \in \{1, \dots, m(j') + n(j')\}$ so that $S_{j'k'}$ is contractible in $R^3 - \bigcup_{i \neq j'} \mathcal{S}_i^{(1)}$. Let $g_2 : D^3 \rightarrow R^3 - \bigcup_{i \neq j'} \mathcal{S}_i^{(1)}$ be a continuous map with $g_2(\partial D^3) = S_{j'k'}$, and we denote $g_2(D^3)$ by E_2 . We set $\mathcal{S}_{j'}^{(2)} = \mathcal{S}_{j'}^{(1)} - S_{j'k'}$, and $\mathcal{S}_i^{(2)} = \mathcal{S}_i^{(1)}$ for $i \neq j'$. We define $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[s_2, s_3]$ as follows:

$$\begin{aligned} (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[s_2] &= (\mathcal{S}_1^{(1)} \cup \dots \cup \mathcal{S}_\mu^{(1)} E_2)[s_2], \\ (\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] &= (\mathcal{S}_1^{(2)} \cup \dots \cup \mathcal{S}_\mu^{(2)})[t] \text{ for } s_2 < t < s_3. \end{aligned}$$

For $R^3[s_3, s_4], \dots, R^3[s_{n+w-1}, s_{n+w}], R^3[s_{n+w}, 2]$, we repeat this

process. It should be noticed that $S_1^{(n+w-1)} \cup \dots \cup S_\mu^{(n+w-1)}$ consists of a single singular-sphere and $S_1^{(n+w)} \cup \dots \cup S_\mu^{(n+w)} = \emptyset$. Therefore, $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[s_{n+w}]$ consists of a singular-ball $E_{n+w}[s_{n+w}]$, and $(\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu) \cap R^3[t] = \emptyset$ for $s_{n+w} < t < 2$.

Thus, we obtain a union of singular-balls $\mathcal{B}_i = B_{i1} \cup \dots \cup B_{in(i)}$ in $R^3[0, \infty)$ for $i = 1, \dots, \mu$ such that $\partial^* B_{ij} = \Sigma_{ij}$. From our construction, it is easily checked that $\mathcal{B}_i \cap \mathcal{B}_h = \emptyset$ for $i \neq h$, and this completes the proof of Theorem 4.1. ■

The relation of link-homotopy was introduced in classical link theory by Milnor [M], and studied higher dimensional links by Massey-Rolfsen [MR] and Koschorke [K], etc. We record a corollary to Theorem 4.1 on link-homotopy.

4.2. Definition. Let P_1, \dots, P_μ be polyhedra, and let $\mathcal{P} = P_1 \amalg \dots \amalg P_\mu$ be their disjoint union, and let X be a manifold. A continuous map $f : \mathcal{P} \rightarrow X$ is said to be a link-map, iff $f(P_i) \cap f(P_h) = \emptyset$ for $i \neq h$. Two link-maps f_0 and f_1 of \mathcal{P} into X will be called link-homotopic, iff there exists a homotopy $\{\eta_t\}_{t \in I} : \mathcal{P} \rightarrow X$ such that $\eta_0 = f_0$, $\eta_1 = f_1$, and $\eta_t(P_i) \cap \eta_t(P_h) = \emptyset$ for $i \neq h$ and each $t \in I = [0, 1]$.

4.3. Theorem. Let $\mathcal{O}_i = O_{i1} \cup \dots \cup O_{in(i)}$ be a trivial link in the 3-space $R^3 = R^3[0] \subset R^3[0, \infty)$ (or $S^3 \subset \partial D^4$) for $i = 1, \dots, \mu$, such that $\mathcal{O}_1 \cup \dots \cup \mathcal{O}_\mu$ is also a trivial link. Let $P_i = D_{i1}^2 \amalg \dots \amalg D_{in(i)}^2$ be the disjoint union of $n(i)$ 2-cells for $i = 1, \dots, \mu$, and we set $\mathcal{P} = P_1 \amalg \dots \amalg P_\mu$. Let f and e be non-degenerate link-maps of \mathcal{P} into R^3 (or S^3) such that $f(\partial D_{ij}^2) = O_{ij} = e(\partial D_{ij}^2)$ for $i = 1, \dots, \mu$ and $j = 1, \dots, n(i)$.

Then, f and e are link-homotopic in $R^3[0, \infty)$ (or D^4) keeping $\mathcal{O}_1 \cup \dots \cup \mathcal{O}_\mu$ fixed.

Proof. Let $f(D_{ij}^2) = D_{ij}$ and $\mathcal{D}_i = D_{i1} \cup \dots \cup D_{in(i)}$ for $i = 1, \dots, \mu$ and $j = 1, \dots, n(i)$. Let $g : \mathcal{P} \rightarrow R^3$ be an embedding, and let $g(D_{ij}^2) = D_{ij}^*$ and $\mathcal{D}_i^* = D_{i1}^* \cup \dots \cup D_{in(i)}^*$. In this notation, it suffices to show that f and g are link-homotopic in $R^3[0, \infty)$ keeping $\mathcal{O}_1 \cup \dots \cup \mathcal{O}_\mu$.

In the notation of Theorem 4.1, we have a finite union of singular-balls $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_\mu$, $\mathcal{B}_i = B_{i1} \cup \dots \cup B_{in(i)}$ in $R^3[0, \infty)$ such that $\mathcal{B}_i \cap \mathcal{B}_h = \emptyset$ for $i \neq h$ and $\partial^* B_{ij} = \Sigma_{ij}$. Let $b_{ij} : D^2 \times I \rightarrow R^3[0, \infty)$ be a continuous

map of the 3-cell $D^2 \times I$ such that $b_{ij}(D^2 \times I) = B_{ij}$. We may assume that $b_{ij}|_{D^2 \times 0} = f|_{D_{ij}^2}$ and $b_{ij}|_{D^2 \times 1} = g|_{D_{ij}^2}$. Then, associating with these b_{ij} , we have a link-homotopy $\{\eta_t\}_{t \in I} : \mathcal{P} \rightarrow R^3[0, \infty)$ defined by

$$\eta_t(D_{ij}^2) = b_{ij}(D^2 \times t)$$

for every $t \in I$. From the condition of the singular-balls $B_1 \cup \dots \cup B_\mu$ in Theorem 4.1, it is easily checked that this homotopy $\{\eta_t\}_{t \in I}$ between f and g satisfies our required condition, and completing the proof of Theorem 4.3. ■

References

- [F] Fox, R.H.: *On the imbedding of polyhedra in 3-space*. Ann. of Math. (2), **49** (1948), 462-470.
- [H] Hudson, J.F.P.: *Piecewise Linear Topology*. W.A. Benjamin, New York, 1969.
- [KSS] Kawauchi, A., Shibuya, T. and Suzuki, S.: *Descriptions on surfaces in four-space I*. Math. Sem. Notes Kobe Univ., **10** (1982), 75-125.
- [K] Koschorke, U.: *Higher-order homotopy invariants for higher-dimensional link maps*. Lecture Notes in Math., **1172** (1985), Springer-Verlag, 116-129.
- [MR] Massey, W.S. and Rolfsen, D.: *Homotopy classification of higher dimensional links*. Indiana Univ. Math. J., **34** (1985), 375-391.
- [M] Milnor, J.: *Link groups*. Ann. of Math. (2), **59** (1954), 177-195.
- [RS] Rourke, C.P. and Sanderson, B.J.: *Introduction to Piecewise-Linear Topology*. Ergebn. Math. u. ihrer Grenzgeb. Bd. **69**, Springer-Verlag, Berlin-Heidelberg-New York, 1972.
- [P] Papakyriakopoulos, C.D.: *On Dehn's lemma and the asphericity of knots*. Ann. of Math. (2), **66** (1957), 1-26.

Department of Mathematics
Kobe University
Nada-ku, Kobe 657, Japan

Department of Mathematics
Waseda University
Shinjuku-ku, Tokyo 169-50, Japan

Recibido: 25 de Febrero de 1994